

Higher-Spin Fermionic Gauge Fields and Their Electromagnetic Coupling

**Marc Henneaux
Gustavo Lucena Gómez
Rakibur Rahman**

Vienna, Preprint ESI 2380 (2012)

August 20, 2012

Supported by the Austrian Federal Ministry of Education, Science and Culture
Available online at <http://www.esi.ac.at>

Higher-Spin Fermionic Gauge Fields and Their Electromagnetic Coupling

Marc Henneaux, Gustavo Lucena Gómez and Rakibur Rahman

Physique Théorique et Mathématique & International Solvay Institutes
Université Libre de Bruxelles, Campus Plaine C.P. 231, B-1050 Bruxelles, Belgium

Abstract

We study the electromagnetic coupling of massless higher-spin fermions in flat space. Under the assumptions of locality and Poincaré invariance, we employ the BRST-BV cohomological methods to construct consistent parity-preserving off-shell cubic $1-s-s$ vertices. Consistency and non-triviality of the deformations not only rule out minimal coupling, but also restrict the possible number of derivatives. Our findings are in complete agreement with, but derived in a manner independent from, the light-cone-formulation results of Metsaev and the string-theory-inspired results of Sagnotti-Taronna. We prove that any gauge-algebra-preserving vertex cannot deform the gauge transformations. We also show that in a local theory, without additional dynamical higher-spin gauge fields, the non-abelian vertices are eliminated by the lack of consistent second-order deformations.

1 Introduction

Consistent interacting theories of massless higher-spin fields in flat space are difficult to construct. Severe restrictions arise from powerful no-go theorems [1, 2], which prohibit, in Minkowski space, minimal coupling to gravity, when the particle's spin $s \geq \frac{5}{2}$, as well as to electromagnetism (EM), when $s \geq \frac{3}{2}$. However, these particles may still interact through gravitational and EM multipoles. Indeed, $\mathcal{N} = 2$ SUGRA [3] allows massless gravitini to have dipole and higher-multipole couplings, but forbids a non-zero $U(1)$ charge in flat space. Gravitational and EM multipole interactions also show up, for example, as the $2 - s - s$ and $1 - s - s$ trilinear vertices constructed in [4, 5] for bosonic fields¹.

These cubic vertices are but special cases of the general form $s - s' - s''$, that involves massless fields of arbitrary spins. Metsaev's light-cone formulation [6, 7] puts restrictions on the number of derivatives in these vertices, and thereby provides a way of classifying them. For bosonic fields, while the complete list of such vertices was given in [8], Noether procedure has been employed in [9] to explicitly construct off-shell vertices, which do obey the number-of-derivative restrictions. Also, the tensionless limit of string theory gives rise to a set of cubic vertices, which are in one-to-one correspondence with the ones of Metsaev, as has been noticed by Sagnotti-Taronna in [10], where generating functions for off-shell trilinear vertices for both bosonic and fermionic fields were presented.

In this paper, we consider the coupling of a massless fermion of arbitrary spin to a $U(1)$ gauge field, in flat spacetime of dimension $D \geq 4$. Such a study is important in that fermionic fields are required by supersymmetry, which plays a crucial role in string theory, which in turn involves higher-spin fields. This fills a gap in the higher-spin literature, most of which is about bosons only (with [7, 10, 11] among the exceptions). We do not consider mixed-symmetry fields, and restrict our attention to totally symmetric Dirac fermions $\psi_{\mu_1 \dots \mu_n}$, of spin $s = n + \frac{1}{2}$. For these fields, we employ the powerful machinery of BRST-BV cohomological methods [12] to construct systematically consistent interaction vertices², with the underlying assumptions of locality, Poincaré invariance and conservation of parity, and without relying on other methods. The would-be off-shell $1 - s - s$ cubic vertices will complement their bosonic counterparts constructed in [5].

The organization of the paper is as follows. We clarify our conventions and notations, and present our main results in the next two Subsections. In Section 2, we briefly recall the BRST deformation scheme [12] for irreducible gauge theories. With this knowledge, we then move on to constructing consistent off-shell $1 - s - s$ vertices in the following three Sections. In particular, Section 3 considers the massless Rarita-Schwinger field, while

¹In a local theory, some of these vertices may not be extended beyond the cubic order [2, 4].

²The BRST-BV approach, in general, is very useful in obtaining gauge-invariant manifestly Lorentz-invariant off-shell vertices for higher-spin fields [13], as has been emphasized recently in [14].

Section 4 pertains to $s = \frac{5}{2}$, and Section 5 generalizes the results, rather straightforwardly, to arbitrary spin, $s = n + \frac{1}{2}$. In Section 6, we prove an interesting property of the vertices under study: an abelian $1 - s - s$ vertex, i.e., a $1 - s - s$ vertex that does not deform the original abelian gauge algebra, never deforms the gauge transformations. Section 7 is a comparative study of our results with those of Metsaev [7] and Sagnotti-Taronna [10], where we explicitly show their equivalence. Section 8 investigates whether there are obstructions to the existence of second-order deformations corresponding the non-abelian vertices, i.e., if they are consistent beyond the cubic order. We conclude with some remarks in Section 9. Two appendices are added to present some useful technical details, much required for the bulk of the paper.

1.1 Conventions & Notations

We work in Minkowski spacetime with mostly positive metric. The Clifford algebra is $\{\gamma^\mu, \gamma^\nu\} = +2\eta^{\mu\nu}$, and $\gamma^{\mu\dagger} = \eta^{\mu\mu}\gamma^\mu$. The Dirac adjoint is defined as $\bar{\psi}_\mu = \psi_\mu^\dagger\gamma^0$. The D -dimensional Levi-Civita tensor, $\epsilon_{\mu_1\mu_2\dots\mu_D}$, is normalized as $\epsilon_{01\dots(D-1)} = +1$. We define $\gamma^{\mu_1\dots\mu_n} = \gamma^{[\mu_1}\gamma^{\mu_2}\dots\gamma^{\mu_n]}$, where the notation $[i_1\dots i_n]$ means totally antisymmetric expression in all the indices i_1, \dots, i_n with the normalization factor $\frac{1}{n!}$. The totally symmetric expression $(i_1\dots i_n)$ has the same normalization. We use the slash notation: $\gamma^\mu Q_\mu \equiv \not{Q}$.

The curvature for the spin-1 field is its 1-curl, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, which is just the EM field strength. Its contraction with two γ -matrices, $\gamma^{\mu\nu}F_{\mu\nu}$, is denoted as \not{F} . Similarly, the curvature for the spin- $\frac{3}{2}$ field is given by the 1-curl, $\Psi_{\mu\nu} = \partial_\mu\psi_\nu - \partial_\nu\psi_\mu$. For arbitrary spin $s = n + \frac{1}{2}$, we have a totally symmetric rank- n tensor-spinor $\psi_{\mu_1\dots\mu_n}$, whose curvature is a rank- $2n$ tensor-spinor, $\Psi_{\mu_1\nu_1|\mu_2\nu_2|\dots|\mu_n\nu_n}$, defined as the n -curl,

$$\Psi_{\mu_1\nu_1|\mu_2\nu_2|\dots|\mu_n\nu_n} \equiv [\dots [[\partial_{\mu_1}\dots\partial_{\mu_n}\psi_{\nu_1\dots\nu_n} - (\mu_1 \leftrightarrow \nu_1)] - (\mu_2 \leftrightarrow \nu_2)] \dots] - (\mu_n \leftrightarrow \nu_n).$$

This is the Weinberg curvature tensor [15], and we discuss more about it in Appendix A.

More generally, the rank- n field $\psi_{\nu_1\dots\nu_n}$ can have an m -curl, for any $0 \leq m \leq n$,

$$\psi_{\mu_1\nu_1|\dots|\mu_m\nu_m|\nu_{m+1}\dots\nu_n}^{(m)} \equiv [\dots [[\partial_{\mu_1}\dots\partial_{\mu_m}\psi_{\nu_1\dots\nu_n} - (\mu_1 \leftrightarrow \nu_1)] - (\mu_2 \leftrightarrow \nu_2)] \dots] - (\mu_m \leftrightarrow \nu_m).$$

When $m = n$, this is nothing but the curvature tensor, $\psi_{\mu_1\nu_1|\dots|\mu_n\nu_n}^{(n)} = \Psi_{\mu_1\nu_1|\dots|\mu_n\nu_n}$, whereas $m = 0$ corresponds to the original field itself, $\psi_{\nu_1\dots\nu_n}^{(0)} = \psi_{\nu_1\nu_2\dots\nu_n}$.

The Fronsdal tensor for the fermionic field [16] will be denoted as $\mathcal{S}_{\mu_1\dots\mu_n}$, so that

$$\mathcal{S}_{\mu_1\dots\mu_n} = i [\not{\partial}\psi_{\mu_1\dots\mu_n} - n\partial_{(\mu_1}\psi_{\mu_2\dots\mu_n)}]. \quad (1)$$

The symbol “ \approx ” will mean off-shell equivalence of two vertices, whereas “ \sim ” will stand for equivalence in the transverse-traceless gauge (up to an overall factor).

1.2 Results

- For massless fermions, we present a cohomological proof of the well-known fact that minimal EM coupling in flat space is ruled out for $s \geq \frac{3}{2}$ [2, 7].
- We find restrictions on the possible number of derivatives in a cubic $1 - s - s$ vertex, with $s = n + \frac{1}{2}$. There are only three allowed values: $2n - 1$, $2n$, and $2n + 1$. This is in complete agreement with the results of Metsaev [7].
- The $(2n - 1)$ -derivative vertex is non-abelian – the only one that deforms the gauge symmetry. With $F^{+\mu\nu} \equiv F^{\mu\nu} + \frac{1}{2}\gamma^{\mu\nu\alpha\beta}F_{\alpha\beta}$, we find that the off-shell vertex is simply $s = \frac{3}{2} : \bar{\psi}_\mu F^{+\mu\nu} \psi_\nu, \dots, s = n + \frac{1}{2} : \bar{\psi}_{\mu_1\nu_1|\dots|\mu_{n-1}\nu_{n-1}}^{(n-1)} F^{+\mu}{}_\nu \psi^{(n-1)\mu_1\nu_1|\dots|\mu_{n-1}\nu_{n-1}} \nu$.

We see that the $(n - 1)$ -curl of the fermionic field appears in the vertex.

- The $2n$ -derivative vertex is abelian. It exists only for $D > 4$, and is gauge invariant up to a total derivative. It involves the curvature tensor, and takes the form

$$s = n + \frac{1}{2} : (\bar{\Psi}_{\mu_1\nu_1|\mu_2\nu_2|\dots|\mu_n\nu_n} \gamma^{\mu_1\nu_1\alpha_1\beta_1\lambda} \Psi_{\alpha_1\beta_1|\mu_2\nu_2|\dots|\mu_n\nu_n}) A_\lambda.$$

It can be interpreted as a Chern-Simons term. To see this, let us start with the spin- $\frac{3}{2}$ case. For any choice of spinor indices a, b , the expression $\bar{\Psi}_a \wedge \Psi^b$ defines a closed 4-form, $d(\bar{\Psi}_a \wedge \Psi^b) = 0$. Here $\Psi^b = \frac{1}{2}\Psi_{\alpha\beta}^b dx^\alpha \wedge dx^\beta$ and a similar expression holds for $\bar{\Psi}_a$. The associated Chern-Simons 5-form

$$\bar{\Psi}_a \wedge \Psi^b \wedge A = \frac{1}{4}\bar{\Psi}_{a|\mu_1\nu_1} \Psi_{\alpha_1\beta_1}^b A_\lambda dx^{\mu_1} \wedge dx^{\nu_1} \wedge dx^{\alpha_1} \wedge dx^{\beta_1} \wedge dx^\lambda,$$

with $A = A_\lambda dx^\lambda$, is therefore gauge invariant up to the exterior derivative of a 4-form. Replacing $dx^{\mu_1} \wedge dx^{\nu_1} \wedge dx^{\alpha_1} \wedge dx^{\beta_1} \wedge dx^\lambda$ by $(\gamma^{\mu_1\nu_1\alpha_1\beta_1\lambda})^a{}_b$ and summing over the spinor indices give a scalar, which is gauge invariant up to a total divergence. This understanding of the $2n$ -derivative vertex explains why it exists only in $D \geq 5$. For higher-spin fields, the curvatures are not exterior forms since they are described by mixed-symmetry Young tableaux. However, the contracted expression

$$\bar{\Psi}_a \wedge \Psi^b = \frac{1}{4}\bar{\Psi}_{a|\mu_1\nu_1|\mu_2\nu_2|\dots|\mu_n\nu_n} \Psi_{\alpha_1\beta_1}^b dx^{\mu_1} \wedge dx^{\nu_1} \wedge dx^{\alpha_1} \wedge dx^{\beta_1}$$

is a closed 4-form, and the construction proceeds then in the same way.

- The $(2n + 1)$ -derivative vertex, which is the highest-derivative one, is a 3-curvature term (Born-Infeld type).
- The non-abelian cubic vertices generically get obstructed, in a local theory, at second order deformation. In special cases, such vertices may extend beyond the cubic order, if additional dynamical higher-spin gauge fields are present.

2 The BRST Deformation Scheme

As pointed out in [12], one can reformulate the classical problem of introducing consistent interactions in a gauge theory in terms of the BRST differential and the BRST cohomology. The advantage is that the search for all possible consistent interactions becomes systematic, thanks to the cohomological approach. Obstructions to deforming a gauge-invariant action also become related to precise cohomological classes of the BRST differential. In what follows, we briefly explain the BRST deformation scheme.

Let us consider an irreducible gauge theory of a collection of fields $\{\phi^i\}$, with m gauge invariances, $\delta_\varepsilon \phi^i = R_\alpha^i \varepsilon^\alpha$, $\alpha = 1, 2, \dots, m$. Corresponding to each gauge parameter ε^α , one introduces a ghost field \mathcal{C}^α , with the same algebraic symmetries but opposite Grassmann parity (ϵ). The original fields and ghosts are collectively called fields, denoted by Φ^A . The configuration space is further enlarged by introducing, for each field and ghost, an antifield Φ_A^* , that has the same algebraic symmetries (in its indices when A is a multi-index) but opposite Grassmann parity.

In the algebra generated by the fields and antifields, we introduce two gradings: the pure ghost number (pgh) and the antighost number (agh). The former is non-zero only for the ghost fields. In particular, for irreducible gauge theories, $pgh(\mathcal{C}^\alpha) = 1$, while $pgh(\phi^i) = 0$ for any original field. The antighost number, on the other hand, is non-zero only for the antifields Φ_A^* . Explicitly, $agh(\Phi_A^*) = pgh(\Phi^A) + 1$, $agh(\Phi^A) = 0 = pgh(\Phi_A^*)$. The ghost number (gh) is another grading, defined as $gh = pgh - agh$.

On the space of fields and antifields, one defines an odd symplectic structure

$$(X, Y) \equiv \frac{\delta^R X}{\delta \Phi^A} \frac{\delta^L Y}{\delta \Phi_A^*} - \frac{\delta^R X}{\delta \Phi_A^*} \frac{\delta^L Y}{\delta \Phi^A}, \quad (2)$$

called the antibracket³. It satisfies the graded Jacobi identity.

The original gauge-invariant action $S^{(0)}[\phi^i]$ is then extended to a new action $S[\Phi^A, \Phi_A^*]$, called the master action, that includes terms involving ghosts and antifields,

$$S[\Phi^A, \Phi_A^*] = S^{(0)}[\phi^i] + \phi_i^* R_\alpha^i \mathcal{C}^\alpha + \dots, \quad (3)$$

which, by virtue of the Noether identities and the higher-order gauge structure equations, satisfies the classical master equation

$$(S, S) = 0. \quad (4)$$

In other words, the master action S incorporates compactly all the consistency conditions pertaining to the gauge transformations. This also plays role as the generator of the

³This definition gives $(\Phi^A, \Phi_B^*) = \delta_B^A$, which is real. Because a field and its antifield have opposite Grassmann parity, it follows that if Φ^A is real, Φ_B^* must be purely imaginary, and vice versa.

BRST differential \mathfrak{s} , which is defined as

$$\mathfrak{s}X \equiv (S, X). \quad (5)$$

Notice that S is BRST-closed, as a simple consequence of the master equation. From the properties of the antibracket, it also follows that \mathfrak{s} is nilpotent,

$$\mathfrak{s}^2 = 0. \quad (6)$$

Therefore, the master action S belongs to the cohomology of \mathfrak{s} in the space of local functionals of the fields, antifields, and their finite number of derivatives.

As we know, the existence of the master action S as a solution of the master equation is completely equivalent to the gauge invariance of the original action $S^{(0)}[\phi^i]$. Therefore, one can reformulate the problem of introducing consistent interactions in a gauge theory as that of deforming the solution S of the master equation. Let S be the solution of the deformed master equation, $(S, S) = 0$. This must be a deformation of the solution S_0 of the master equation of the free gauge theory, in the deformation parameter g ,

$$S = S_0 + gS_1 + g^2S_2 + \mathcal{O}(g^3). \quad (7)$$

The master equation for S splits, up to $\mathcal{O}(g^2)$, into

$$(S_0, S_0) = 0, \quad (8)$$

$$(S_0, S_1) = 0, \quad (9)$$

$$(S_1, S_1) = -2(S_0, S_2). \quad (10)$$

Eqn. (8) is fulfilled by assumption, and in fact S_0 is the generator of the BRST differential for the free theory, which we will denote as s . Thus, Eqn. (9) translates to

$$sS_1 = 0, \quad (11)$$

i.e., S_1 is BRST-closed. If the first-order local deformations are given by $S_1 = \int a$, where a is a top-form of ghost number 0, then one has the cocycle condition

$$sa + db = 0. \quad (12)$$

Non-trivial deformations therefore belong to $H^0(s|d)$ – the cohomology of the zeroth-order BRST differential s , modulo total derivative d , at ghost number 0. Now, if one makes an antighost-number expansion of the local form a , it stops at $agh = 2$ [5, 17, 18],

$$a = a_0 + a_1 + a_2, \quad agh(a_i) = i = pgh(a_i). \quad (13)$$

For cubic deformations $S_1 = \int a$, it is indeed impossible to construct an object with $agh > 2$ [5]. The result is however more general and holds in fact also for higher order deformations, as it follows from the results of [17, 18, 19].

The significance of the various terms is worth recalling. a_0 is the deformation of the Lagrangian, while a_1 and a_2 encode information about the deformations of the gauge transformations and the gauge algebra respectively [12]. Thus, if a_2 is not trivial, the algebra of the gauge transformations is deformed and becomes non-abelian. On the other hand, if $a_2 = 0$ (up to redefinitions), the algebra remains abelian to first order in the deformation parameter. In that case, if a_1 is not trivial, the gauge transformations are deformed (remaining abelian), while if $a_1 = 0$ (up to redefinitions), the gauge transformations remain the same as in the undeformed case.

The various gradings are of relevance as s decomposes into the sum of the Koszul-Tate differential, Δ , and the longitudinal derivative along the gauge orbits, Γ :

$$s = \Delta + \Gamma. \quad (14)$$

Δ implements the equations of motion (EoM) by acting only on the antifields. It decreases the antighost number by one unit while keeping unchanged the pure ghost number. Γ acts only on the original fields and produces the gauge transformations. It increases the pure ghost number by one unit without modifying the antighost number. Accordingly, all three Δ , Γ and s increase the ghost number by one unit, $gh(\Delta) = gh(\Gamma) = gh(s) = 1$. Note that Δ and Γ are nilpotent and anticommuting,

$$\Gamma^2 = \Delta^2 = 0, \quad \Gamma\Delta + \Delta\Gamma = 0. \quad (15)$$

Given the expansion (13) and the decomposition (14), the cocycle condition (12) yields the following cascade of relations, that a consistent deformation must obey

$$\Gamma a_2 = 0, \quad (16)$$

$$\Delta a_2 + \Gamma a_1 + db_1 = 0, \quad (17)$$

$$\Delta a_1 + \Gamma a_0 + db_0 = 0, \quad (18)$$

where $agh(b_i) = i$, $pgh(b_i) = i + 1$. Note that a_2 has been chosen to be Γ -closed, instead of Γ -closed modulo d , as is always possible [17].

We now analyze the conditions under which a_2 and a_1 are non-trivial.

- Non-triviality of the deformation of the gauge algebra: The highest-order term a_2 will be trivial (i.e., removable by redefinitions) if and only if one can get rid of it by adding to a an s -exact term modulo d , $sm + dn$. Expanding m and n according to the antighost number, and taking into account the fact that m and n also stop

at $agh = 2$ since they are both cubic, one finds that a_2 is trivial if and only if $a_2 = \Gamma m_2 + dn_2$. We see that the cohomology of Γ modulo d plays an important role. The cubic vertex will deform the gauge algebra if and only if a_2 is a non-trivial element of the cohomology of Γ modulo d . Otherwise, can always choose $a_2 = 0$, and $a_1 = \Gamma$ -closed [17]. Note that since a_2 is a cocycle of the cohomology of Γ modulo d , which can be chosen to be Γ -closed [17], one can investigate the general form of a_2 by studying the elements in the cohomology of Γ that are not d -exact.

- Non-triviality of the deformation of the gauge transformations: We now assume $a_2 = 0$. In this case, a_1 can be chosen to be a non-trivial cocycle of Γ . The vertex deforms the gauge transformations unless a_1 is Δ -exact modulo d , $a_1 = \Delta m_2 + dn_1$, where m_2 can be assumed to be invariant [17, 18, 19]. In that instance, one can remove a_1 , and so one can take a_0 to be Γ -closed modulo d : the vertex only deforms the action without deforming the gauge transformations. The cohomology of Δ is also relevant in that the Lagrangian deformation a_0 is Δ -closed, whereas trivial interactions are given by Δ -exact terms.

Finally, while the graded Jacobi identity for the antibracket renders (S_1, S_1) BRST-closed, the second-order consistency condition (10) requires that this actually be s -exact:

$$(S_1, S_1) = -2sS_2. \quad (19)$$

This condition determines whether or not, in a local theory, a consistent first-order deformation gets obstructed at the second order. Such obstructions are controlled by the local BRST cohomology group $H^1(s|d)$.

2.1 The Cohomology of Γ

In this Subsection we present some facts about the cohomology of Γ , which will be very useful in the latter parts of the paper. Details will be relegated to Appendix B.

The system of gauge fields under consideration consists of a photon A_μ and a rank- n spinor-tensor $\psi_{\mu_1 \dots \mu_n}$. Corresponding to them, there will be a Grassmann-odd ghost field C and a Grassmann-even rank- $(n-1)$ spinorial ghost $\xi_{\mu_1 \dots \mu_{n-1}}$, which obeys $\xi_{\mu_1 \dots \mu_{n-2}} = 0$. The set of antifields is $\Phi_A^* = \{A^{*\mu}, C^*, \bar{\psi}^{*\mu_1 \dots \mu_n}, \bar{\xi}^{*\mu_1 \dots \mu_{n-1}}\}$. We note that the cohomology of Γ is isomorphic to the space of local functions depending on

- The curvatures $\{F_{\mu\nu}, \Psi_{\mu_1\nu_1|\mu_2\nu_2|\dots|\mu_n\nu_n}\}$, and their derivatives.
- The antifields $\{A^{*\mu}, C^*, \bar{\psi}^{*\mu_1 \dots \mu_n}, \bar{\xi}^{*\mu_1 \dots \mu_{n-1}}\}$, and their derivatives.

- The undifferentiated ghosts $\{C, \xi_{\mu_1 \dots \mu_{n-1}}\}$, and the γ -traceless part of all possible curls of the spinorial ghost $\left\{ \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m}^{\nu_{m+1} \dots \nu_{n-1}}, m \leq n-1 \right\}$.
- The Fronsdal tensor $\mathcal{S}_{\mu_1 \dots \mu_n}$, and its derivatives.

The derivative of the curl, $\partial_{\nu_n} \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m}^{\nu_{m+1} \dots \nu_{n-1}}$, is of special interest. If and only if symmetrized w.r.t. the indices $\{\nu_{m+1}, \dots, \nu_n\}$, does this quantity become Γ -exact:

$$\partial_{(\nu_n} \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m}^{\nu_{m+1} \dots \nu_{n-1})} = \frac{1}{n-m} \Gamma \psi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m}^{\nu_{m+1} \dots \nu_n}, \quad 0 \leq m \leq n-1.$$

Incidentally, for the $(n-1)$ -curl one has $\partial_{\nu_n} \xi_{\mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1}}^{\nu_n} = \Gamma \psi_{\mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1}}^{\nu_n}$.

3 EM Coupling of Massless Spin 3/2

In this Section we construct parity-preserving off-shell $1 - \frac{3}{2} - \frac{3}{2}$ vertices by employing the BRST-BV cohomological methods. The spin- $\frac{3}{2}$ system is simple enough so that one can implement the BRST deformation scheme with ease, while it captures many non-trivial features that could serve as guidelines as one moves on to higher spins.

The starting point is the free theory, which contains a photon A_μ and a massless Rarita-Schwinger field ψ_μ , described by the action

$$S^{(0)}[A_\mu, \psi_\mu] = \int d^D x \left[-\frac{1}{4} F_{\mu\nu}^2 - i \bar{\psi}_\mu \gamma^{\mu\nu\rho} \partial_\nu \psi_\rho \right], \quad (20)$$

which enjoys two abelian gauge invariances:

$$\delta_\lambda A_\mu = \partial_\mu \lambda, \quad \delta_\varepsilon \psi_\mu = \partial_\mu \varepsilon. \quad (21)$$

For the Grassmann-even bosonic gauge parameter λ , we introduce the Grassmann-odd bosonic ghost C . Corresponding to the Grassmann-odd fermionic gauge parameter ε , we have the Grassmann-even fermionic ghost ξ . Therefore, the set of fields becomes

$$\Phi^A = \{A_\mu, C, \psi_\mu, \xi\}. \quad (22)$$

For each of these fields, we introduce an antifield with the same algebraic symmetries in its indices but opposite Grassmann parity. The set of antifields is

$$\Phi_A^* = \{A^{*\mu}, C^*, \bar{\psi}^{*\mu}, \bar{\xi}^*\}. \quad (23)$$

Now we construct the free master action S_0 , which is an extension of the original gauge-invariant action (20) by terms involving ghosts and antifields. Explicitly,

$$S_0 = \int d^D x \left[-\frac{1}{4} F_{\mu\nu}^2 - i \bar{\psi}_\mu \gamma^{\mu\nu\rho} \partial_\nu \psi_\rho + A^{*\mu} \partial_\mu C - (\bar{\psi}^{*\mu} \partial_\mu \xi - \partial_\mu \bar{\xi} \psi^{*\mu}) \right]. \quad (24)$$

Notice that the antifields appear as sources for the “gauge” variations, with gauge parameters replaced by corresponding ghosts. It is easy to verify that (24) indeed solves the master equation $(S_0, S_0) = 0$. The different gradings and Grassmann parity of the various fields and antifields, along with the action of Γ and Δ on them, are given in Table 1.

Table 1: Properties of the Various Fields & Antifields ($n = 1$)

Z	$\Gamma(Z)$	$\Delta(Z)$	$pgh(Z)$	$agh(Z)$	$gh(Z)$	$\epsilon(Z)$
A_μ	$\partial_\mu C$	0	0	0	0	0
C	0	0	1	0	1	1
$A^{*\mu}$	0	$-\partial_\nu F^{\mu\nu}$	0	1	-1	1
C^*	0	$-\partial_\mu A^{*\mu}$	0	2	-2	0
ψ_μ	$\partial_\mu \xi$	0	0	0	0	1
ξ	0	0	1	0	1	0
$\bar{\psi}^{*\mu}$	0	$-\frac{i}{2} \bar{\Psi}_{\alpha\beta} \gamma^{\alpha\beta\mu}$	0	1	-1	0
$\bar{\xi}^*$	0	$\partial_\mu \bar{\psi}^{*\mu}$	0	2	-2	1

For the spin- $\frac{3}{2}$ field the Fronsdal tensor is $\mathcal{S}_\mu = i[\not{\partial}\psi_\mu - \partial_\mu\psi] = -i\gamma^\nu\Psi_{\mu\nu}$, i.e., the γ -trace of the curvature. The cohomology of Γ is isomorphic to the space of functions of

- The undifferentiated ghosts $\{C, \xi\}$,
- The antifields $\{A^{*\mu}, C^*, \bar{\psi}^{*\mu}, \bar{\xi}^*\}$ and their derivatives,
- The curvatures $\{F_{\mu\nu}, \Psi_{\mu\nu}\}$ and their derivatives.

3.1 Gauge-Algebra Deformation

The next step is to consider, for the first-order deformation, the most general form of a_2 – the term with $agh = 2$, that contains information about the deformation of the gauge algebra. a_2 must satisfy $\Gamma a_2 = 0$, and be Grassmann even with $gh(a_2) = 0$. Besides, we require that a_2 be a parity-even Lorentz scalar. Then, the most general possibility is

$$a_2 = -g_0 C (\bar{\xi}^* \xi + \bar{\xi} \xi^*) - g_1 C^* \bar{\xi} \xi, \quad (25)$$

which is a linear combination of two independent terms: one contains both the bosonic ghost C and the fermionic ghost ξ , while the other contains only ξ but not C . The former one potentially gives rise to minimal coupling, while the latter could produce dipole interaction. This can be understood by first noting that the corresponding Lagrangian deformation, a_0 , is obtained through the consistency cascade (16)–(18). From the action

of Γ and Δ on the fields and antifields, it is then easy to see that the respective a_0 would contain no derivative and one derivative respectively.

3.2 Deformation of Gauge Transformations

Next, we would like to see if a_2 can be lifted to certain a_1 , i.e., with the given a_2 , if one could solve Eqn. (17) to find an a_1 . Indeed, one finds that⁴

$$\begin{aligned}\Delta a_2 &= +g_0 C [(\partial_\mu \bar{\psi}^{*\mu})\xi - \bar{\xi}(\partial_\mu \psi^{*\mu})] + g_1 (\partial_\mu A^{*\mu})\bar{\xi}\xi \\ &= -g_0 [\bar{\psi}^{*\mu}\partial_\mu(C\xi) - \partial_\mu(C\bar{\xi})\psi^{*\mu}] - g_1 A^{*\mu}\partial_\mu(\bar{\xi}\xi) + d(\dots) \\ &= -\Gamma [g_0(\bar{\psi}^{*\mu}\psi_\mu + \bar{\psi}_\mu\psi^{*\mu})C + g_0(\bar{\psi}^{*\mu}A_\mu\xi - \bar{\xi}A_\mu\psi^{*\mu}) + g_1 A^{*\mu}(\bar{\psi}_\mu\xi - \bar{\xi}\psi_\mu)] + d(\dots).\end{aligned}$$

Therefore, in view of Eqn. (17), one must have

$$a_1 = g_0 [\bar{\psi}^{*\mu}(\psi_\mu C + \xi A_\mu) + \text{h.c.}] + g_1 A^{*\mu}(\bar{\psi}_\mu\xi - \bar{\xi}\psi_\mu) + \tilde{a}_1, \quad \Gamma\tilde{a}_1 = 0, \quad (26)$$

where the ambiguity, \tilde{a}_1 , belongs to the cohomology of Γ . Its most general form will be

$$\tilde{a}_1 = [\bar{\psi}^{*\mu} X_{\mu\nu\rho} \Psi^{\nu\rho}] C + [\bar{\psi}^{*\mu} Y_{\mu\nu\rho} F^{\nu\rho} + \bar{\Psi}^{\mu\nu} Z_{\mu\nu\rho} A^{*\rho}] \xi + \text{h.c.}, \quad (27)$$

where X, Y and Z may contain derivatives and spinor indices.

3.3 Lagrangian Deformation

We note that Δa_1 must be Γ -closed modulo d , since

$$\Gamma(\Delta a_1) = \Delta(-\Gamma a_1) = \Delta[\Delta a_2 + d(\dots)] = d(\dots). \quad (28)$$

Condition (18), however, requires that Δa_1 be Γ -exact modulo d . The Δ -variation of neither of the unambiguous pieces in a_1 is Γ -exact modulo d , and the non-trivial part must be killed by $\Delta\tilde{a}_1$, if (18) holds at all. But such a cancelation is impossible for the first piece, i.e., the would-be minimal coupling, simply because \tilde{a}_1 contains too many derivatives. Thus, minimal coupling is ruled out, and we must set $g_0 = 0$. Then, we have

$$\Delta a_1 = -\Gamma(g_1 \bar{\psi}_\mu F^{\mu\nu} \psi_\nu) - \frac{1}{2}g_1 F^{\mu\nu}(\bar{\Psi}_{\mu\nu}\xi - \bar{\xi}\bar{\Psi}_{\mu\nu}) + \Delta\tilde{a}_1 + d(\dots). \quad (29)$$

The second term on the right hand side is in the cohomology of Γ , and must be canceled by $\Delta\tilde{a}_1$. To see if this is possible or not, we make use of the identity

$$\eta^{\mu\nu|\alpha\beta} \equiv \frac{1}{2}(\eta^{\mu\alpha}\eta^{\nu\beta} - \eta^{\mu\beta}\eta^{\nu\alpha}) = \frac{1}{2}\gamma^{\mu\nu}\gamma^{\alpha\beta} - 2\gamma^{[\mu}\eta^{\nu][\alpha}\gamma^{\beta]} - \frac{1}{2}\gamma^{\mu\nu\alpha\beta}, \quad (30)$$

⁴Here one also needs the relations $\Delta\xi^* = -\partial_\mu\psi^{*\mu}$, $\Gamma\bar{\psi}_\mu = -\partial_\mu\bar{\xi}$, which follow from Table 1.

to rewrite the term as

$$\begin{aligned}
F^{\mu\nu}(\bar{\Psi}_{\mu\nu}\xi - \bar{\xi}\Psi_{\mu\nu}) &= +\frac{1}{2}(\bar{\Psi}\mathcal{F} - 4\bar{\Psi}_{\mu\nu}\gamma^\mu F^{\nu\rho}\gamma_\rho)\xi - \frac{1}{2}\bar{\xi}(F\Psi - 4\gamma_\mu F^{\mu\alpha}\gamma^\beta\Psi_{\alpha\beta}) \\
&\quad -\frac{1}{2}(\bar{\Psi}_{\mu\nu}\gamma^{\mu\nu\alpha\beta}F_{\alpha\beta}\xi - \bar{\xi}F_{\mu\nu}\gamma^{\mu\nu\alpha\beta}\Psi_{\alpha\beta}) \\
&= +\frac{1}{2}(\bar{\Psi}\mathcal{F} - 4\bar{\Psi}_{\mu\nu}\gamma^\mu F^{\nu\rho}\gamma_\rho)\xi - \frac{1}{2}\bar{\xi}(F\Psi - 4\gamma_\mu F^{\mu\alpha}\gamma^\beta\Psi_{\alpha\beta}) \\
&\quad +\Gamma(\bar{\psi}_\mu\gamma^{\mu\nu\alpha\beta}F_{\alpha\beta}\psi_\nu) + d(\dots). \tag{31}
\end{aligned}$$

Notice that, we have rendered the second line in the first step Γ -exact modulo d , by virtue of the Bianchi identity, $\partial_{[\mu}F_{\nu\rho]} = 0$. We plug Eqn. (31) into (29) to obtain

$$\begin{aligned}
\Delta a_1 &= -\Gamma(g_1\bar{\psi}_\mu F^{+\mu\nu}\psi_\nu) + \Delta\tilde{a}_1 + d(\dots) \\
&\quad -\frac{1}{4}g_1[(\bar{\Psi}\mathcal{F} - 4\bar{\Psi}_{\mu\nu}\gamma^\mu F^{\nu\rho}\gamma_\rho)\xi - \bar{\xi}(F\Psi - 4\gamma_\mu F^{\mu\alpha}\gamma^\beta\Psi_{\alpha\beta})]. \tag{32}
\end{aligned}$$

Now, the most important point is that, the terms in the second line of the above expression are Δ -exact, such that it is consistent to set

$$\Delta\tilde{a}_1 = \frac{1}{4}g_1[(\bar{\Psi}\mathcal{F} - 4\bar{\Psi}_{\mu\nu}\gamma^\mu F^{\nu\rho}\gamma_\rho)\xi - \bar{\xi}(F\Psi - 4\gamma_\mu F^{\mu\alpha}\gamma^\beta\Psi_{\alpha\beta})]. \tag{33}$$

This is tantamount to setting

$$\tilde{a}_1 = ig_1\left[\bar{\psi}^{*\mu}\gamma^\nu F_{\mu\nu} - \frac{1}{2(D-2)}\bar{\psi}^*F\right]\xi + \text{h.c.}, \tag{34}$$

which, of course, is in the cohomology of Γ . Then, Eqn. (32) reduces to

$$\Delta a_1 = -\Gamma(g_1\bar{\psi}_\mu F^{+\mu\nu}\psi_\nu) + d(\dots), \tag{35}$$

so that we have a consistent Lagrangian deformation a_0 . To summarize, we have

$$a_0 = g_1\bar{\psi}_\mu F^{+\mu\nu}\psi_\nu, \quad a_1 = g_1 A^{*\mu}(\bar{\psi}_\mu\xi - \bar{\xi}\psi_\mu) + \tilde{a}_1, \quad a_2 = -g_1 C^*\bar{\xi}\xi. \tag{36}$$

3.4 Abelian Vertices

Now that we have exhausted all the possibilities for a_2 , any other vertex can only have a trivial a_2 . In this case, as we will show in Section 6, one can always choose to write a vertex as the photon field A_μ contracted with a gauge-invariant current j^μ ,

$$a_0 = j^\mu A_\mu, \quad \Gamma j^\mu = 0, \tag{37}$$

where the divergence of the current is Δ -exact:

$$\partial_\mu j^\mu = \Delta M, \quad \Gamma M = 0, \tag{38}$$

so that one has $a_1 = MC$. If, however, M happens to be Δ -exact modulo d in the space of invariants, one can add a Δ -exact term in a_0 , so that the new current is identically conserved [17]. In the latter case, the vertex does not deform the gauge symmetry at all.

Now the most general vertex of the form (37) contains the current

$$j^\lambda = \bar{\Psi}_{\mu\nu} X^{\mu\nu\alpha\beta\lambda} \Psi_{\alpha\beta}, \quad (39)$$

whose divergence is required to obey the condition (38). Here X may contain Dirac matrices as well as derivatives. It is not difficult to see if X contains more than one derivatives, a_0 is Δ -exact modulo d , i.e., trivial. First, if X contains the Laplacian, \square , the contribution is always Δ -exact, by the EoM $\square\Psi_{\mu\nu} = 0$. We can also forgo the Dirac operator, $\not{\partial}$, because by using the relation $\not{\partial}\gamma^\mu = 2\partial^\mu - \gamma^\mu\not{\partial}$, one can always make $\not{\partial}$ act on the curvature to get Δ -exact terms, thanks to the EoM $\not{\partial}\Psi_{\mu\nu} = 0$. Therefore, any derivative contained in $X^{\mu\nu\alpha\beta\lambda}$ must carry one of the five indices. Given the EoM $\partial^\mu\Psi_{\mu\nu} = 0$, the antisymmetry of the field strength $\Psi_{\mu\nu}$, and the commutativity of ordinary derivatives, the only potentially non-trivial way to have more-than-one derivatives is

$$a_0 = \left(\bar{\Psi}_{\mu\alpha} \overleftarrow{\partial}_\nu \gamma^\lambda \partial^\mu \Psi^{\alpha\nu} \right) A_\lambda. \quad (40)$$

But algebraic manipulations show that this vertex is actually Δ -exact modulo d , i.e., trivial. To see this, we use $\Psi^{\alpha\nu} = \partial^\alpha\psi^\nu - \partial^\nu\psi^\alpha$, and rewrite (40) as

$$a_0 = \left[\bar{\Psi}_{\mu\alpha} \overleftarrow{\partial}_\nu \gamma^\lambda \partial^\mu \partial^\alpha \psi^\nu - \frac{1}{2} \bar{\Psi}_{\mu\alpha} \overleftarrow{\partial}_\nu \gamma^\lambda \partial^\nu \Psi^{\mu\alpha} \right] A_\lambda.$$

While the first term is identically zero, in the second term, one can use the 3-box rule, $2\partial_\mu X \partial^\mu Y = \square(XY) - X(\square Y) - (\square X)Y$, so that

$$a_0 = -\frac{1}{4} \left[\square \left(\bar{\Psi}_{\mu\alpha} \gamma^\lambda \Psi^{\mu\alpha} \right) - \left(\square \bar{\Psi}_{\mu\alpha} \right) \gamma^\lambda \Psi^{\mu\alpha} - \bar{\Psi}_{\mu\alpha} \gamma^\lambda \left(\square \Psi^{\mu\alpha} \right) \right] A_\lambda.$$

Here, the last two terms are Δ -exact, whereas in the first term a double integration by parts gives $\square A_\lambda$, which is equal to $\partial_\lambda(\partial \cdot A)$ by the photon EoM. Then, one is left with

$$a_0 = -\frac{1}{4} \left(\bar{\Psi}_{\mu\alpha} \gamma^\lambda \Psi^{\mu\alpha} \right) \partial_\lambda(\partial \cdot A) + \Delta\text{-exact} + d(\dots).$$

Upon integrating by parts w.r.t. ∂_λ , this indeed becomes Δ -exact modulo d ,

$$a_0 = \left(\bar{\Psi}_{\mu\alpha} \overleftarrow{\partial}_\nu \gamma^\lambda \partial^\mu \Psi^{\alpha\nu} \right) A_\lambda = \Delta\text{-exact} + d(\dots). \quad (41)$$

The only possibilities are therefore that X contains either no derivative or one derivative. For the former case, we have the candidate $X^{\mu\nu\alpha\beta\lambda} = -2\eta^{\mu\nu|\alpha\beta}\gamma^\lambda$. This gives

$$M = -4i\bar{\Psi}_{\mu\nu}\partial^\mu \left(\psi^{*\nu} - \frac{1}{D-2}\gamma^\nu \not{\psi} \right) - \text{h.c.}, \quad (42)$$

which is obviously gauge invariant: $\Gamma M = 0$. However, explicit computation easily shows that M is actually Δ -exact modulo d . Therefore, one can render the current identically conserved by adding a Δ -exact term. In fact, in view of identity (30), our candidate j^μ is

$$j^\mu = \frac{1}{2} \bar{\Psi}_{\mu\nu} (\gamma^{\mu\nu\alpha\beta} \gamma^\lambda + \gamma^\lambda \gamma^{\mu\nu\alpha\beta}) \Psi_{\alpha\beta} + \Delta\text{-exact}. \quad (43)$$

Then, it is clear from the identity

$$\frac{1}{2} \gamma^{\mu\nu\alpha\beta} \gamma^\lambda + \frac{1}{2} \gamma^\lambda \gamma^{\mu\nu\alpha\beta} = \gamma^{\mu\nu\alpha\beta\lambda}, \quad (44)$$

that our 2-derivative vertex is actually off-shell equivalent (\approx) to

$$a_0 \approx (\bar{\Psi}_{\mu\nu} \gamma^{\mu\nu\alpha\beta\lambda} \Psi_{\alpha\beta}) A_\lambda. \quad (45)$$

This vertex does not deform the gauge symmetry, and is gauge invariant up to a total derivative. Note that the vertex does not exist in $D = 4$, because of the presence of $\gamma^{\mu\nu\alpha\beta\lambda}$. This is in complete agreement with Metsaev's results [7].

Finally, we are left with the possibility of having just one derivative in X , which would correspond to a 3-derivative vertex. The only candidate is $X^{\mu\nu\alpha\beta\lambda} = \frac{1}{2} \eta^{\mu\nu|\alpha\beta} \overleftrightarrow{\partial}^\lambda$, which is equivalent to $-\frac{1}{4} \gamma^{\mu\nu\alpha\beta\lambda} \overleftrightarrow{\partial}^\lambda$, up to Δ -exact terms, thanks to the identity (30). We have

$$a_0 = \frac{1}{2} (\bar{\Psi}_{\mu\nu} \eta^{\mu\nu|\alpha\beta} \overleftrightarrow{\partial}^\lambda \Psi_{\alpha\beta}) A_\lambda = \frac{1}{2} (\bar{\Psi}_{\mu\nu} \partial^\lambda \Psi^{\mu\nu} - \bar{\Psi}_{\mu\nu} \overleftarrow{\partial}^\lambda \Psi^{\mu\nu}) A_\lambda. \quad (46)$$

In this case too, our candidate current reduces on-shell to an identically conserved one, so that the vertex actually does not deform the gauge symmetry. To see this, we use the Bianchi identity $\partial^\lambda \Psi^{\mu\nu} = -\partial^\mu \Psi^{\nu\lambda} + \partial^\nu \Psi^{\mu\lambda}$, to write the vertex as

$$a_0 = \left(-\bar{\Psi}_{\mu\nu} \partial^\mu \Psi^{\nu\lambda} + \bar{\Psi}_\nu{}^\lambda \overleftarrow{\partial}_\mu \Psi^{\mu\nu} \right) A_\lambda.$$

Thanks to the EoM $\partial^\mu \Psi_{\mu\nu} = 0$, up to Δ -exact terms, the current reduces to the total derivative of a fermion bilinear, which is identically conserved:

$$a_0 \approx 2\partial_\nu (\bar{\Psi}_\alpha{}^{[\mu} \Psi^{\nu]\alpha}) A_\mu. \quad (47)$$

Upon integration by parts, this is just a 3-curvature term (Born-Infeld type),

$$a_0 \approx \bar{\Psi}_{\mu\alpha} \Psi_\nu{}^\alpha F^{\mu\nu}. \quad (48)$$

This exhausts all possible $1 - \frac{3}{2} - \frac{3}{2}$ vertices. Below we present a summary table.

Here we parenthetically comment about the nature of the abelian vertices. As it turned out, the vertices that do not deform the gauge algebra do not deform the gauge transformations either. In other words, if a_2 is trivial, so is a_1 . This is not accidental at all. In fact, in Section 6 we are going to show that, for a massless particle of arbitrary spin $s = n + \frac{1}{2}$ coupled to a $U(1)$ vector field, the cubic couplings that do not deform the gauge algebra actually do not deform the gauge transformations and hence only deform the Lagrangian.

Table 2: Summary of $1 - \frac{3}{2} - \frac{3}{2}$ Vertices

# of Derivatives	Vertex	Nature	Exists in
1	$\bar{\psi}_\mu F^{+\mu\nu} \psi_\nu$	Non-abelian	$D \geq 4$
2	$(\bar{\Psi}_{\mu\nu} \gamma^{\mu\nu\alpha\beta\lambda} \Psi_{\alpha\beta}) A_\lambda$	Abelian	$D \geq 5$
3	$\bar{\Psi}_{\mu\alpha} \Psi_{\nu}^\alpha F^{\mu\nu}$	Abelian	$D \geq 4$

4 Massless Spin 5/2 Coupled to EM

Now we move on to constructing parity-preserving off-shell cubic vertices for a spin- $\frac{5}{2}$ gauge field, which is a symmetric rank-2 tensor-spinor $\psi_{\mu\nu}$. The original free action is

$$S^{(0)}[A_\mu, \psi_{\mu\nu}] = \int d^D x \left[-\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{2} (\bar{\psi}_{\mu\nu} \mathcal{R}^{\mu\nu} - \bar{\mathcal{R}}^{\mu\nu} \psi_{\mu\nu}) \right], \quad (49)$$

where the tensor $\mathcal{R}^{\mu\nu}$ is related to the spin- $\frac{5}{2}$ Fronsdal tensor, $\mathcal{S}^{\mu\nu}$, as follows.

$$\mathcal{R}^{\mu\nu} = \mathcal{S}^{\mu\nu} - \gamma^{(\mu} \mathcal{S}^{\nu)} - \frac{1}{2} \eta^{\mu\nu} \mathcal{S}', \quad \mathcal{S}' \equiv \mathcal{S}_\mu^\mu. \quad (50)$$

Here the photon gauge invariance is as usual, while the fermionic part is gauge invariant under a constrained vector-spinor gauge parameter, ε_μ ,

$$\delta_\varepsilon \psi_{\mu\nu} = 2\partial_{(\mu} \varepsilon_{\nu)}, \quad \not{\varepsilon} = 0. \quad (51)$$

Then, the corresponding Grassmann-even fermionic ghost, ξ_μ , must also be γ -traceless:

$$\not{\xi} = 0, \quad (52)$$

and so will be its antighost. The set of fields and antifields under study are given below.

$$\Phi^A = \{A_\mu, C, \psi_{\mu\nu}, \xi_\mu\}, \quad \Phi_A^* = \{A^{*\mu}, C^*, \bar{\psi}^{*\mu\nu}, \bar{\xi}^{*\mu}\}. \quad (53)$$

The free master action, S_0 , takes the form

$$S_0 = \int d^D x \left[-\frac{1}{4} F_{\mu\nu}^2 - \frac{1}{2} (\bar{\psi}_{\mu\nu} \mathcal{R}^{\mu\nu} - \bar{\mathcal{R}}^{\mu\nu} \psi_{\mu\nu}) + A^{*\mu} \partial_\mu C - 2(\bar{\psi}^{*\mu\nu} \partial_\mu \xi_\nu - \partial_\mu \bar{\xi}_\nu \psi^{*\mu\nu}) \right]. \quad (54)$$

Properties of the various fields and antifields are given in Table 3. Note that the spin- $\frac{5}{2}$ curvature tensor is the 2-curl (see Appendix A for its properties),

$$\Psi_{\mu_1\nu_1|\mu_2\nu_2} = [\partial_{\mu_1} \partial_{\mu_2} \psi_{\nu_1\nu_2} - (\mu_1 \leftrightarrow \nu_1)] - (\mu_2 \leftrightarrow \nu_2). \quad (55)$$

The cohomology of Γ is isomorphic to the space of functions of (see Appendix B)

- The undifferentiated ghosts $\{C, \xi_\mu\}$, and the γ -traceless part of the 1-curl of the spinorial ghost $\xi_{\mu\nu}^{(1)} = 2\partial_{[\mu}\xi_{\nu]}$,
- The antifields $\{A^{*\mu}, C^*, \bar{\psi}^{*\mu\nu}, \bar{\xi}^{*\mu}\}$, and their derivatives,
- The curvatures $\{F_{\mu\nu}, \Psi_{\mu_1\nu_1|\mu_2\nu_2}\}$, and their derivatives,
- The Fronsdal tensor $\mathcal{S}_{\mu\nu}$, and its symmetrized derivatives.

Table 3: Properties of the Various Fields & Antifields ($n = 2$)

Z	$\Gamma(Z)$	$\Delta(Z)$	$pgh(Z)$	$agh(Z)$	$gh(Z)$	$\epsilon(Z)$
A_μ	$\partial_\mu C$	0	0	0	0	0
C	0	0	1	0	1	1
$A^{*\mu}$	0	$-\partial_\nu F^{\mu\nu}$	0	1	-1	1
C^*	0	$-\partial_\mu A^{*\mu}$	0	2	-2	0
$\psi_{\mu\nu}$	$2\partial_{(\mu}\xi_{\nu)}$	0	0	0	0	1
ξ_μ	0	0	1	0	1	0
$\bar{\psi}^{*\mu\nu}$	0	$\bar{\mathcal{R}}^{\mu\nu}$	0	1	-1	0
$\bar{\xi}^{*\mu}$	0	$2\partial_\nu \bar{\psi}^{*\mu\nu}$	0	2	-2	1

4.1 Non-Abelian Vertices

The set of all possible non-trivial a_2 's falls into two subsets: Subset-1 contains both the bosonic ghost C and the fermionic ghost ξ_μ , while Subset-2 contains only ξ_μ but not C .

- Subset-1 = $\left\{ C \left(\bar{\xi}_\mu^* \xi^\mu + \bar{\xi}_\mu \xi^{*\mu} \right), C \left(\bar{\xi}_{\mu\nu}^{*(1)} \xi^{(1)\mu\nu} + \bar{\xi}_{\mu\nu}^{(1)} \xi^{*(1)\mu\nu} \right) \right\}$,
- Subset-2 = $\left\{ C^* \bar{\xi}_\mu \xi^\mu, C^* \bar{\xi}_{\mu\nu}^{(1)} \xi^{(1)\mu\nu} \right\}$.

One can easily verify that other possible rearrangements of derivatives or other possible contractions of the indices, e.g., by γ -matrices, all give trivial terms, thanks to the γ -tracelessness of the fermionic ghost and its antighost. Here, the term $C \bar{\xi}_\mu^* \xi^\mu$ corresponds to potential minimal coupling, while the other candidate a_2 's to multipole interactions.

To see which of the a_2 's can be lifted to a_1 , let us solve Eqn. (17). A computation, similar to what leads one from Eqn. (25) to Eqn. (26), shows that both the elements in Subset-1 enjoy such a lift, thanks to the relations (B.9)–(B.10) among others. Explicitly,

$$a_2 = \begin{cases} C \bar{\xi}_\mu^* \xi^\mu \\ C \bar{\xi}_{\mu\nu}^{*(1)} \xi^{(1)\mu\nu} \end{cases} \longrightarrow a_1 = \begin{cases} -\bar{\psi}^{*\mu\nu} (\psi_{\mu\nu} C + 2\xi_\mu A_\nu) + \tilde{a}_1 \\ -\bar{\psi}^{*(1)\mu\nu|\rho} (\psi_{\mu\nu|\rho}^{(1)} C + 2\xi_{\mu\nu}^{(1)} A_\rho) + \tilde{a}_1, \end{cases} \quad (56)$$

and similarly for the hermitian conjugate terms. Here \tilde{a}_1 is the usual ambiguity: $\Gamma\tilde{a}_1 = 0$. To see whether these could further be lifted to a_0 's, we write

$$\Delta a_1 = \begin{cases} -\bar{\mathcal{R}}^{\mu\nu} (\psi_{\mu\nu} C + 2\xi_\mu A_\nu) + \Delta\tilde{a}_1 \\ -\bar{\mathcal{R}}^{(1)\mu\nu\|\rho} \left(\psi_{\mu\nu\|\rho}^{(1)} C + 2\xi_{\mu\nu}^{(1)} A_\rho \right) + \Delta\tilde{a}_1. \end{cases} \quad (57)$$

It is important to notice that, up to total derivatives, the Δa_1 's have an expansion in the basis of undifferentiated ghosts, $\omega_I = \{C, \xi_\mu\}$. Because $\Gamma(\Delta\tilde{a}_1) = -\Delta(\Gamma\tilde{a}_1) = 0$, the coefficients α^I in the expansion of the ambiguity will be Γ -cocycles, i.e., they will be “invariant polynomials”. Clearly, this is not the case for the unambiguous pieces; in fact, their expansion coefficients β^I are not even cocycles of $H^0(\Gamma|d)$ ⁵. Schematically,

$$\Delta a_1 = (\alpha^I + \beta^I) \omega_I + d(\dots); \quad \Gamma\alpha^I = 0, \quad \Gamma\beta^I \neq d(\dots). \quad (58)$$

Now, Γa_0 is a pg h -1 object that can be expanded, up to a total derivative, in the basis of $\{\partial_\mu C, \partial_{(\mu}\xi_{\nu)}\}$. Then, obviously, one can also expand it in the undifferentiated ghosts ω_I :

$$\Gamma a_0 = -(\partial \cdot J)^I \omega_I + d(\dots). \quad (59)$$

One can plug the respective expansions (58) and (59) for Δa_1 and Γa_0 into the consistency condition (18), and then take a functional derivative w.r.t. $\omega_I = \{C, \xi_\mu\}$ to find that

$$\alpha^I + \beta^I = \partial \cdot J^I = d(\dots). \quad (60)$$

But if this is true, then $\Gamma(\alpha^I + \beta^I) = d(\dots)$, which is in direct contradiction with the properties of α^I and β^I , given in (58)⁶. The conclusion is that none of the a_2 's in Subset-1 can be lifted all the way to a_0 . It is important to notice that this obstruction originates from the very nature of the a_2 's themselves, namely each of them contains both the ghosts.

For Subset-2, the analysis simplifies because only one term, $C^{*\bar{\xi}_{\mu\nu}^{(1)}} \xi^{(1)\mu\nu}$, with the maximum number of derivatives, can be lifted to an a_1 . For the other term we have

$$\Delta(C^{*\bar{\xi}_\nu} \xi^\nu) = A^{*\mu} (\bar{\xi}^\nu \partial_\mu \xi_\nu + \partial_\mu \bar{\xi}_\nu \xi^\nu) + d(\dots) \quad (61)$$

Because one can write $\partial_\mu \xi_\nu = \partial_{[\mu}\xi_{\nu]} + \partial_{(\mu}\xi_{\nu)}$, which is the sum of a non-trivial and a trivial element in the cohomology of Γ , the right hand side of Eqn. (61) cannot be Γ -exact modulo d . Therefore, the candidate $C^{*\bar{\xi}_\mu} \xi^\mu$ is ruled out. However, one finds that

$$\begin{aligned} \Delta(C^{*\bar{\xi}_{\mu\nu}^{(1)}} \xi^{(1)\mu\nu}) &= A^{*\rho} (\bar{\xi}^{(1)\mu\nu} \partial_\rho \xi_{\mu\nu}^{(1)} + \partial_\rho \bar{\xi}_{\mu\nu}^{(1)} \xi^{(1)\mu\nu}) + d(\dots) \\ &= \Gamma \left[A^{*\rho} \left(\bar{\psi}_{\mu\nu\|\rho}^{(1)} \xi^{(1)\mu\nu} - \bar{\xi}^{(1)\mu\nu} \psi_{\mu\nu\|\rho}^{(1)} \right) \right] + d(\dots), \end{aligned} \quad (62)$$

⁵But still, because of Eqn. (28), one must have $[\beta^I \omega_I] \in H^1(\Gamma|d)$, and indeed this is the case.

⁶For the would-be minimal coupling, the impossibility can also be seen as a consequence of α^I containing too many derivatives compared to β^I . We have used this argument for spin $\frac{3}{2}$.

thanks to the relation (B.10). Thus, indeed, $C^* \bar{\xi}_{\mu\nu}^{(1)} \xi^{(1)\mu\nu}$ gets lifted to an a_1 :

$$a_2 = C^* \bar{\xi}_{\mu\nu}^{(1)} \xi^{(1)\mu\nu} \longrightarrow a_1 = -A^{*\rho} \left(\bar{\psi}_{\mu\nu|\rho}^{(1)} \xi^{(1)\mu\nu} - \bar{\xi}^{(1)\mu\nu} \psi_{\mu\nu|\rho}^{(1)} \right) + \tilde{a}_1. \quad (63)$$

To see if this a_1 can be lifted to an a_0 , we compute its Δ variation,

$$\Delta a_1 = \Gamma \left(\bar{\psi}_{\alpha\beta|\mu}^{(1)} F^{\mu\nu} \psi^{(1)\alpha\beta|}_{\nu} \right) + \frac{1}{2} F^{\mu\nu} \left(\bar{\Psi}_{\mu\nu|\alpha\beta} \xi^{(1)\alpha\beta} - \bar{\xi}^{(1)\alpha\beta} \Psi_{\mu\nu|\alpha\beta} \right) + \Delta \tilde{a}_1 + d(\dots). \quad (64)$$

This equation bears striking resemblance with its spin- $\frac{3}{2}$ counterpart Eqn. (29). We recall that, in the latter, cancelation of non- Γ -exact terms was possible by the insertion of identity (30) in the contraction of curvatures, the Bianchi identity $\partial_{[\mu} F_{\nu\rho]} = 0$, and the fermion EoMs in terms of curvature, $\gamma^\mu \Psi_{\mu\nu} = 0$, $\gamma^{\mu\nu} \Psi_{\mu\nu} = 0$. In the present case as well, as shown in Appendix A, the fermion EoMs can be written as the γ -traces of the curvature, $\gamma^\mu \Psi_{\mu\nu|\alpha\beta} = 0$, $\gamma^{\mu\nu} \Psi_{\mu\nu|\alpha\beta} = 0$. Therefore, the non- Γ -exact terms from the unambiguous piece in (64) can indeed be canceled by the Δ variation of a Γ -closed ambiguity,

$$\Delta \tilde{a}_1 = -\frac{1}{4} \left(\bar{\Psi}_{\alpha\beta} \mathcal{F} - 4 \bar{\Psi}_{\mu\nu|\alpha\beta} \gamma^\mu F^{\nu\rho} \gamma_\rho \right) \xi^{(1)\alpha\beta} + \text{h.c.} \quad (65)$$

Thus, we have a lift all the way to a_0 , the latter being a 3-derivative non-abelian vertex

$$a_0 = -\bar{\psi}_{\alpha\beta|\mu}^{(1)} F^{+\mu\nu} \psi^{(1)\alpha\beta|}_{\nu}. \quad (66)$$

4.2 Abelian Vertices

In this case, all the statements (37)–(38) hold, and the current in the vertex, $a_0 = j^\mu A_\mu$, is an invariant polynomial, which takes the most general form

$$j^\lambda = \bar{\Psi}_{\mu_1\nu_1|\mu_2\nu_2} X^{\mu_1\nu_1\alpha_1\beta_1\lambda\mu_2\nu_2\alpha_2\beta_2} \Psi_{\alpha_1\beta_1|\alpha_2\beta_2}. \quad (67)$$

Notice that the Fronsdal tensor, although allowed in principle, cannot appear in the current simply because it would render the vertex Δ -exact. In view of the spin- $\frac{5}{2}$ EoMs and the symmetry properties of the field strength, one can show, like in Section 3.4, that $X^{\mu_1\nu_1\alpha_1\beta_1\lambda\mu_2\nu_2\alpha_2\beta_2}$ can contain at most one derivative, which must carry one of the indices.

When X does not contain any derivative, the corresponding vertex will contain 4. In this case, we have the candidate $X^{\mu_1\nu_1\alpha_1\beta_1\lambda\mu_2\nu_2\alpha_2\beta_2} = -2\eta^{\mu_1\nu_1|\alpha_1\beta_1} \eta^{\mu_2\nu_2|\alpha_2\beta_2} \gamma^\lambda$. But again, the identities (30) and (44) tell us that the resulting vertex deforms nothing:

$$a_0 \approx \left(\bar{\Psi}_{\mu_1\nu_1|\mu_2\nu_2} \gamma^{\mu_1\nu_1\alpha_1\beta_1\lambda} \Psi_{\alpha_1\beta_1|}^{\mu_2\nu_2} \right) A_\lambda. \quad (68)$$

Finally, the 1-derivative candidate is $X^{\mu_1\nu_1\alpha_1\beta_1\lambda\mu_2\nu_2\alpha_2\beta_2} = \frac{1}{2} \eta^{\mu_1\nu_1|\alpha_1\beta_1} \eta^{\mu_2\nu_2|\alpha_2\beta_2} \overleftrightarrow{\partial}^\lambda$, which is equivalent to a 5-derivative 3-curvature term (Born-Infeld type),

$$a_0 \approx \bar{\Psi}_{\mu_1\nu_1|\mu_2\rho} \Psi^{\mu_1\nu_1|\rho}_{\nu_2} F^{\mu_2\nu_2}. \quad (69)$$

Below we present a summary table for all possible $1 - \frac{5}{2} - \frac{5}{2}$ vertices.

Table 4: Summary of $1 - \frac{5}{2} - \frac{5}{2}$ Vertices

# of Derivatives	Vertex	Nature	Exists in
3	$\bar{\psi}_{\alpha\beta \mu}^{(1)} F^{+\mu\nu} \psi^{(1)\alpha\beta }_{\nu}$	Non-abelian	$D \geq 4$
4	$\left(\bar{\Psi}_{\mu\nu \rho\sigma} \gamma^{\mu\nu\alpha\beta\lambda} \Psi_{\alpha\beta \rho\sigma} \right) A_{\lambda}$	Abelian	$D \geq 5$
5	$\bar{\Psi}_{\alpha\beta \mu\rho} \Psi^{\alpha\beta \rho}_{\nu} F^{\mu\nu}$	Abelian	$D \geq 4$

5 Arbitrary Spin: $s = n + \frac{1}{2}$

The set of fields and antifields in this case are

$$\Phi^A = \{A_{\mu}, C, \psi_{\mu_1 \dots \mu_n}, \xi_{\mu_1 \dots \mu_{n-1}}\}, \quad \Phi_A^* = \{A^{*\mu}, C^*, \bar{\psi}^{*\mu_1 \dots \mu_n}, \bar{\xi}^{*\mu_1 \dots \mu_{n-1}}\}. \quad (70)$$

For $n > 2$, there is an additional constraint that the field-antifield are triply γ -traceless:

$$\psi'_{\mu_1 \mu_3 \dots \mu_{n-3}} = 0, \quad \bar{\psi}'_{\mu_1 \mu_3 \dots \mu_{n-3}} = 0, \quad (71)$$

where prime denotes trace w.r.t Minkowski metric. Besides, the rank- $(n-1)$ fermionic ghost and its antighost are γ -traceless:

$$\xi_{\mu_1 \dots \mu_{n-2}} = 0, \quad \bar{\xi}_{\mu_1 \dots \mu_{n-2}} = 0. \quad (72)$$

Properties of the various fields and antifields are given in Table 5.

 Table 5: Properties of the Various Fields & Antifields ($n = \text{arbitrary}$)

Z	$\Gamma(Z)$	$\Delta(Z)$	$pgh(Z)$	$agh(Z)$	$gh(Z)$	$\epsilon(Z)$
A_{μ}	$\partial_{\mu} C$	0	0	0	0	0
C	0	0	1	0	1	1
$A^{*\mu}$	0	$-\partial_{\nu} F^{\mu\nu}$	0	1	-1	1
C^*	0	$-\partial_{\mu} A^{*\mu}$	0	2	-2	0
$\psi_{\mu_1 \dots \mu_n}$	$n \partial_{(\mu_1} \xi_{\mu_2 \dots \mu_n)}$	0	0	0	0	1
$\xi_{\mu_1 \dots \mu_{n-1}}$	0	0	1	0	1	0
$\bar{\psi}^{*\mu_1 \dots \mu_n}$	0	$\bar{\mathcal{R}}^{\mu_1 \dots \mu_n}$	0	1	-1	0
$\bar{\xi}^{*\mu_1 \dots \mu_{n-1}}$	0	$n \partial_{\mu_n} \bar{\psi}^{*\mu_1 \dots \mu_n}$	0	2	-2	1

The rank- n tensor-spinor $\mathcal{R}_{\mu_1 \dots \mu_n}$ appearing in the spin- s EoMs is an arbitrary-spin generalization of (50); it is related to the Fronsdal tensor as

$$\mathcal{R}_{\mu_1 \dots \mu_n} = \mathcal{S}_{\mu_1 \dots \mu_n} - \frac{1}{2} n \gamma_{(\mu_1} \mathcal{S}_{\mu_2 \dots \mu_n)} - \frac{1}{4} n(n-1) \eta_{(\mu_1 \mu_2} \mathcal{S}'_{\mu_3 \dots \mu_n)}. \quad (73)$$

The cohomology of Γ has already been given in Section 2.1, with the details appearing in Appendix B. One can immediately write down the set of all possible non-trivial a_2 's. Again, they fall into two subsets: Subset-1 contains both the bosonic ghost C and the fermionic ghost $\xi_{\mu_1 \dots \mu_{n-1}}$, while Subset-2 contains only $\xi_{\mu_1 \dots \mu_{n-1}}$ but not C .

- Subset-1 = $\left\{ C \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}}^{\bar{*}(m)} \xi^{(m) \mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}} + \text{h.c.} \right\}$,
- Subset-2 = $\left\{ C^* \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}}^{\bar{*}(m)} \xi^{(m) \mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}} \right\}$.

Here, $0 \leq m \leq n-1$. As a straightforward generalization of the spin- $\frac{5}{2}$ case, one finds that each element in Subset-1 gets lifted to a_1 :

$$a_1 = -\bar{\psi}^{*(m) \mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_n} \psi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_n}^{(m)} C - n \bar{\psi}^{*(m) \mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_n} \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m \| (\nu_{m+1} \dots \nu_{n-1})}^{(m)} A_{\nu_n} + \text{h.c.} + \tilde{a}_1. \quad (74)$$

Now, one can compute Δa_1 and expand it in the basis of $pg\hbar$ -1 objects in the cohomology of Γ , namely $\omega_I = \left\{ C, \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}}^{(m)} \mid 0 \leq m \leq n-1 \right\}$. Upon comparing the expansion coefficients for the unambiguous piece and the ambiguity \tilde{a}_1 , again one can conclude that none of these a_1 's can be lifted to an a_0 . On the other hand, for the elements of Subset-2, one notices that

$$\Delta \left(C^* \xi_{\mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}}^{\bar{*}(m)} \xi^{(m) \mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}} \right) = A^{*\nu_n} \bar{\xi}^{(m) \mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}} \partial_{\nu_n} \xi^{(m) \mu_1 \nu_1 | \dots | \mu_m \nu_m \| \nu_{m+1} \dots \nu_{n-1}} + \text{h.c.} + d(\dots). \quad (75)$$

Then, in view of Eqn. (B.9)–(B.10), it is clear that the right side of the above equation is Γ -exact modulo d only for $m = n-1$. This rules out, in particular, the would-be minimal coupling corresponding to $m = 0$. Therefore, one is left with the lift:

$$a_1 = -A^{*\nu_n} \bar{\psi}_{\mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1} \| \nu_n}^{(n-1)} \xi^{(n-1) \mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1}} + \text{h.c.} + \tilde{a}_1, \quad (76)$$

whose Δ -variation is given by

$$\Delta a_1 = \Gamma \left(\bar{\psi}_{\mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1} \| \mu_n}^{(n-1)} F_{\nu_n}^{\mu_n} \psi^{(n-1) \mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1} \| \nu_n} \right) + \Delta \tilde{a}_1 + d(\dots) + \frac{1}{2} F^{\mu_n \nu_n} \left(\bar{\Psi}_{\mu_1 \nu_1 | \dots | \mu_n \nu_n} \xi^{(n-1) \mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1}} + \text{h.c.} \right). \quad (77)$$

In view of Eqn. (29) and (64), pertaining respectively to the spin- $\frac{3}{2}$ and spin- $\frac{5}{2}$ cases, and the subsequent steps, we realize that it is possible to cancel the non- Γ -exact terms in (77) by inserting identity (30) in the contraction of curvatures, thanks to the Bianchi identity $\partial_{[\mu} F_{\nu\rho]} = 0$, and to the fermion EoMs in terms of curvature (see Appendix A), $\gamma^{\mu_1} \Psi_{\mu_1 \nu_1 | \dots | \mu_n \nu_n} = 0$, $\gamma^{\mu_1 \nu_1} \Psi_{\mu_1 \nu_1 | \dots | \mu_n \nu_n} = 0$. In other words, Δa_1 is rendered Γ -exact modulo d by an appropriate choice of the ambiguity \tilde{a}_1 , so that one finally has

$$a_0 = -\bar{\psi}_{\mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1} \| \mu_n}^{(n-1)} F_{\nu_n}^{+\mu_n} \psi^{(n-1) \mu_1 \nu_1 | \dots | \mu_{n-1} \nu_{n-1} \| \nu_n}. \quad (78)$$

This $(2n-1)$ -derivative non-abelian vertex contains the $(n-1)$ -curl of the fermionic field.

For an abelian vertex, $a_0 = j^\mu A_\mu$, the gauge-invariant current does not contain the Fronsdal tensor nor its derivatives, since their presence would render the vertex Δ -exact. Again, non-triviality of the abelian deformation allows only two possible values for the number of derivatives in the vertex: $2n$ and $2n+1$. The off-shell vertices can be obtained exactly the same way as for spins $\frac{3}{2}$ and $\frac{5}{2}$, considered in Sections 3.4 and 4.2 respectively. A summary table for all possible $1-s-s$ vertices is given below.

Table 6: Summary of $1-s-s$ Vertices with p Derivatives

p	Vertex	Nature	Exists in
$2n-1$	$\bar{\psi}_{\mu_1\nu_1 \dots \mu_{n-1}\nu_{n-1}}^{(n-1)} F^{+\mu_n}_{\nu_n} \psi^{(n-1)\mu_1\nu_1 \dots \mu_{n-1}\nu_{n-1}} \nu_n$	Non-abelian	$D \geq 4$
$2n$	$(\bar{\Psi}_{\mu_1\nu_1 \mu_2\nu_2 \dots \mu_n\nu_n} \gamma^{\mu_1\nu_1\alpha_1\beta_1\lambda} \Psi_{\alpha_1\beta_1 \mu_2\nu_2 \dots \mu_n\nu_n}) A_\lambda$	Abelian	$D \geq 5$
$2n+1$	$\bar{\Psi}_{\mu_1\nu_1 \mu_2\nu_2 \dots \mu_n\alpha} \Psi^{\mu_1\nu_1 \mu_2\nu_2 \dots \alpha\nu_n} F^{\mu_n}_{\nu_n}$	Abelian	$D \geq 4$

6 Abelian Vertices Preserve Gauge Symmetries

Abelian vertices are those that do not deform the gauge algebra, i.e., they can only have a trivial a_2 . For such a vertex, it is always possible to choose a_1 to be Γ -closed [17]:

$$\Gamma a_1 = 0. \quad (79)$$

If this gets lifted to an a_0 , one has the cocycle condition (18),

$$\Delta a_1 + \Gamma a_0 + db_0 = 0. \quad (80)$$

For the $1-s-s$ vertices under study, one can always write a vertex as the photon field A_μ contracted with a current j^μ , which is a fermion bilinear:

$$a_0 = j^\mu A_\mu. \quad (81)$$

One can always choose the current such that it satisfies

$$\Gamma j^\mu = 0, \quad \partial_\mu j^\mu = \Delta M, \quad \Gamma M = 0. \quad (82)$$

To see this, let us note that the a_1 corresponding to (81) has the general form

$$a_1 = MC + (\bar{P}_{\mu_1\dots\mu_{n-1}} \xi^{\mu_1\dots\mu_{n-1}} - \bar{\xi}_{\mu_1\dots\mu_{n-1}} P^{\mu_1\dots\mu_{n-1}}) + a'_1, \quad (83)$$

where M and $P_{\mu_1 \dots \mu_{n-1}}$ belong to $H(\Gamma)$, with $pg h = 0$, $agh = 1$, and a'_1 stands for expansion terms in the ghost-curls. Given (81) and (83), the condition (80) reads

$$\Gamma(j^\mu A_\mu) + \Delta MC + (\Delta \bar{P}_{\mu_1 \dots \mu_{n-1}} \xi^{\mu_1 \dots \mu_{n-1}} - \bar{\xi}_{\mu_1 \dots \mu_{n-1}} \Delta P^{\mu_1 \dots \mu_{n-1}}) + \Delta a'_1 + db_0 = 0. \quad (84)$$

Now, $P_{\mu_1 \dots \mu_{n-1}}$ consists of two kinds of terms: one contains the antifield $A^{*\mu}$ and its derivatives, and the other contains the antifield $\psi^{*\nu_1 \dots \nu_n}$ and its derivatives. The former one also contains (derivatives of) the Fronsdal tensor $\mathcal{S}_{\nu_1 \dots \nu_n}$, or (derivatives of) the curvature $\Psi_{\rho_1 \nu_1 | \dots | \rho_n \nu_n}$, while the latter one contains (derivatives of) the EM field strength $F_{\mu\nu}$. One can choose to get rid of derivatives on $A^{*\mu}$ and $F_{\mu\nu}$ by using the Leibniz rule,

$$\begin{aligned} P_{\mu_1 \dots \mu_{n-1}} &= A^{*\mu} \left(\vec{P}_{\mu, \mu_1 \dots \mu_{n-1}}^{(\mathcal{S}) \nu_1 \dots \nu_n} \mathcal{S}_{\nu_1 \dots \nu_n} + \vec{P}_{\mu, \mu_1 \dots \mu_{n-1}}^{(\Psi) \rho_1 \nu_1 | \dots | \rho_n \nu_n} \Psi_{\rho_1 \nu_1 | \dots | \rho_n \nu_n} \right) \\ &\quad + F^{\mu\nu} \left(\vec{P}_{\mu\nu, \mu_1 \dots \mu_{n-1}}^{(\psi^*) \nu_1 \dots \nu_n} \psi_{\nu_1 \dots \nu_n}^* \right) + \partial^{\mu_n} p_{\mu_1 \dots \mu_n}, \end{aligned} \quad (85)$$

where $\Gamma p_{\mu_1 \dots \mu_n} = 0$, and the \vec{P} 's are operators acting to the right. Notice that the quantity in the parentheses in the first line is both Γ -closed and Δ -exact⁷. One can take the Δ -variation of (85), and then add a total derivative in order to cast it in the form

$$\Delta P_{\mu_1 \dots \mu_{n-1}} = \frac{1}{2} F^{\mu\nu} \Delta Q_{[\mu\nu], \mu_1 \dots \mu_{n-1}} + \partial^{\mu_n} \Delta q_{\mu_1 \dots \mu_n}, \quad (86)$$

where, $\Gamma Q_{[\mu\nu], \mu_1 \dots \mu_{n-1}} = 0$, $\Gamma q_{\mu_1 \dots \mu_n} = 0$. Therefore, we have

$$\bar{\xi}_{\mu_1 \dots \mu_{n-1}} \Delta P^{\mu_1 \dots \mu_{n-1}} = A_\mu \Delta \left[\partial_\nu (\bar{\xi}_{\mu_1 \dots \mu_{n-1}} Q^{[\mu\nu], \mu_1 \dots \mu_{n-1}}) \right] - \bar{\xi}_{\mu_1 \dots \mu_{n-1}} \overleftarrow{\partial}_{\mu_n} \Delta q^{\mu_1 \dots \mu_n} + d(\dots). \quad (87)$$

The second term on the right side is Γ -closed, and can be broken as a Γ -exact piece plus terms involving the ghost-curls. The latter can always be canceled in the cocycle condition (84) by appropriately choosing the similar terms coming from a'_1 . Thus,

$$\begin{aligned} \Gamma \left[j^\mu A_\mu + \Delta (\bar{\psi}_{\mu_1 \dots \mu_n} q^{\mu_1 \dots \mu_n} + \text{h.c.}) \right] + \Delta MC \\ - A_\mu \Delta \left[\partial_\nu (\bar{\xi}_{\mu_1 \dots \mu_{n-1}} Q^{[\mu\nu], \mu_1 \dots \mu_{n-1}}) + \text{h.c.} \right] + d(\dots) = 0. \end{aligned} \quad (88)$$

The Δ -exact term added to the original vertex $j^\mu A_\mu$ is trivial, and therefore can be dropped. Now we are left with

$$A_\mu \left[\Gamma j^\mu - \Delta (\partial_\nu (\bar{\xi}_{\mu_1 \dots \mu_{n-1}} Q^{[\mu\nu], \mu_1 \dots \mu_{n-1}}) + \text{h.c.}) \right] + (\Delta M - \partial_\mu j^\mu) C + d(\dots) = 0. \quad (89)$$

Taking functional derivative w.r.t. C produces part of the sought-after conditions (82),

$$\partial_\mu j^\mu = \Delta M, \quad \Gamma M = 0, \quad (90)$$

⁷ Δ -exactness of the first term is manifest, while in the second, the presence of the curvature admits only Δ -exact terms, like its own γ -traces and divergences (see Appendix A).

while the functional derivative w.r.t. A_μ gives

$$\Gamma j^\mu = \partial_\nu \left(\bar{\xi}_{\mu_1 \dots \mu_{n-1}} \Delta Q^{[\mu\nu], \mu_1 \dots \mu_{n-1}} \right) + \text{h.c.}, \quad \Gamma Q^{[\mu\nu], \mu_1 \dots \mu_{n-1}} = 0. \quad (91)$$

The expression for Γj^μ has to be Γ -exact. This demands that $\partial_\nu Q^{[\mu\nu], \mu_1 \dots \mu_{n-1}}$ be Δ -closed, and that $Q^{[\mu\nu], \mu_1 \dots \mu_{n-1}}$ have the interchange symmetry $\nu \leftrightarrow \mu_i$, $i = 1, \dots, n-1$. Then,

$$j^\alpha = \tilde{j}^\alpha + \Delta \left(\frac{1}{n} \bar{\psi}_{\mu_1 \dots \mu_n} Q^{[\alpha\mu_1], \mu_2 \dots \mu_n} + \text{h.c.} \right), \quad \Gamma \tilde{j}^\alpha = 0. \quad (92)$$

Therefore, by field redefinitions, the current can always be made gauge invariant:

$$\Gamma j^\mu = 0. \quad (93)$$

This completes the proof of (82). Then, from (80), one obtains the lift:

$$a_1 = MC. \quad (94)$$

Now we will show that M must be Δ -exact modulo d . We recall that M belongs to the cohomology of Γ , with $pgh = 0$, $agh = 1$. It will contain (derivatives of) the fermionic antifield, and (derivatives of) the Fronsdal tensor $\mathcal{S}_{\nu_1 \dots \nu_n}$ or the curvature $\Psi_{\rho_1 \nu_1 | \dots | \rho_n \nu_n}$. However, one can choose to have no derivatives of the antifield by using the Leibniz rule. Thus M has the most general form

$$M = \bar{\psi}^{*\mu_1 \dots \mu_n} \left(\vec{M}_{\mu_1 \dots \mu_n}^{(\mathcal{S}) \nu_1 \dots \nu_n} \mathcal{S}_{\nu_1 \dots \nu_n} + \vec{M}_{\mu_1 \dots \mu_n}^{(\Psi) \rho_1 \nu_1 | \dots | \rho_n \nu_n} \Psi_{\rho_1 \nu_1 | \dots | \rho_n \nu_n} \right) + \partial^\mu m_\mu - \text{h.c.}, \quad (95)$$

where $\Gamma m_\mu = 0$, and the operators \vec{M} 's act to the right. The first term in the parentheses is manifestly Δ -exact, while the second one must contain either a γ -trace and or a divergence of the curvature, which are Δ -exact too (see Appendix A). Therefore, M must be Δ -exact modulo d . This means that a_1 , given in (94), can be rendered trivial by adding a Δ -exact piece in a_0 [17], and so the vertex will be gauge invariant up to a total derivative:

$$\Gamma a_0 + db_0 = 0. \quad (96)$$

In other words, one can always add a Δ -exact term in a_0 , so that the new current is identically conserved [17]:

$$j^\mu \rightarrow j'^\mu = j^\mu + \Delta k^\mu = \partial_\nu \mathcal{A}^{\mu\nu}, \quad \mathcal{A}^{\mu\nu} = -\mathcal{A}^{\nu\mu}. \quad (97)$$

Thus we have proved that no abelian vertex can deform the gauge transformations.

7 Comparative Study of Vertices

We have found that the possible number of derivatives in a $1-s-s$ vertex, with $s = n + \frac{1}{2}$, is restricted to the values: $2n-1$, $2n$, and $2n+1$. Moreover, the $2n$ -derivative vertex exists only in $D \geq 5$. These are in complete agreement with Metsaev's light-cone-formulation results [7]. While the light-cone vertices are maximally gauge fixed, the corresponding covariant on-shell vertices were also written down in [7] for lower spins, from previously known results. These on-shell vertices are partially gauge fixed, with the gauge choice being the transverse-traceless gauge (TT gauge),

$$\partial^{\mu_1} \psi_{\mu_1 \dots \mu_n} = 0, \quad \gamma^{\mu_1} \psi_{\mu_1 \dots \mu_n} = 0, \quad \partial^\mu A_\mu = 0. \quad (98)$$

Note that in this gauge the fermion and photon EoMs boil down to

$$\not{\partial} \psi_{\mu_1 \dots \mu_n} = 0, \quad \square A_\mu = 0. \quad (99)$$

We will find that our off-shell vertices reduce in the TT gauge to the on-shell ones given in [7]. So do the Sagnotti-Taronna (ST) off-shell vertices [10], as we will see. If two vertices are shown to match in a particular gauge, say the TT gauge, the full off-shell ones must be equivalent, i.e., differ only by terms that are Δ -exact modulo d . Still, for the simplest case of spin $\frac{3}{2}$, we will make explicit the off-shell equivalence of the ST vertices with ours. For $s \geq \frac{5}{2}$, we match our vertices with the ST ones in the TT gauge.

The ST off-shell vertices, when read off in the most naive way, contain many terms, and it is not straightforward at all to see that some of them actually vanish in $D = 4$. In comparison, the off-shell vertices we present for arbitrary spin are rather neat in form, and the absence of some of them in $D = 4$ is obvious from inspection.

We will denote a p -derivative off-shell vertex of ours as $V^{(p)}$, and its Sagnotti-Taronna counterpart as $V_{\text{ST}}^{(p)}$. The corresponding TT-gauge vertex will be denoted as $V_{\text{TT}}^{(p)}$ ⁸.

7.1 1—3/2—3/2 Vertices

Our 1-derivative off-shell $1 - \frac{3}{2} - \frac{3}{2}$ vertex is given by

$$V^{(1)} = \bar{\psi}_\mu F^{+\mu\nu} \psi_\nu = \bar{\psi}_\mu \left(\eta^{\mu\nu|\alpha\beta} + \frac{1}{2} \gamma^{\mu\nu\alpha\beta} \right) F_{\alpha\beta} \psi_\nu. \quad (100)$$

To see what it reduces to in the TT gauge, let us rewrite identity (30) as

$$\begin{aligned} \eta^{\mu\nu|\alpha\beta} + \frac{1}{2} \gamma^{\mu\nu\alpha\beta} &= -\frac{1}{2} \eta^{\mu\nu} \gamma^{\alpha\beta} + \frac{1}{2} \gamma^\mu \gamma^\nu \gamma^{\alpha\beta} - 2\gamma^{[\mu} \eta^{\nu][\alpha} \gamma^{\beta]} \\ &= 2 \left(\eta^{\mu\nu|\alpha\beta} - \frac{1}{4} \eta^{\mu\nu} \gamma^{\alpha\beta} \right) + \frac{1}{4} \left(\gamma^\mu \gamma^\nu \gamma^{\alpha\beta} + \gamma^{\alpha\beta} \gamma^\mu \gamma^\nu \right), \end{aligned} \quad (101)$$

⁸Both the spins $m + \frac{1}{2}$ and $m + \frac{3}{2}$ will respectively have one vertex with $2m + 1$ derivatives. Our notation should not cause any confusion, as we will be considering one spin at a time.

where in the second line we have used $2\gamma^{[\mu}\eta^{\nu][\alpha}\gamma^{\beta]} = \frac{1}{4}(\gamma^\mu\gamma^\nu\gamma^{\alpha\beta} - \gamma^{\alpha\beta}\gamma^\mu\gamma^\nu) - 2\eta^{\mu\nu|\alpha\beta}$. Upon insertion of identity (101) into the vertex (100), one finds

$$V^{(1)} = 2(\bar{\psi}_\mu F^{\mu\nu}\psi_\nu - \frac{1}{4}\bar{\psi}_\mu \mathcal{F}\psi^\mu) + \frac{1}{4}(\bar{\psi}\gamma^\mu \mathcal{F}\psi_\mu + \bar{\psi}_\mu \mathcal{F}\gamma^\mu \psi). \quad (102)$$

On the other hand, the 1-derivative ST vertex reads [10]

$$V_{\text{ST}}^{(1)} = \bar{\psi}^\mu(\partial_\nu\psi_\mu)A^\nu - (\partial_\mu\bar{\psi}_\nu)\psi^\nu A^\mu + \bar{\psi}_\mu\psi_\nu(\partial^\mu A^\nu) - \bar{\psi}_\mu(\partial^\mu\psi^\nu)A_\nu + (\partial_\mu\bar{\psi}_\nu)\psi^\mu A^\nu - \bar{\psi}_\mu\psi_\nu(\partial^\nu A^\mu). \quad (103)$$

Integrating by parts the 2nd, 4th and 5th terms on the right hand side, we obtain

$$V_{\text{ST}}^{(1)} = 2\bar{\psi}_\mu F^{\mu\nu}\psi_\nu + 2\bar{\psi}_\mu A \cdot \partial\psi^\mu + \bar{\psi}_\mu(\partial \cdot A)\psi^\mu + (\partial \cdot \bar{\psi})A \cdot \psi - \bar{\psi} \cdot A(\partial \cdot \psi) + d(\dots). \quad (104)$$

Let us take the 2nd term on the right hand side and replace $\eta^{\alpha\beta} = \gamma^{(\alpha}\gamma^{\beta)}$ in the operator $(A \cdot \partial)$. Also in the 3rd term we replace $\eta^{\alpha\beta} = \gamma^\alpha\gamma^\beta - \gamma^{\alpha\beta}$ in $(\partial \cdot A)$. The result is

$$2\bar{\psi}_\mu A \cdot \partial\psi^\mu + \bar{\psi}_\mu(\partial \cdot A)\psi^\mu = -\frac{1}{2}\bar{\psi}_\mu \mathcal{F}\psi^\mu + \bar{\psi}_\mu \mathcal{A}(\not{\partial}\psi^\mu) - (\not{\partial}\bar{\psi}_\mu)\mathcal{A}\psi^\mu + d(\dots), \quad (105)$$

which, when plugged into the vertex (104) gives

$$V_{\text{ST}}^{(1)} = 2(\bar{\psi}_\mu F^{\mu\nu}\psi_\nu - \frac{1}{4}\bar{\psi}_\mu \mathcal{F}\psi^\mu) + [\bar{\psi}_\mu \mathcal{A}(\not{\partial}\psi^\mu) - \bar{\psi} \cdot A(\partial \cdot \psi) + \text{h.c.}] + d(\dots). \quad (106)$$

It is obvious that both the off-shell vertices (102) and (106) reduce in the TT gauge to

$$V_{\text{TT}}^{(1)} = 2(\bar{\psi}_\mu F^{\mu\nu}\psi_\nu - \frac{1}{4}\bar{\psi}_\mu \mathcal{F}\psi^\mu). \quad (107)$$

This is precisely the on-shell 1-derivative vertex reported by Metsaev [7]. To see explicitly that the off-shell vertices are also equivalent, we subtract (106) from (102) to get

$$V^{(1)} - V_{\text{ST}}^{(1)} = \frac{1}{4}(\bar{\psi}\gamma^\mu \mathcal{F}\psi_\mu + \bar{\psi}_\mu \mathcal{F}\gamma^\mu \psi) - [\bar{\psi}_\mu \mathcal{A}(\not{\partial}\psi^\mu) - \bar{\psi} \cdot A(\partial \cdot \psi) + \text{h.c.}] + d(\dots). \quad (108)$$

Now we make use of the identity

$$[\gamma^\mu, \gamma^{\alpha\beta}] = 4\eta^{\mu[\alpha}\gamma^{\beta]}, \quad (109)$$

in order to be able to pass γ^μ past \mathcal{F} in both the terms in the parentheses on the right hand side of Eqn. (108). As a result, we will obtain, among others, the term $\frac{1}{2}\bar{\psi}\mathcal{F}\psi$, in which we replace $\mathcal{F} = \not{\partial}\mathcal{A} - \partial \cdot A$. Now in all the resulting terms we perform integrations by parts such that no derivative acts on the photon field. The final result is

$$V^{(1)} - V_{\text{ST}}^{(1)} = [2\bar{\psi}^{[\mu}A^{\nu]}\gamma_\mu(\not{\partial}\psi_\nu - \partial_\nu\psi) - \bar{\psi}^\mu A^\nu\gamma_{\mu\nu}(\partial \cdot \psi - \not{\partial}\psi) + \text{h.c.}] + d(\dots). \quad (110)$$

This is manifestly Δ -exact modulo d , which proves the equivalence of the off-shell vertices:

$$V^{(1)} \approx V_{\text{ST}}^{(1)}. \quad (111)$$

Next, we consider the 2-derivative vertex,

$$V^{(2)} = (\bar{\Psi}_{\mu\nu} \gamma^{\mu\nu\alpha\beta\lambda} \Psi_{\alpha\beta}) A_\lambda \approx -2 (\bar{\Psi}_{\mu\nu} \gamma^\rho \Psi^{\mu\nu}) A_\rho. \quad (112)$$

One can use the definition $\Psi_{\mu\nu} = 2\partial_{[\mu}\psi_{\nu]}$ to rewrite it as

$$V^{(2)} \approx -4\bar{\psi}_\alpha \overleftarrow{\partial}_\mu \not{A} \partial^\mu \psi^\alpha + 2 \left(\bar{\psi}_\alpha \overleftarrow{\partial}_\mu \not{A} \partial^\alpha \psi^\mu + \text{h.c.} \right). \quad (113)$$

In the 1st term, we can use the 3-box rule, already given in Section (3.4),

$$2\partial_\mu X \partial^\mu Y = \square(XY) - X(\square Y) - (\square X)Y, \quad (114)$$

and perform a double integration by parts in order to have a \square acting on the photon field. In the 2nd term on the right hand side of (112) one can integrate by parts w.r.t. any of the derivatives. When the derivative acts on the photon field, one can use $\partial_\mu A_\nu = F_{\mu\nu} + \partial_\nu A_\mu$ to rewrite it in terms of the field strength. The result is

$$\begin{aligned} V^{(2)} \approx & 2 \left(\bar{\psi}_\alpha \gamma^\mu \partial^\alpha \psi^\nu - \bar{\psi}^\mu \overleftarrow{\partial}^\alpha \gamma^\nu \psi_\alpha \right) F_{\mu\nu} - 2 \left[(\bar{\psi}_\alpha \gamma^\mu \partial^\alpha \psi^\nu) \partial_\mu A_\nu + \text{h.c.} \right] \\ & - 2\bar{\psi}_\alpha \square \not{A} \psi^\alpha + 2 \left[\bar{\psi}_\alpha \not{A} (\square \psi^\alpha - \partial^\alpha \partial \cdot \psi) + \text{h.c.} \right]. \end{aligned} \quad (115)$$

Now, in the last term of the first line we perform integration by parts so that no derivative acts on the photon field. On the other hand, the last term in the second line is Δ -exact, and therefore can be dropped. Thus we are left with

$$\begin{aligned} V^{(2)} \approx & 2 \left(\bar{\psi}_\alpha \gamma^\mu \partial^\alpha \psi^\nu - \bar{\psi}^\mu \overleftarrow{\partial}^\alpha \gamma^\nu \psi_\alpha \right) F_{\mu\nu} \\ & + 2 \left[\left(\bar{\psi}_\alpha \partial^\alpha \not{\partial} \psi^\nu + \bar{\psi}_\alpha \overleftarrow{\not{\partial}} \partial^\alpha \psi^\nu \right) A_\nu + \text{h.c.} \right] - 2\bar{\psi}_\alpha \square \not{A} \psi^\alpha. \end{aligned} \quad (116)$$

As one reads off the 2-derivative ST vertex, it gives

$$\begin{aligned} V_{\text{ST}}^{(2)} = & -\bar{\psi}^\mu \gamma^\alpha (\partial_\mu \psi_\nu) \partial^\nu A_\alpha + \bar{\psi}^\mu \gamma^\alpha \psi^\nu \partial_\mu \partial_\nu A_\alpha + \bar{\psi}^\mu \overleftarrow{\partial}^\nu \gamma^\alpha (\partial_\mu \psi_\nu) A_\alpha - \bar{\psi}^\mu \overleftarrow{\partial}^\nu \gamma^\alpha \psi_\nu \partial_\mu A_\alpha \\ & - (\partial \cdot \bar{\psi}) \not{A} \partial \cdot \psi - \bar{\psi} \psi^\nu \partial_\nu \partial \cdot A + \bar{\psi} \overleftarrow{\partial}_\nu \psi^\nu \partial \cdot A + \bar{\psi}^\mu (\partial_\mu \not{\psi}) \partial \cdot A \\ & - \bar{\psi}^\mu \not{\psi} \partial_\mu \partial \cdot A - \bar{\psi} (\partial \cdot \psi) \partial \cdot A - (\partial \cdot \bar{\psi}) \not{\psi} \partial \cdot A. \end{aligned} \quad (117)$$

As we mentioned already, in this form it is not evident at all that this vertex vanishes for $D = 4$. Let us integrate by parts the 2nd and 3rd terms in the first line of Eqn. (117), w.r.t. ∂_μ . The 2nd term in the second line and the 1st term in the third line contain the gradient of $\partial \cdot A$; we integrate by parts the gradient in both these terms. Thus we have

$$\begin{aligned} V_{\text{ST}}^{(2)} = & -2\bar{\psi}_\mu \gamma^\alpha (\partial^\mu \psi^\nu) \partial_\nu A_\alpha - (\partial \cdot \bar{\psi}) (\partial_\nu \not{A}) \psi^\nu - 2\bar{\psi}^\mu \overleftarrow{\partial}^\nu \gamma^\alpha \psi_\nu \partial_\mu A_\alpha - (\partial \cdot \bar{\psi}) \overleftarrow{\partial}_\nu \not{A} \psi^\nu \\ & - (\partial \cdot \bar{\psi}) \not{A} \partial \cdot \psi + 2(\partial \cdot A) \left(\bar{\psi}^\mu \partial_\mu \not{\psi} + \bar{\psi} \overleftarrow{\partial}^\mu \psi_\mu \right) + d(\dots). \end{aligned} \quad (118)$$

Notice that the 2nd, 4th and 5th terms combine into a total derivative. One can rewrite the 1st and 3rd terms in terms of the photon field strength by using $\partial_\mu A_\nu = F_{\mu\nu} + \partial_\nu A_\mu$. Also, one can extract a Δ -exact piece, by using EoMs: $\not{\partial}\psi_\mu - \partial_\mu\psi = 0$, in the term containing $(\partial \cdot A)$. This leaves us with

$$V_{\text{ST}}^{(2)} \approx 2 \left(\bar{\psi}_\alpha \gamma^\mu \partial^\alpha \psi^\nu - \bar{\psi}^\mu \overleftarrow{\partial}^\alpha \gamma^\nu \psi_\alpha \right) F_{\mu\nu} - 2 \left[(\bar{\psi}_\alpha \gamma^\mu \partial^\alpha \psi^\nu) \partial_\mu A_\nu + \text{h.c.} \right] + 2(\partial \cdot A) \left(\bar{\psi}^\mu \not{\partial} \psi_\mu + \bar{\psi}^\mu \overleftarrow{\not{\partial}} \psi_\mu \right). \quad (119)$$

Again, we integrate by parts the last term of the first line, so that no derivatives act on the photon field. In the second line as well we perform integration by parts to have 2 derivatives acting on the photon field. This finally gives

$$V_{\text{ST}}^{(2)} \approx 2 \left(\bar{\psi}_\alpha \gamma^\mu \partial^\alpha \psi^\nu - \bar{\psi}^\mu \overleftarrow{\partial}^\alpha \gamma^\nu \psi_\alpha \right) F_{\mu\nu} + 2 \left[\left(\bar{\psi}_\alpha \partial^\alpha \not{\partial} \psi^\nu + \bar{\psi}_\alpha \overleftarrow{\not{\partial}} \partial^\alpha \psi^\nu \right) A_\nu + \text{h.c.} \right] - 2\bar{\psi}_\alpha (\not{\partial} \partial \cdot A) \psi^\alpha. \quad (120)$$

It is clear that, in the TT gauge, both the off-shell vertices (116) and (120) reduce to

$$V_{\text{TT}}^{(2)} = 2 \left(\bar{\psi}_\alpha \gamma^\mu \partial^\alpha \psi^\nu - \bar{\psi}^\mu \overleftarrow{\partial}^\alpha \gamma^\nu \psi_\alpha \right) F_{\mu\nu}, \quad (121)$$

which is nothing but the 2-derivative on-shell vertex given in [7]. The equivalence of the two off-shell vertices is also evident as, upon subtracting (120) from (116), we have

$$V^{(2)} - V_{\text{ST}}^{(2)} = 2\bar{\psi}_\alpha \gamma^\mu (\partial^\nu F_{\mu\nu}) \psi^\alpha = \Delta\text{-exact}. \quad (122)$$

Finally, we consider the vertex with 3 derivatives, which reads

$$V^{(3)} = \bar{\Psi}_{\mu\alpha} \Psi_\nu^\alpha F^{\mu\nu} = (\partial_\mu \bar{\psi}^\alpha - \partial^\alpha \bar{\psi}_\mu) (\partial_\alpha \psi_\nu - \partial_\nu \psi_\alpha) F^{\mu\nu}. \quad (123)$$

Integration by parts w.r.t. ∂_μ , appearing in the 1st term inside the first parentheses, gives

$$V^{(3)} \approx -\bar{\psi}^\alpha (\partial_\mu \partial_\alpha \psi_\nu - \partial_\mu \partial_\nu \psi_\alpha) F^{\mu\nu} - \bar{\psi}^\alpha \Psi_{\alpha\nu} \partial_\mu F^{\mu\nu} - \bar{\psi}_\mu \overleftarrow{\partial}^\alpha \partial_\alpha \psi_\nu F^{\mu\nu} + \bar{\psi}_\mu \overleftarrow{\partial}^\alpha \partial_\nu \psi_\alpha F^{\mu\nu}. \quad (124)$$

Here the 2nd term inside the parentheses on the right side is identically zero, while the term containing $\partial_\mu F^{\mu\nu}$ is Δ -exact. We use the 3-box rule (114) in the penultimate term. Also we integrate by parts w.r.t. ∂_ν in the last term, and it produces a Δ -exact piece, containing $\partial_\nu F^{\mu\nu}$, that we discard. The result is

$$V^{(3)} \approx - \left(\bar{\psi}^\alpha \partial_\alpha \partial_\mu \psi_\nu + \bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \right) F^{\mu\nu} + \frac{1}{2} \left[\bar{\psi}_\mu \square \psi_\nu + \bar{\psi}_\mu \overleftarrow{\square} \psi_\nu - \square (\bar{\psi}_\mu \psi_\nu) \right] F^{\mu\nu}. \quad (125)$$

Now, one can perform a double integration by parts in the last term in the brackets in order to have $\square F^{\mu\nu}$, which gives a Δ -exact piece, so that we finally have

$$V^{(3)} \approx - \left(\bar{\psi}^\alpha \partial_\alpha \partial_\mu \psi_\nu + \bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \right) F^{\mu\nu} + \frac{1}{2} \left(\bar{\psi}_\mu \square \psi_\nu + \bar{\psi}_\mu \overleftarrow{\square} \psi_\nu \right) F^{\mu\nu}. \quad (126)$$

On the other hand, the 3-derivative off-shell ST vertex contains as many as 14 terms:

$$\begin{aligned}
V_{\text{ST}}^{(3)} = & (\partial_\alpha \bar{\psi}^\mu)(\partial_\mu \psi^\nu) \partial_\nu A^\alpha - \bar{\psi}^\mu (\partial_\alpha \partial_\mu \psi^\nu) \partial_\nu A^\alpha - (\partial_\alpha \partial_\nu \bar{\psi}^\mu)(\partial_\mu \psi^\nu) A^\alpha + (\partial_\nu \bar{\psi}^\mu)(\partial_\mu \partial_\alpha \psi^\nu) A^\alpha \\
& - (\partial_\alpha \bar{\psi}^\mu) \psi^\nu \partial_\mu \partial_\nu A^\alpha + \bar{\psi}^\mu (\partial_\alpha \psi^\nu) \partial_\mu \partial_\nu A^\alpha + (\partial_\alpha \partial_\nu \bar{\psi}^\mu) \psi^\nu \partial_\mu A^\alpha - (\partial_\nu \bar{\psi}^\mu)(\partial_\alpha \psi^\nu) \partial_\mu A^\alpha \\
& + (\partial_\mu \partial \cdot \bar{\psi})(\partial \cdot \psi) A^\mu - (\partial \cdot \bar{\psi})(\partial_\mu \partial \cdot \psi) A^\mu + \bar{\psi}^\mu (\partial_\mu \partial_\alpha \psi^\alpha) \partial \cdot A \\
& - \bar{\psi}^\mu (\partial \cdot \psi) \partial_\mu \partial_\alpha A^\alpha - (\partial_\mu \partial_\alpha \bar{\psi}^\alpha) \psi^\mu \partial \cdot A + (\partial \cdot \bar{\psi}) \psi^\mu \partial_\mu \partial_\alpha A^\alpha.
\end{aligned}$$

Here we will perform a number of integrations by parts. In the first line, we integrate by parts the 1st term w.r.t. ∂_α , the 3rd w.r.t. ∂_μ , and the 4th w.r.t. ∂_ν . In the second line, the 1st, 2nd and 4th terms are integrated by parts respectively w.r.t. ∂_ν , ∂_μ and ∂_α . In the third line, this is done only on the 3rd term w.r.t. ∂_α . Finally, in the fourth line, the 1st and 3rd terms are integrated by parts w.r.t. both ∂_μ and ∂_α , while the 2nd one only w.r.t. ∂_α . Dropping total derivatives, the result is

$$\begin{aligned}
V_{\text{ST}}^{(3)} \approx & -4\bar{\psi}^\alpha (\partial_\alpha \partial_\mu \psi_\nu) \partial^\nu A^\mu + 4\bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \partial^\mu A^\nu + 2 \left(\bar{\psi}^\mu \overleftarrow{\partial}_\alpha \psi^\alpha - \bar{\psi}^\alpha \partial_\alpha \psi^\mu \right) \partial_\mu \partial \cdot A \\
& + 2 \left(\bar{\psi} \cdot \overleftarrow{\partial} \overleftarrow{\partial}_\mu \overleftarrow{\partial}_\alpha \psi^\alpha - \bar{\psi}^\alpha \partial_\alpha \partial_\mu \partial \cdot \psi \right) A^\mu + \left(\bar{\psi} \cdot \overleftarrow{\partial} \overleftarrow{\partial}_\mu \partial_\alpha \psi^\mu - \bar{\psi}^\mu \overleftarrow{\partial}_\alpha \partial_\mu \partial \cdot \psi \right) A^\alpha \\
& + [(\partial_\alpha \partial \cdot \bar{\psi}) \partial \cdot \psi + \text{h.c.}] A^\alpha + [(\partial_\alpha \bar{\psi}_\mu) \partial \cdot \psi + \text{h.c.}] \partial^\mu A^\alpha. \tag{127}
\end{aligned}$$

Let us rewrite the first two terms in the first line in terms of the photon field strength by using $\partial_\mu A_\nu = F_{\mu\nu} + \partial_\nu A_\mu$, and use the 3-box rule (114) in the additional terms. Also, we notice that the last line in (127) reduces exactly to the 2nd term on the second line, up to a total derivative. Then, the vertex reads

$$\begin{aligned}
V_{\text{ST}}^{(3)} \approx & 4 \left(\bar{\psi}^\alpha \partial_\alpha \partial_\mu \psi_\nu + \bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \right) F^{\mu\nu} + 2 \left(\bar{\psi} \cdot \overleftarrow{\partial} \overleftarrow{\partial}_\mu \partial_\alpha \psi^\mu - \bar{\psi}^\mu \overleftarrow{\partial}_\alpha \partial_\mu \partial \cdot \psi \right) A^\alpha \\
& + 2 \left(\bar{\psi}_\mu \overleftarrow{\partial}_\nu \square \psi^\nu - \bar{\psi}^\nu \square \partial_\nu \psi_\mu \right) A^\mu - 2 \left(\bar{\psi}^\mu \overleftarrow{\partial}_\alpha \psi^\alpha - \bar{\psi}^\alpha \partial_\alpha \psi^\mu \right) (\square A_\mu - \partial_\mu \partial \cdot A) \\
& + 2 \left[\bar{\psi}^\alpha \partial_\alpha (\square \psi_\mu - \partial_\mu \partial \cdot \psi) - \left(\bar{\psi}_\mu \overleftarrow{\square} - \bar{\psi} \cdot \overleftarrow{\partial} \overleftarrow{\partial}_\mu \right) \overleftarrow{\partial}_\alpha \psi^\alpha \right] A^\mu. \tag{128}
\end{aligned}$$

Clearly, the 2nd term in the second line and the entire third line are Δ -exact, while, modulo Δ -exact pieces, the 1st term in the second line can have $\square \psi^\nu$ replaced by $\partial^\nu \partial \cdot \psi$. The latter result can be combined with the 2nd term in the first line to give

$$\begin{aligned}
V_{\text{ST}}^{(3)} \approx & 4 \left(\bar{\psi}^\alpha \partial_\alpha \partial_\mu \psi_\nu + \bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \right) F^{\mu\nu} - 2 \left(\bar{\psi} \cdot \overleftarrow{\partial} \overleftarrow{\partial}_\mu \Psi^{\mu\nu} - \bar{\Psi}^{\mu\nu} \partial_\mu \partial \cdot \psi \right) A_\nu \\
\approx & 4 \left(\bar{\psi}^\alpha \partial_\alpha \partial_\mu \psi_\nu + \bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \right) F^{\mu\nu} + \left(\bar{\psi} \cdot \overleftarrow{\partial} \Psi_{\mu\nu} - \bar{\Psi}_{\mu\nu} \partial \cdot \psi \right) F^{\mu\nu}, \tag{129}
\end{aligned}$$

where we have reached the second step by performing integration by parts w.r.t. ∂_μ in the 2nd term of the first step, and dropping Δ -exact terms containing $\partial_\mu \Psi^{\mu\nu}$. In the 2nd term of the second step, one can write $\Psi_{\mu\nu} = 2\partial_{[\mu} \psi_{\nu]}$, and integrate by parts to obtain, among others, Δ -exact terms containing $\partial_\mu F^{\mu\nu}$, which can be dropped. The result is

$$V_{\text{ST}}^{(3)} \approx - \left(\bar{\psi}^\alpha \partial_\alpha \partial_\mu \psi_\nu + \bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \right) F^{\mu\nu} + \frac{1}{2} \left(\bar{\psi}_\mu \partial_\nu \partial \cdot \psi + \bar{\psi} \cdot \overleftarrow{\partial} \overleftarrow{\partial}_\mu \psi_\nu \right) F^{\mu\nu}, \tag{130}$$

where we have made the rescaling $A_\mu \rightarrow -\frac{1}{4}A_\mu$, for convenience of comparison with our vertex $V^{(3)}$. One finds that both the vertices reduce in the TT gauge to

$$V_{\text{TT}}^{(3)} = - \left(\bar{\psi}^\alpha \partial_\alpha \partial_\mu \psi_\nu + \bar{\psi}_\mu \overleftarrow{\partial}_\nu \overleftarrow{\partial}_\alpha \psi^\alpha \right) F^{\mu\nu}, \quad (131)$$

which indeed is the 3-derivative on-shell vertex reported in [7]. In view of Eqn. (126) and (130), one also finds that the two vertices differ by Δ -exact terms:

$$V^{(3)} - V_{\text{ST}}^{(3)} \approx \frac{1}{2} \left[\bar{\psi}_\mu (\square \psi_\nu - \partial_\nu \partial \cdot \psi) + \left(\bar{\psi}_\mu \overleftarrow{\square} - \bar{\psi} \cdot \overleftarrow{\partial} \overleftarrow{\partial}_\mu \right) \psi_\nu \right] F^{\mu\nu} = \Delta\text{-exact}. \quad (132)$$

This shows the equivalence of the full off-shell vertices.

7.2 1—s—s Vertices: $s \geq 5/2$

For the sake of simplicity, from now on we restrict our attention to on-shell equivalence of vertices. As we already mentioned, if two vertices match in some gauge, say the TT one, they should also be off-shell equivalent. With this end in view, we read off the ST vertices [10], which would generally contain a bunch of terms to begin with even in the TT gauge. However, one can perform integrations by parts to see that actually the on-shell vertices are extremely simple, containing no more than a few non-trivial terms.

For example, one can take the 3-derivative $1 - \frac{5}{2} - \frac{5}{2}$ ST vertex in the TT gauge, and integrate by parts in order to have one derivative on each field. The result is simply

$$V_{\text{ST}}^{(3)} \sim \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_\beta F^{\mu\nu} \partial^\alpha \psi^\beta{}_\nu + \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_\rho (\partial_\beta A^\rho) \partial^\alpha \psi^{\beta\mu}, \quad (133)$$

where \sim means equivalence in the TT gauge up to an overall factor. In the 2nd term we integrate by parts to avoid derivatives on the photon field. We get

$$V_{\text{ST}}^{(3)} \sim \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_\beta F^{\mu\nu} \partial^\alpha \psi^\beta{}_\nu - \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_\beta \left(\overleftarrow{\partial} \cdot A \right) \partial^\alpha \psi^{\beta\mu}. \quad (134)$$

One can make use of the Clifford algebra to write $\overleftarrow{\partial} \cdot A = \frac{1}{2} \overleftarrow{\partial}_\rho A_\sigma (\gamma^\rho \gamma^\sigma + \gamma^\sigma \gamma^\rho)$, in the 2nd term on the right hand side of Eqn. (134), and then integrate by parts w.r.t. this derivative. Dropping some Δ -exact terms in the TT gauge, we get

$$V_{\text{ST}}^{(3)} \sim \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_\beta F^{\mu\nu} \partial^\alpha \psi^\beta{}_\nu + \frac{1}{2} \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_\beta (\partial_\rho A_\sigma) \gamma^\sigma \gamma^\rho \partial^\alpha \psi^{\beta\mu}. \quad (135)$$

Because $\partial \cdot A = 0$ in our gauge choice, we can write $(\partial_\rho A_\sigma) \gamma^\sigma \gamma^\rho = -\frac{1}{2} \mathcal{F}$, by making use of the identity $\gamma^\sigma \gamma^\rho = \eta^{\sigma\rho} - \gamma^{\sigma\rho}$. Therefore, we are left with

$$V_{\text{ST}}^{(3)} \sim \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_\beta \left(F^{\mu\nu} - \frac{1}{4} \eta^{\mu\nu} \mathcal{F} \right) \partial^\alpha \psi^\beta{}_\nu. \quad (136)$$

We would like to see how this compares with our 3-derivative $1 - \frac{5}{2} - \frac{5}{2}$ vertex,

$$V^{(3)} = \bar{\psi}_{\alpha\beta\parallel\mu}^{(1)} F^{+\mu\nu} \psi^{(1)\alpha\beta\parallel}{}_\nu. \quad (137)$$

The same steps as took us from Eqn. (100) to Eqn. (102) for the spin- $\frac{3}{2}$ case lead to

$$V^{(3)} \sim 2 \bar{\psi}_{\alpha\beta\|\mu}^{(1)} \left(F^{\mu\nu} - \frac{1}{4} \eta^{\mu\nu} F \right) \psi^{(1)\alpha\beta\|}_{\nu}. \quad (138)$$

Now, one can rewrite the fermionic 1-curl in terms of the original field. There will be terms that have at least one pair of mutually contracted derivatives: one acting on $\bar{\psi}_{\mu}$ and the other on ψ_{μ} . For such terms one can make use of the 3-box rule (114) to see that they are trivial in the TT gauge. Up to a trivial factor, one then has

$$V^{(3)} \sim \bar{\psi}_{\mu\alpha} \overleftarrow{\partial}_{\beta} \left(F^{\mu\nu} - \frac{1}{4} \eta^{\mu\nu} F \right) \partial^{\alpha} \psi^{\beta}_{\nu}. \quad (139)$$

From Eqn. (136) and (139), we see that the two vertices are indeed on-shell equivalent.

Let us move on to the 4-derivative $1 - \frac{5}{2} - \frac{5}{2}$ vertex. The ST one is found to be

$$V_{\text{ST}}^{(4)} \sim \bar{\psi}_{\mu\nu} \overleftarrow{\partial}_{\rho} \overleftarrow{\partial}_{\sigma} \not{A} \partial^{\mu} \partial^{\nu} \psi^{\rho\sigma}, \quad (140)$$

whereas our one is given by

$$V^{(4)} = \left(\bar{\Psi}_{\mu\nu|\rho\sigma} \gamma^{\mu\nu\alpha\beta\lambda} \Psi_{\alpha\beta|\rho\sigma} \right) A_{\lambda} \approx -2 \left(\bar{\Psi}_{\mu\nu|\rho\sigma} \gamma^{\lambda} \Psi^{\mu\nu|\rho\sigma} \right) A_{\lambda}. \quad (141)$$

We rewrite the curvature in terms of the spin- $\frac{5}{2}$ field. Among the resulting terms those with contracted pair(s) of derivatives are, again, trivial in the TT gauge, thanks to the 3-box rule (114). The other terms clearly add up to reproduce the expression (140). Therefore,

$$V^{(4)} \approx V_{\text{ST}}^{(4)} \sim \bar{\psi}_{\mu\nu} \overleftarrow{\partial}_{\rho} \overleftarrow{\partial}_{\sigma} \not{A} \partial^{\mu} \partial^{\nu} \psi^{\rho\sigma}. \quad (142)$$

For spin $\frac{5}{2}$, the only other vertex is the 5-derivative one. The ST one reads

$$V_{\text{ST}}^{(5)} \sim \left(\bar{\psi}_{\mu\nu} \overleftarrow{\partial}_{\rho} \overleftarrow{\partial}_{\sigma} \overleftrightarrow{\partial}^{\lambda} \partial^{\mu} \partial^{\nu} \psi^{\rho\sigma} \right) A_{\lambda}. \quad (143)$$

On the other hand, we have the 5-derivative Born-Infeld type vertex:

$$V^{(5)} = \bar{\Psi}_{\alpha\beta|\mu\rho} \Psi^{\alpha\beta|\rho}_{\nu} F^{\mu\nu} \approx \frac{1}{2} \left(\bar{\Psi}_{\mu\nu|\rho\sigma} \overleftrightarrow{\partial}^{\lambda} \Psi^{\mu\nu|\rho\sigma} \right) A_{\lambda}. \quad (144)$$

The off-shell equivalence can be understood in view of Eqn. (46)–(48), which pertain to spin $\frac{3}{2}$. In the equivalent vertex, again, we rewrite the fermionic curvature in terms of the spin- $\frac{5}{2}$ field, and massage the resulting terms the same way as was done for $V^{(4)}$. Thus, up to overall factors, we reproduce on-shell (143), so that

$$V^{(5)} \approx V_{\text{ST}}^{(5)} \sim \left(\bar{\psi}_{\mu\nu} \overleftarrow{\partial}_{\rho} \overleftarrow{\partial}_{\sigma} \overleftrightarrow{\partial}^{\lambda} \partial^{\mu} \partial^{\nu} \psi^{\rho\sigma} \right) A_{\lambda}. \quad (145)$$

For arbitrary spin, $s = n + \frac{1}{2}$, the story is very similar, and there are no further complications other than cluttering of indices. One can write down the ST vertices in the TT gauge from Eqn. (A.16) of [10]. They turn out to be

$$V_{\text{ST}}^{(2n-1)} \sim \bar{\psi}_{\mu\alpha_1\dots\alpha_{n-1}} \overleftarrow{\partial}_{\beta_1}\dots\overleftarrow{\partial}_{\beta_{n-1}} F^{\mu\nu} \partial^{\alpha_1}\dots\partial^{\alpha_{n-1}} \psi^{\beta_1\dots\beta_{n-1}}{}_{\nu} - \bar{\psi}_{\mu\alpha_1\dots\alpha_{n-1}} \overleftarrow{\partial}_{\beta_1}\dots\overleftarrow{\partial}_{\beta_{n-1}} \left(\overleftarrow{\partial} \cdot A \right) \partial^{\alpha_1}\dots\partial^{\alpha_{n-1}} \psi^{\beta_1\dots\beta_{n-1}}{}_{\mu}, \quad (146)$$

$$V_{\text{ST}}^{(2n)} \sim \bar{\psi}_{\mu_1\dots\mu_n} \overleftarrow{\partial}_{\nu_1}\dots\overleftarrow{\partial}_{\nu_n} A \partial^{\mu_1}\dots\partial^{\mu_n} \psi^{\nu_1\dots\nu_n}, \quad (147)$$

$$V_{\text{ST}}^{(2n+1)} \sim \left(\bar{\psi}_{\mu_1\dots\mu_n} \overleftarrow{\partial}_{\nu_1}\dots\overleftarrow{\partial}_{\nu_n} \overleftrightarrow{\partial}^{\lambda} \partial^{\mu_1}\dots\partial^{\mu_n} \psi^{\nu_1\dots\nu_n} \right) A_{\lambda}. \quad (148)$$

Their similarity with the lower-spin counterparts is obvious. Indeed, setting $n = 2$ produces exactly the respective $1 - \frac{5}{2} - \frac{5}{2}$ vertices given in Eqn. (134), (140) and (143). One can massage the $(2n - 1)$ -derivative vertex, in particular, the same way as its spin- $\frac{5}{2}$ counterpart to obtain an arbitrary-spin generalization of Eqn. (136), namely

$$V_{\text{ST}}^{(2n-1)} \sim \bar{\psi}_{\mu\alpha_1\dots\alpha_{n-1}} \overleftarrow{\partial}_{\beta_1}\dots\overleftarrow{\partial}_{\beta_{n-1}} \left(F^{\mu\nu} - \frac{1}{4} \eta^{\mu\nu} F \right) \partial^{\alpha_1}\dots\partial^{\alpha_{n-1}} \psi^{\beta_1\dots\beta_{n-1}}{}_{\nu}. \quad (149)$$

Our arbitrary-spin vertices are also straightforward generalizations of their lower-spin examples. In view of the spin- $\frac{5}{2}$ counterparts, Eqn. (137), (141) and (144), one can write

$$V^{(2n-1)} \approx \bar{\psi}_{\alpha_1\beta_1|\dots|\alpha_{n-1}\beta_{n-1}} \overleftarrow{\partial}_{\mu} F^{+\mu\nu} \psi^{(n-1)\alpha_1\beta_1|\dots|\alpha_{n-1}\beta_{n-1}}{}_{\nu}, \quad (150)$$

$$V^{(2n)} \approx \left(\bar{\Psi}_{\mu_1\nu_1|\dots|\mu_n\nu_n} \gamma^{\lambda} \Psi^{\mu_1\nu_1|\dots|\mu_n\nu_n} \right) A_{\lambda}, \quad (151)$$

$$V^{(2n+1)} \approx \left(\bar{\Psi}_{\mu_1\nu_1|\dots|\mu_n\nu_n} \overleftrightarrow{\partial}^{\lambda} \Psi^{\mu_1\nu_1|\dots|\mu_n\nu_n} \right) A_{\lambda}. \quad (152)$$

Again, one can use the 2nd identity in (101) to rewrite the $F^{+\mu\nu}$ in the first vertex, and express the fermionic $(n - 1)$ - and n -curls in all the vertices (150)–(152) in terms of the original field. The terms with contracted pair(s) of derivatives are, as usual, subject to the 3-box rule (114), and hence trivial in the TT gauge. One finds that our vertices indeed reduce on shell respectively to (149), (147) and (148). This proves the on-shell (and therefore off-shell) equivalence of the $1 - s - s$ vertices:

$$V^{(p)} \sim V_{\text{ST}}^{(p)}, \quad p = 2n - 1, 2n, 2n + 1. \quad (153)$$

8 Second-Order Consistency

We recall that consistent second-order deformation requires (S_1, S_1) to be s -exact:

$$(S_1, S_1) = -2sS_2 = -2\Delta S_2 - 2\Gamma S_2. \quad (154)$$

For abelian vertices, this antibracket is zero, so that the first-order deformations always go unobstructed. Non-abelian vertices, however, are more interesting in this respect.

We can see that there is obstruction for the non-abelian vertices we have obtained, which do not obey Eqn. (154). We prove our claim by contradiction. If Eqn. (154) holds, then the most general form of the antibracket evaluated at zero antifields is

$$[(S_1, S_1)]_{\Phi_A^*=0} = \Delta M + \Gamma N, \quad (155)$$

where $M = -2[S_2]_{C_\alpha^*=0}$ and $N = -2[S_2]_{\Phi_A^*=0}$. Note that M is obtained by setting to zero only the antighosts in S_2 . Furthermore, the equality (155) holds precisely because S_2 is linear in the antifields. The Γ -variation of (155) is therefore Δ -exact:

$$\Gamma [(S_1, S_1)]_{\Phi_A^*=0} = \Gamma \Delta M = -\Delta (\Gamma M). \quad (156)$$

It is relatively easier to compute the left hand side of (156) for our non-abelian vertices. For spin $\frac{3}{2}$, we recall that

$$a_2 = -C^* \bar{\xi} \xi, \quad a_1 = A^{*\mu} (\bar{\psi}_\mu \xi - \bar{\xi} \psi_\mu) + \tilde{a}_1, \quad a_0 = \bar{\psi}_\mu F^{+\mu\nu} \psi_\nu, \quad (157)$$

$$\tilde{a}_1 = i \left[\bar{\psi}^{*\mu} \gamma^\nu F_{\mu\nu} - \frac{1}{2(D-2)} \bar{\psi}^* \not{F} \right] \xi + \text{h.c.} \quad (158)$$

To compute the antibracket of $S_1 = \int (a_2 + a_1 + a_0)$ with itself, we notice that a field-antifield pair shows up only in $\int a_1$, and between $\int a_0$ and $\int a_1$, so that it reduces to

$$(S_1, S_1) = 2 \left(\int a_0, \int a_1 \right) + \left(\int a_1, \int a_1 \right). \quad (159)$$

Now, the second antibracket on the right hand side necessarily contains antifields, while the first one will not contain any. Thus we have

$$[(S_1, S_1)]_{\Phi_A^*=0} = 2 \left(\int a_0, \int a_1 \right). \quad (160)$$

Notice that, while the unambiguous piece in a_1 contains the antifield $A^{*\mu}$, the ambiguity, \tilde{a}_1 , contains instead the antifield $\bar{\psi}^{*\mu}$. Correspondingly, $(\int a_0, \int a_1)$ will contain two distinct kinds of pieces: 4-Fermion terms and Fermion bilinears. Explicitly,

$$\begin{aligned} [(S_1, S_1)]_{\Phi_A^*=0} &= \int d^D x \left\{ 4(\bar{\psi}_\mu \xi - \bar{\xi} \psi_\mu) \partial_\nu (\bar{\psi}^{[\mu} \psi^{\nu]}) + \frac{1}{2} \bar{\psi}_\alpha \gamma^{\mu\nu\alpha\beta} \psi_\beta \right\} \\ &+ \int d^D x \left\{ i \bar{\psi}_\mu F^{+\mu\nu} \left[2\gamma^\rho F_{\nu\rho} - \frac{1}{(D-2)} \gamma_\nu \not{F} \right] \xi + \text{h.c.} \right\}. \end{aligned} \quad (161)$$

If the vertex is unobstructed and Eqn. (156) holds, then the Γ -variation of each of these terms should independently be Δ -exact. Let us consider the Fermion bilinears appearing in the second line of Eqn. (161), originating from $(\int a_0, \int \tilde{a}_1)$. It is easy to see that their Γ -variation is not Δ -exact. We conclude that the non-abelian $1 - \frac{3}{2} - \frac{3}{2}$ vertex gets obstructed beyond the cubic order. The proof for arbitrary spin will be very similar.

Notice that the vertex $\bar{\psi}_\mu F^{+\mu\nu} \psi_\nu$ is precisely the Pauli term appearing in $\mathcal{N} = 2$ SUGRA [3]. The theory, however, contains additional degrees of freedom, namely graviton, on top of a complex massless spin $\frac{3}{2}$ and a $U(1)$ field. It is this new dynamical field that renders the vertex unobstructed, while keeping locality intact. If one decouples gravity by taking $M_{\text{P}} \rightarrow \infty$, the Pauli term vanishes because the dimensionful coupling constant is nothing but $1/M_{\text{P}}$ [3]. One could integrate out the massless graviton to obtain a system of spin- $\frac{3}{2}$ and spin-1 fields only. The resulting theory contains the Pauli term, but is necessarily non-local. Thus, higher-order consistency of the non-abelian vertex is possible either by forgoing locality or by adding a new dynamical field (graviton).

9 Remarks & Future Perspectives

In this paper, we have employed the BRST-BV cohomological methods to construct consistent parity-preserving off-shell cubic vertices for fermionic gauge fields coupled to EM in flat space. We have shown that consistency and non-triviality of the deformations forbid minimal coupling, and pose number-of-derivative restrictions on a $1 - s - s$ vertex, in accordance with Metsaev's light-cone-formulation results [7].

The vertices either deform the gauge algebra or, when they do not deform the gauge algebra, turn out not to deform the gauge transformations either and to deform only the Lagrangian. The non-abelian ones get obstructed in a local theory beyond the cubic order in the absence of additional higher-spin gauge fields.

Our off-shell cubic vertices are equivalent to the string-theory-inspired ones of Sagnotti-Taronna [10]. Note that in [10] there appears just one dimensionful coupling constant, which can be set to unity. Then, each cubic vertex will come with a fixed known numerical coefficient. This is apparently in contrast with our results, where each of the three $1 - s - s$ vertices has an independent coupling constant. However, it is well known that higher-order consistency requirements may impose restrictions on the cubic couplings by relating them with one another [17]. Because the consistency of string theory is not limited to the cubic order, then it should not come as a surprise that the cubic vertices it gives rise to have no freedom in the coupling constant. At any rate, string theory may not be the unique consistent theory of higher-spin fields. If this is true, other possible choices of the cubic couplings would pertain to other consistent theories.

The number of possible $1 - s - s$ vertices for fermions differ from that for bosons. While in both cases there is only one non-abelian vertex, fermions have, beside the usual Born-Infeld type 3-curvature term, another abelian vertex in $D \geq 5$, which is gauge invariant up to a total derivative. In this respect, fermionic $1 - s - s$ vertices are strikingly similar in nature to the bosonic $2 - s - s$ ones. The latter include one non-abelian $(2s - 2)$ -derivative

vertex, a $2s$ -derivative abelian one that is gauge invariant up to a total derivative and exists in $D \geq 5$, and a Born-Infeld type abelian one containing $2s + 2$ derivatives. This could be seen by using either the light-cone method [6] or the cohomological methods [4, 5].

For gravitational coupling, spin $\frac{3}{2}$ has no consistency issues, but consistent deformations of the free theory uniquely lead one to $\mathcal{N} = 1$ SUGRA [19], under certain reasonable assumptions. Fermionic gauge fields with higher spin, $s \geq \frac{5}{2}$, and their coupling to gravity are more interesting, for which one can also employ the BRST-deformation technique [20]. Another interesting avenue to pursue are the mixed-symmetry fields.

It is instructive to consider the EM coupling of massive higher-spin fields in flat space, which has been discussed by various authors⁹. If Lorentz, parity and time-reversal symmetries hold good, a massive spin- s particle will have $2s + 1$ EM multipoles [23]. This immediately sets for the possible number of derivatives in a $1 - s - s$ vertex an upper bound, which remains the same in the massless limit. The assumption of light-cone helicity conservation in $D = 4$ uniquely determines all the multipoles [23]. However, only the highest multipole survives in an appropriate massless chargeless scaling limit. This observation is in harmony with our results, since any of our lower-derivative vertices either vanishes in 4D or is not consistent by itself in a local theory.

On the other hand, causal propagation of a charged massive field may call for certain non-minimal terms. Indeed, for a massive spin $\frac{3}{2}$ in flat space, causality analysis in a constant external EM background [24] or in the case of $\mathcal{N} = 2$ broken SUGRA [25] reveals the crucial role played by the Pauli term, $\bar{\psi}_\mu F^{+\mu\nu} \psi_\nu$. In the massless case, as we have seen, the same term arises as the unique non-abelian deformation of the gauge theory. These facts go in favor of the gauge-invariant (Stückelberg-invariant) formulation, adopted in [22], for constructing consistent EM interactions of massive higher spins.

Our non-abelian vertices are seen to be inconsistent beyond the cubic order in a local theory. Such obstructions are rather common for massless higher-spin vertices in flat space, and some could not even be removed by the inclusion of an (in)finite number of higher-spin fields, as has been argued in [26]. Non-locality may therefore be essential. In fact, as noticed in [27], evidence for non-locality shows up already at the quartic level. The geometric formulation of free massless higher spins also hints towards the same, as they generically yield non-local EoMs [28] if higher-derivative terms are not considered [29].

If one has to give up locality, what becomes relevant for studying higher-spin interactions is a formulation that does not require locality as an input, e.g., the old S-matrix theory, or perhaps the more powerful BCFW construction [30] and generalizations thereof. The latter seem promising for the systematic search of consistent interactions of massless higher-spin particles in 4D Minkowski space [31].

⁹See, for example, Ref. [21, 22, 23, 24] and references therein.

There are certain technical difficulties in extending the applicability of the BRST-BV cohomological methods to constant curvature spaces. For AdS space, in particular, those could be avoided by using the ambient-space formulation [32]. If so, one would be able to construct off-shell vertices for AdS, and compare them with the recently-obtained results of [33, 34]. This would be one step towards finding a standard action for the Vasiliev systems [35], which are a consistent set of non-linear equations for symmetric tensors of arbitrary rank in any number of dimensions. We leave this as future work.

Acknowledgments

We would like to thank N. Boulanger, D. Francia, E. Joung, K. Mkrtchyan, M. Porrati and M. Taronna for useful discussions. MH and RR are grateful to The Erwin Schrödinger Institute, Vienna for its kind hospitality, during the Workshop on Higher Spin Gravity, while this work was ongoing. MH gratefully acknowledges support from the Alexander von Humboldt Foundation through a Humboldt Research Award and support from the ERC through the “SyDuGraM” Advanced Grant. RR is a Postdoctoral Fellow of the Fonds de la Recherche Scientifique-FNRS. This work is also partially supported by IISN - Belgium (conventions 4.4511.06 and 4.4514.08) and by the “Communauté Française de Belgique” through the ARC program.

A Curvatures & Equations of Motion

Let us recall that for arbitrary spin $s = n + \frac{1}{2}$, we have a totally symmetric rank- n tensor-spinor $\psi_{\mu_1 \dots \mu_n}$, whose curvature is its n -curl, i.e., the rank- $2n$ tensor

$$\Psi_{\mu_1 \nu_1 | \mu_2 \nu_2 | \dots | \mu_n \nu_n} = [\dots [[\partial_{\mu_1} \dots \partial_{\mu_n} \psi_{\nu_1 \dots \nu_n} - (\mu_1 \leftrightarrow \nu_1)] - (\mu_2 \leftrightarrow \nu_2)] \dots] - (\mu_n \leftrightarrow \nu_n). \quad (\text{A.1})$$

Notice that, unlike the Fronsdal tensor, $\mathcal{S}_{\mu_1 \dots \mu_n}$, the curvature tensor (A.1) is gauge invariant even for an unconstrained gauge parameter. Its properties can be found in [15]. The curvature is antisymmetric under the interchange of “paired” indices, e.g.,

$$\Psi_{\mu_1 \nu_1 | \mu_2 \nu_2 | \dots | \mu_n \nu_n} = -\Psi_{\nu_1 \mu_1 | \mu_2 \nu_2 | \dots | \mu_n \nu_n}, \quad (\text{A.2})$$

but symmetric under the interchange of any two sets of paired indices, e.g.,

$$\Psi_{\mu_1 \nu_1 | \mu_2 \nu_2 | \dots | \mu_{n-1} \nu_{n-1} | \mu_n \nu_n} = \Psi_{\mu_n \nu_n | \mu_2 \nu_2 | \dots | \mu_{n-1} \nu_{n-1} | \mu_1 \nu_1}. \quad (\text{A.3})$$

These symmetries actually hold good for any m -curl of the field, $m \leq n$. Another important property of the curvature is that it obeys the Bianchi identity

$$\partial_{[\rho} \Psi_{\mu_1 \nu_1 | \mu_2 \nu_2 | \dots | \mu_n \nu_n} = 0. \quad (\text{A.4})$$

The (Weinberg) curvature (A.1) and the EM field strength $F_{\mu\nu}$ are useful in casting the EoMs into a variety of forms, which, among others, can help one identify Δ -exact pieces. First, we write down these various forms for the photon field. Next, we do the same for spin $\frac{3}{2}$, explaining as well how to derive them and writing them explicitly as Δ -variations. Then we move on to spin $\frac{5}{2}$, and finally to arbitrary spin.

A.1 The Photon

The original photon EoMs are given by

$$\partial^\mu F_{\mu\nu} = \square A_\nu - \partial_\nu(\partial \cdot A) = \Delta A_\nu^*. \quad (\text{A.5})$$

One can take its 1-curl to obtain

$$\square F_{\mu\nu} = 2\Delta(\partial_{[\mu} A_{\nu]}^*). \quad (\text{A.6})$$

A.2 Spin 3/2

For spin $\frac{3}{2}$, the original EoMs can be obtained directly from the master action (24)

$$\gamma^{\mu\alpha\beta}\Psi_{\alpha\beta} = -2i\Delta\psi^{*\mu}, \quad (\text{A.7a})$$

$$\bar{\Psi}_{\alpha\beta}\gamma^{\alpha\beta\mu} = 2i\Delta\bar{\psi}^{*\mu}. \quad (\text{A.7b})$$

One can take the γ -trace of Eqn. (A.7a), and use $\gamma_\mu\gamma^{\mu\alpha\beta} = (D-2)\gamma^{\alpha\beta}$, to obtain

$$\gamma^{\mu\nu}\Psi_{\mu\nu} = 2(\not{\partial}\psi - \partial \cdot \psi) = -2i\Delta\left(\frac{1}{D-2}\not{\psi}^*\right). \quad (\text{A.8})$$

Now, in Eqn. (A.7a), one can use the identity $\gamma^{\mu\alpha\beta} = \gamma^\mu\gamma^{\alpha\beta} - 2\eta^{\mu[\alpha}\gamma^{\beta]}$, and then the EoM. (A.8), to obtain another very useful form

$$\gamma^\mu\Psi_{\mu\nu} = \not{\partial}\psi_\nu - \partial_\nu\psi = -i\Delta\left(\psi_\nu^* - \frac{1}{D-2}\gamma_\nu\not{\psi}^*\right). \quad (\text{A.9})$$

One can take a curl of the above equation to get

$$\not{\partial}\Psi_{\mu\nu} = -2i\Delta\left(\partial_{[\mu}\psi_{\nu]}^* - \frac{1}{D-2}\gamma_{[\nu}\partial_{\mu]}\not{\psi}^*\right). \quad (\text{A.10})$$

Another useful form can be obtained by applying the Dirac operator on (A.9), and then getting rid of $\not{\partial}\psi$ in the resulting expression by using (A.8). The result is

$$\partial^\mu\Psi_{\mu\nu} = \square\psi_\nu - \partial_\nu(\partial \cdot \psi) = -i\Delta\left[\not{\partial}\psi_\nu^* + \frac{1}{D-2}\gamma_{\nu\rho}\partial^\rho\not{\psi}^*\right]. \quad (\text{A.11})$$

Similarly, one could have started with (A.7b) to derive the following.

$$\bar{\Psi}_{\mu\nu}\gamma^{\mu\nu} = 2\left(\bar{\psi}\cdot\overleftarrow{\partial}-\bar{\psi}\overleftarrow{\partial}\right) = 2i\Delta\left(\frac{1}{D-2}\bar{\psi}^*\right), \quad (\text{A.12})$$

$$\bar{\Psi}_{\mu\nu}\gamma^\nu = \bar{\psi}\overleftarrow{\partial}_\mu - \bar{\psi}_\mu\overleftarrow{\partial} = i\Delta\left(\bar{\psi}_\mu^* - \frac{1}{D-2}\bar{\psi}^*\gamma_\mu\right), \quad (\text{A.13})$$

$$\bar{\Psi}_{\mu\nu}\overleftarrow{\partial} = 2i\Delta\left(\bar{\psi}_{[\mu}^*\overleftarrow{\partial}_{\nu]} - \frac{1}{D-2}\bar{\psi}^*\gamma_{[\mu}\overleftarrow{\partial}_{\nu]}\right), \quad (\text{A.14})$$

$$\bar{\Psi}_{\mu\nu}\overleftarrow{\partial}^\nu = \left(\bar{\psi}\cdot\overleftarrow{\partial}\right)\overleftarrow{\partial}_\mu - \bar{\psi}_\mu\overleftarrow{\partial} = i\Delta\left[\bar{\psi}_\mu^*\overleftarrow{\partial} + \frac{1}{D-2}\bar{\psi}^*\overleftarrow{\partial}^\rho\gamma_{\rho\mu}\right]. \quad (\text{A.15})$$

A.3 Spin 5/2

For the spin- $\frac{5}{2}$ case, let us recall from Section 4 that the original EoMs are given by

$$\mathcal{R}_{\mu\nu} = \mathcal{S}_{\mu\nu} - \gamma_{(\mu}\mathcal{F}_{\nu)} - \frac{1}{2}\eta_{\mu\nu}\mathcal{S}' = \Delta\psi_{\mu\nu}^*, \quad (\text{A.16a})$$

$$\bar{\mathcal{R}}_{\mu\nu} = \bar{\mathcal{S}}_{\mu\nu} - \bar{\mathcal{F}}_{(\mu}\gamma_{\nu)} - \frac{1}{2}\eta_{\mu\nu}\bar{\mathcal{S}}' = \Delta\bar{\psi}_{\mu\nu}^*, \quad (\text{A.16b})$$

which one can easily rewrite in terms of the Fronsdal tensor,

$$\mathcal{S}_{\nu_1\nu_2} = i\left[\not{\partial}\psi_{\nu_1\nu_2} - 2\partial_{(\nu_1}\psi_{\nu_2)}\right] = \Delta\left[\psi_{\nu_1\nu_2}^* - \frac{2}{D}\gamma_{(\nu_1}\psi_{\nu_2)}^* - \frac{1}{D}\eta_{\nu_1\nu_2}\psi^{*\prime}\right], \quad (\text{A.17a})$$

$$\bar{\mathcal{S}}_{\nu_1\nu_2} = i\left[\bar{\psi}_{\nu_1\nu_2}\overleftarrow{\not{\partial}} - 2\bar{\psi}_{(\nu_1}\overleftarrow{\partial}_{\nu_2)}\right] = \Delta\left[\bar{\psi}_{\nu_1\nu_2}^* - \frac{2}{D}\bar{\psi}_{(\nu_1}^*\gamma_{\nu_2)} - \frac{1}{D}\eta_{\nu_1\nu_2}\bar{\psi}^{*\prime}\right]. \quad (\text{A.17b})$$

Now we see that the quantity $\gamma^{\mu_1}\Psi_{\mu_1\nu_1|\mu_2\nu_2}$ is given by the 1-curl of the Fronsdal tensor, so that it is Δ -exact as a result of Eqn. (A.17a):

$$\begin{aligned} \gamma^{\mu_1}\Psi_{\mu_1\nu_1|\mu_2\nu_2} &= \not{\partial}\psi_{\mu_2\nu_2|\nu_1}^{(1)} - \partial_{\nu_1}\psi_{\mu_2\nu_2}^{(1)} = -i\mathcal{S}_{\mu_2\nu_2|\nu_1}^{(1)} \\ &= -i\Delta\left[\psi_{\mu_2\nu_2|\nu_1}^{*(1)} - \frac{1}{D}\gamma_{\nu_1}\psi_{\mu_2\nu_2}^{*(1)} + \frac{2}{D}\gamma_{[\mu_2}\partial_{\nu_2]}\psi_{\nu_1}^* + \frac{2}{D}\eta_{\nu_1[\mu_2}\partial_{\nu_2]}\psi^{*\prime}\right]. \end{aligned} \quad (\text{A.18})$$

Contracting this expression with γ^{ν_1} on the left, we obtain another useful form,

$$\gamma^{\mu_1\nu_1}\Psi_{\mu_1\nu_1|\mu_2\nu_2} = 2\left[\not{\partial}\psi_{\mu_2\nu_2}^{(1)} - \partial^\rho\psi_{\mu_2\nu_2|\rho}^{(1)}\right] = i\mathcal{F}_{\mu_2\nu_2}^{(1)} = -2i\Delta\left[\frac{1}{D}\psi_{\mu_2\nu_2}^{*(1)}\right]. \quad (\text{A.19})$$

One finds that taking a curl of (A.18) gives yet another form,

$$\begin{aligned} \not{\partial}\Psi_{\mu_1\nu_1|\mu_2\nu_2} &= -i\Delta\left[\frac{1}{2}\psi_{\mu_1\nu_1|\mu_2\nu_2}^{*(2)} + \frac{2}{D}\gamma_{[\mu_1}\partial_{\nu_1]}\psi_{\mu_2\nu_2}^{*(1)} + \frac{2}{D}\partial_{[\mu_1}\eta_{\nu_1][\mu_2}\partial_{\nu_2]}\psi^{*\prime} + (\mu_1\nu_1 \leftrightarrow \mu_2\nu_2)\right] \\ &= -i\mathcal{S}_{\mu_1\nu_1|\mu_2\nu_2}^{(2)}. \end{aligned} \quad (\text{A.20})$$

Given Eqn. (A.18) and (A.19), one can also write

$$\begin{aligned} \partial^{\mu_1}\Psi_{\mu_1\nu_1|\mu_2\nu_2} &= -i\Delta\left[\not{\partial}\left(\psi_{\mu_2\nu_2|\nu_1}^{*(1)} + \frac{2}{D}\gamma_{[\mu_2}\partial_{\nu_2]}\psi_{\nu_1}^* + \frac{2}{D}\eta_{\nu_1[\mu_2}\partial_{\nu_2]}\psi^{*\prime}\right) + \frac{1}{D}\gamma_{\nu_1\rho}\partial^\rho\psi_{\mu_2\nu_2}^{*(1)}\right] \\ &= -i\not{\partial}\mathcal{S}_{\mu_2\nu_2|\nu_1}^{(1)} + \frac{i}{2}\partial_{\nu_1}\mathcal{F}_{\mu_2\nu_2}^{(1)}. \end{aligned} \quad (\text{A.21})$$

Similarly, Eqn. (A.17b) gives the various forms of the EoMs for the Dirac conjugate spinor.

A.4 Arbitrary Spin

For arbitrary spin $s = n + \frac{1}{2}$, we recall from Section 5 that the original EoMs read

$$\mathcal{R}_{\mu_1 \dots \mu_n} = \mathcal{S}_{\mu_1 \dots \mu_n} - \frac{1}{2}n \gamma_{(\mu_1} \mathcal{S}_{\mu_2 \dots \mu_n)} - \frac{1}{4}n(n-1) \eta_{(\mu_1 \mu_2} \mathcal{S}'_{\mu_3 \dots \mu_n)} = \Delta \psi_{\mu_1 \dots \mu_n}^*, \quad (\text{A.22a})$$

$$\bar{\mathcal{R}}_{\mu_1 \dots \mu_n} = \bar{\mathcal{S}}_{\mu_1 \dots \mu_n} - \frac{1}{2}n \bar{\mathcal{S}}_{(\mu_1 \dots \mu_{n-1}} \gamma_{\mu_n)} - \frac{1}{4}n(n-1) \eta_{(\mu_1 \mu_2} \bar{\mathcal{S}}'_{\mu_3 \dots \mu_n)} = \Delta \bar{\psi}_{\mu_1 \dots \mu_n}^*. \quad (\text{A.22b})$$

One can reexpress the EoMs in terms of the Fronsdal tensor as follows.

$$\mathcal{S}_{\nu_1 \dots \nu_n} = \Delta \left[\psi_{\nu_1 \dots \nu_n}^* - \frac{n}{2n+D-4} \gamma_{(\nu_1} \not{\psi}_{\nu_2 \dots \nu_n)}^* - \frac{n(n-1)}{2(n+D-2)} \eta_{(\nu_1 \nu_2} \psi_{\nu_3 \dots \nu_n)}^{*'} \right], \quad (\text{A.23a})$$

$$\bar{\mathcal{S}}_{\nu_1 \dots \nu_n} = \Delta \left[\bar{\psi}_{\nu_1 \dots \nu_n}^* - \frac{n}{2n+D-4} \bar{\not{\psi}}_{(\nu_1 \dots \nu_{n-1}}^* \gamma_{\nu_n)} - \frac{n(n-1)}{2(n+D-2)} \eta_{(\nu_1 \nu_2} \bar{\psi}_{\nu_3 \dots \nu_n)}^{*'} \right]. \quad (\text{A.23b})$$

From the definition (1) of the Fronsdal tensor, it is easy to see that

$$\gamma^{\mu_1} \Psi_{\mu_1 \nu_1 | \dots | \mu_n \nu_n} = -i \mathcal{S}_{\mu_2 \nu_2 | \dots | \mu_n \nu_n}^{(n-1)}, \quad (\text{A.24})$$

whose contraction with γ^{ν_1} on the left gives

$$\gamma^{\mu_1 \nu_1} \Psi_{\mu_1 \nu_1 | \dots | \mu_n \nu_n} = i \mathcal{S}_{\mu_2 \nu_2 | \dots | \mu_n \nu_n}^{(n-1)}. \quad (\text{A.25})$$

Also, a curl of Eqn. (A.24) yields

$$\not{\partial} \Psi_{\mu_1 \nu_1 | \dots | \mu_n \nu_n} = -i \mathcal{S}_{\mu_1 \nu_1 | \dots | \mu_n \nu_n}^{(n)}. \quad (\text{A.26})$$

Finally, from Eqn. (A.24) and (A.25) one obtains

$$\partial^{\mu_1} \Psi_{\mu_1 \nu_1 | \dots | \mu_n \nu_n} = -i \not{\partial} \mathcal{S}_{\mu_2 \nu_2 | \dots | \mu_n \nu_n}^{(n-1)} + \frac{i}{2} \partial_{\nu_1} \mathcal{S}_{\mu_2 \nu_2 | \dots | \mu_n \nu_n}^{(n-1)}. \quad (\text{A.27})$$

In view of Eqn. (A.23a), it is now straightforward to write the EoMs (A.24)–(A.27) as Δ -exact terms. Similar things follow from Eqn. (A.23b) for the Dirac conjugate spinor.

B The Cohomology of Γ

This Appendix is devoted to clarifying and providing proofs of the statements about the cohomology of Γ appearing in Section 2.1. We recall that the action of Γ is defined by

$$\Gamma A_\mu = \partial_\mu C, \quad (\text{B.1a})$$

$$\Gamma \psi_{\nu_1 \dots \nu_n} = n \partial_{(\nu_1} \xi_{\nu_2 \dots \nu_n)}. \quad (\text{B.1b})$$

Note that the non-trivial elements in the cohomology of Γ are nothing but gauge-invariant objects that themselves are not gauge variations of something else. Here we consider one by one all such elements enlisted in Section 2.1. In the process, we also prove the statements made towards the end of Section 2.1 about some Γ -exact terms.

B.1 The Curvatures

The curvatures $\{F_{\mu\nu}, \Psi_{\mu_1\nu_1|\dots|\mu_n\nu_n}\}$ and their derivatives belong to the cohomology of Γ . Seeing that the curvatures are Γ -closed is straightforward. For the photon it follows directly from the commutativity of partial derivatives as one takes a curl of Eqn. (B.1a),

$$\Gamma F_{\mu\nu} = \Gamma (2\partial_{[\mu}A_{\nu]}) = 2\partial_{[\mu}\partial_{\nu]}C = 0. \quad (\text{B.2})$$

On the other hand, taking a 1-curl of Eqn. (B.1b) one obtains

$$\Gamma \psi^{(1)\mu_1\nu_1\|\nu_2\dots\nu_n} = (n-1)\partial_{(\nu_2}\xi^{(1)\mu_1\nu_1\|\nu_3\dots\nu_n)}. \quad (\text{B.3})$$

It is easy to see that, in general, an m -curl of Eqn. (B.1b) gives, for $m \leq n$,

$$\Gamma \psi^{(m)\mu_1\nu_1|\dots|\mu_m\nu_m\|\nu_{m+1}\dots\nu_n} = (n-m)\partial_{(\nu_{m+1}}\xi^{(m)\mu_1\nu_1|\dots|\mu_m\nu_m\|\nu_{m+2}\dots\nu_n)}. \quad (\text{B.4})$$

In particular, when $m = n$, we have the Γ -variation of the curvature, which vanishes:

$$\Gamma \Psi^{\mu_1\nu_1|\dots|\mu_n\nu_n} = 0. \quad (\text{B.5})$$

Notice that the Γ -closedness of the curvature does not require any constraints on the fermionic ghost. That the curvatures are not Γ -exact simply follows from the fact that these are $pg\hbar=0$ objects, whereas any Γ -exact piece must have $pg\hbar > 0$. Therefore, the curvatures are in the cohomology of Γ , and so are their derivatives.

We have seen that only the highest curl (n -curl) of the spinor $\psi_{\nu_1\dots\nu_n}$ is Γ -closed, while the lower curls are not. The key point is the commutativity of partial derivatives, and clearly, any arbitrary derivative of the field will not be Γ -closed in general. However, some particular linear combination of such objects (or γ -traces thereof) can be Γ -closed under the constrained ghost. The latter possibility is exhausted by the Fronsdal tensor and its derivatives, which we will discuss later.

B.2 The Antifields

The antifields $\{A^{*\mu}, C^*, \bar{\psi}^{*\mu_1\dots\mu_n}, \bar{\xi}^{*\mu_1\dots\mu_{n-1}}\}$ and their derivatives also belong to the cohomology of Γ . Clearly, these objects are Γ -closed since Γ does not act on the antifields, while they cannot be Γ -exact because they have $pg\hbar = 0$.

B.3 The Ghosts & Curls Thereof

The undifferentiated ghosts $\{C, \xi_{\mu_1\dots\mu_{n-1}}\}$ are Γ -closed objects simply because Γ does not act on them. Also they cannot be Γ -exact, thanks to Eqn. (B.1), which tells us that any Γ -exact piece must contain at least one derivative of any of the ghosts.

Any derivatives of the ghosts will also be Γ -closed. Some derivatives, however, will be Γ -exact, and therefore trivial in the cohomology of Γ . One can immediately dismiss as trivial any derivative of the bosonic ghost C , because $\partial_\mu C = \Gamma A_\mu$ from Eqn. (B.1a).

Derivatives of the fermionic ghost¹⁰ are more subtle. One can show that only the γ -traceless part of the curls of the ghost $\{\xi_{\mu_1\nu_1|\dots|\mu_m\nu_m\|\nu_{m+1}\dots\nu_{n-1}}, m \leq n-1\}$ are non-trivial elements in the cohomology of Γ . First, one can convince oneself step by step why only the ghost-curls are interesting. In the simplest non-trivial case of $n=2$, we see that

$$\partial_\mu \xi_\nu = \partial_{(\mu} \xi_{\nu)} + \partial_{[\mu} \xi_{\nu]} = \frac{1}{2} \Gamma \psi_{\mu\nu} + \frac{1}{2} \xi_{\mu\nu}^{(1)}. \quad (\text{B.6})$$

For $n=3$, we find

$$\partial_\mu \xi_{\nu\rho} = \partial_{(\mu} \xi_{\nu\rho)} + \frac{4}{3} \partial_{[\mu} \xi_{\nu]\rho} + \frac{2}{3} \partial_{[\nu} \xi_{\rho]\mu} = \frac{1}{3} \Gamma \psi_{\mu\nu\rho} + \frac{2}{3} \xi_{\mu\nu\|\rho}^{(1)} + \frac{1}{3} \xi_{\nu\rho\|\mu}^{(1)}, \quad (\text{B.7})$$

It is easy to generalize this to the arbitrary spin case, for which we obtain

$$\begin{aligned} \partial_\rho \xi_{\nu_1 \dots \nu_{n-1}} &= \partial_{(\rho} \xi_{\nu_1 \dots \nu_{n-1})} + 2 \left(1 - \frac{1}{n}\right) \partial_{[\rho} \xi_{\nu_1] \nu_2 \dots \nu_{n-1}} \\ &\quad + 2 \sum_{m=1}^{n-2} \left(1 - \frac{m+1}{n}\right) \partial_{[\nu_m} \xi_{\nu_{m+1}] \rho \nu_1 \dots \nu_{m-1} \nu_{m+2} \dots \nu_{n-1}} \\ &= \frac{1}{n} \Gamma \psi_{\rho \nu_1 \dots \nu_{n-1}} + \left(1 - \frac{1}{n}\right) \xi_{\rho \nu_1 \|\nu_2 \dots \nu_{n-1}}^{(1)} \\ &\quad + \sum_{m=1}^{n-2} \left(1 - \frac{m+1}{n}\right) \xi_{\nu_m \nu_{m+1} \|\rho \nu_1 \dots \nu_{m-1} \nu_{m+2} \dots \nu_{n-1}}^{(1)}. \end{aligned} \quad (\text{B.8})$$

In view of Eqn. (B.6)–(B.8), we conclude that any first derivative of the ghost is a linear combination of 1-curls, up to Γ -exact terms. Therefore, in the cohomology of Γ it suffices to consider only 1-curls of the ghost. More generally, one can consider only the m -curls in the cohomology of Γ , instead of arbitrary m derivatives, where $m \leq n-1$. The latter can easily be seen by first taking a 1-curl of Eqn. (B.8), and convincing oneself that only 2-curls of the ghost are interesting. In this way, one can continue up to m -curls of Eqn. (B.8), $m \leq n-1$, to show that it suffices to consider only m -curls.

The derivative of an m -curl, $\partial_{\nu_n} \xi_{\mu_1 \nu_1 |\dots| \mu_m \nu_m \|\nu_{m+1} \dots \nu_{n-1}}^{(m)}$, contains non-trivial $(m+1)$ -curls plus trivial terms. It is clear that this quantity can be Γ -exact if and only if symmetrized w.r.t. the indices $\{\nu_{m+1}, \dots, \nu_n\}$. In this case, we have from Eqn. (B.4)

$$\partial_{(\nu_n} \xi_{\mu_1 \nu_1 |\dots| \mu_m \nu_m \|\nu_{m+1} \dots \nu_{n-1})}^{(m)} = \frac{1}{n-m} \Gamma \psi_{\mu_1 \nu_1 |\dots| \mu_m \nu_m \|\nu_{m+1} \dots \nu_n}^{(m)}, \quad 0 \leq m \leq n-1. \quad (\text{B.9})$$

It follows immediately that a derivative of the highest ghost-curl is always Γ -exact. Indeed, in Eqn. (B.9) one can set $m = n-1$, and obtain

$$\partial_{\nu_n} \xi_{\mu_1 \nu_1 |\dots| \mu_{n-1} \nu_{n-1}}^{(n-1)} = \Gamma \psi_{\mu_1 \nu_1 |\dots| \mu_{n-1} \nu_{n-1} \|\nu_n}^{(n-1)}. \quad (\text{B.10})$$

¹⁰The rest of this Appendix will deal only with the fermionic ghost $\xi_{\mu_1 \dots \mu_n}$, and without any source of confusion we will simply call it “ghost”.

Although any m -curl, $\xi_{\mu_1\nu_1|\dots|\mu_m\nu_m\|\nu_{m+1}\dots\nu_{n-1}}^{(m)}$, is in the cohomology of Γ , its γ -trace is always Γ -exact. In fact, the latter vanishes when the γ -matrix to be contracted carries one of the unpaired indices $\{\nu_{m+1}, \dots, \nu_{n-1}\}$, thanks to the γ -tracelessness of the ghost. If the index contraction is otherwise, the same constraint gives

$$\gamma^{\mu_1}\xi_{\mu_1\nu_1|\dots|\mu_m\nu_m\|\nu_{m+1}\dots\nu_{n-1}}^{(m)} = \not{\partial}\xi_{\mu_2\nu_2|\dots|\mu_m\nu_m\|\nu_1\nu_{m+1}\dots\nu_{n-1}}^{(m-1)}. \quad (\text{B.11})$$

On the other hand, one can take a γ -trace of Eqn. (B.9) to have

$$\Gamma\psi_{\mu_1\nu_1|\dots|\mu_m\nu_m\|\nu_{m+1}\dots\nu_{n-1}}^{(m)} = (n-m)\not{\partial}\xi_{\mu_2\nu_2|\dots|\mu_m\nu_m\|\nu_1\nu_{m+1}\dots\nu_{n-1}}^{(m-1)}. \quad (\text{B.12})$$

Comparing Eqn. (B.11) with (B.12), it is clear that $\gamma^{\mu_1}\xi_{\mu_1\nu_1|\dots|\mu_m\nu_m\|\nu_{m+1}\dots\nu_{n-1}}^{(m)}$ is Γ -exact. Therefore, one can exclude the γ -traces of ghost curls from the cohomology of Γ .

B.4 The Fronsdal Tensor

The Fronsdal tensor $\mathcal{S}_{\mu_1\dots\mu_n}$ and its derivatives are also in the cohomology of Γ . From the definition (1), $\mathcal{S}_{\mu_1\dots\mu_n}$ can be shown to be Γ -closed under the constrained ghost. Indeed,

$$\begin{aligned} \Gamma\mathcal{S}_{\mu_1\dots\mu_n} &= i \left[\not{\partial}\Gamma\psi_{\mu_1\dots\mu_n} - n\partial_{(\mu_1}\Gamma\psi_{\mu_2\dots\mu_n)} \right] \\ &= in \left[\not{\partial}\partial_{(\mu_1}\xi_{\mu_2\dots\mu_n)} - n\gamma^\rho\partial_{(\mu_1}\partial_{(\rho}\xi_{\mu_2\dots\mu_n)} \right] \\ &= -in(n-1)\partial_{(\mu_1}\partial_{(\mu_2}\xi_{\mu_3\dots\mu_n)}), \end{aligned} \quad (\text{B.13})$$

which vanishes if the ghost is γ -traceless. Being a $pg\hbar$ -0 object, $\mathcal{S}_{\mu_1\dots\mu_n}$ can also not be Γ -exact. So, the Fronsdal tensor and its derivatives belong to the cohomology of Γ .

However, in view of Eqn. (A.24) and (A.26), the two highest curls of the Fronsdal tensor boil down to objects we have already enlisted in Subsection B.1, and therefore need not be considered separately. These equations are generalizations of the Damour-Deser relations [36] (see also [29]). For spin $\frac{5}{2}$, in particular, they make it sufficient to consider only symmetrized derivatives of the Fronsdal tensor.

References

- [1] S. Weinberg, Phys. Rev. **135**, B1049 (1964). C. Aragone and S. Deser, Phys. Lett. B **86**, 161 (1979); Nucl. Phys. B **170**, 329 (1980). S. Weinberg and E. Witten, Phys. Lett. B **96**, 59 (1980).
- [2] M. Porrati, Phys. Rev. D **78**, 065016 (2008) [arXiv:0804.4672 [hep-th]].
- [3] S. Ferrara and P. van Nieuwenhuizen, Phys. Rev. Lett. **37**, 1669 (1976). D. Z. Freedman, A. K. Das, Nucl. Phys. **B120**, 221 (1977).

- [4] N. Boulanger and S. Leclercq, JHEP **0611**, 034 (2006) [hep-th/0609221].
- [5] N. Boulanger, S. Leclercq and P. Sundell, JHEP **0808**, 056 (2008) [arXiv:0805.2764 [hep-th]].
- [6] R. R. Metsaev, Nucl. Phys. B **759**, 147 (2006) [hep-th/0512342].
- [7] R. R. Metsaev, Nucl. Phys. B **859**, 13 (2012) [arXiv:0712.3526 [hep-th]].
- [8] A. K. H. Bengtsson, I. Bengtsson and N. Linden, Class. Quant. Grav. **4**, 1333 (1987).
- [9] R. Manvelyan, K. Mkrtchyan and W. Ruhl, Nucl. Phys. B **836**, 204 (2010) [arXiv:1003.2877 [hep-th]]; Phys. Lett. B **696**, 410 (2011) [arXiv:1009.1054 [hep-th]]. K. Mkrtchyan, arXiv:1101.5643 [hep-th].
- [10] A. Sagnotti and M. Taronna, Nucl. Phys. B **842**, 299 (2011) [arXiv:1006.5242 [hep-th]]; M. Taronna, arXiv:1005.3061 [hep-th].
- [11] E. S. Fradkin and M. A. Vasiliev, Nucl. Phys. B **291**, 141 (1987); Phys. Lett. B **189**, 89 (1987).
- [12] G. Barnich and M. Henneaux, Phys. Lett. B **311**, 123 (1993) [hep-th/9304057]. M. Henneaux, Contemp. Math. **219**, 93 (1998) [hep-th/9712226].
- [13] I. L. Buchbinder, A. Fotopoulos, A. C. Petkou and M. Tsulaia, Phys. Rev. D **74**, 105018 (2006) [hep-th/0609082]. A. Fotopoulos and M. Tsulaia, Phys. Rev. D **76**, 025014 (2007) [arXiv:0705.2939 [hep-th]]; JHEP **1011**, 086 (2010) [arXiv:1009.0727 [hep-th]]. A. Fotopoulos, N. Irges, A. C. Petkou and M. Tsulaia, JHEP **0710**, 021 (2007) [arXiv:0708.1399 [hep-th]]. P. Dempster and M. Tsulaia, arXiv:1203.5597 [hep-th].
- [14] R. R. Metsaev, arXiv:1205.3131 [hep-th].
- [15] S. Weinberg, Phys. Rev. **138**, B988 (1965). B. de Wit and D. Z. Freedman, Phys. Rev. D **21**, 358 (1980). M. Dubois-Violette and M. Henneaux, Lett. Math. Phys. **49**, 245 (1999) [math/9907135]; Commun. Math. Phys. **226**, 393 (2002) [math/0110088 [math-qa]].
- [16] C. Fronsdal, Phys. Rev. D **18**, 3624 (1978). J. Fang and C. Fronsdal, Phys. Rev. D **18**, 3630 (1978).
- [17] G. Barnich, F. Brandt and M. Henneaux, Commun. Math. Phys. **174**, 57 (1995) [hep-th/9405109]; Commun. Math. Phys. **174**, 93 (1995) [hep-th/9405194]; Phys. Lett. B **346**, 81 (1995) [hep-th/9411202]; Phys. Rept. **338**, 439 (2000) [hep-th/0002245].

- [18] N. Boulanger, T. Damour, L. Gualtieri and M. Henneaux, Nucl. Phys. B **597**, 127 (2001) [hep-th/0007220].
- [19] N. Boulanger and M. Esole, Class. Quant. Grav. **19**, 2107 (2002) [gr-qc/0110072].
- [20] G. Lucena Gómez, M. Henneaux and R. Rahman, Work in progress.
- [21] M. Porrati and R. Rahman, Nucl. Phys. B **801**, 174 (2008) [arXiv:0801.2581 [hep-th]]; arXiv:0809.2807 [hep-th]; Nucl. Phys. B **814**, 370 (2009) [arXiv:0812.4254 [hep-th]]; Phys. Rev. D **84**, 045013 (2011) [arXiv:1103.6027 [hep-th]]. M. Porrati, R. Rahman and A. Sagnotti, Nucl. Phys. B **846**, 250 (2011) [arXiv:1011.6411 [hep-th]]. R. Rahman, arXiv:1111.3366 [hep-th].
- [22] Y. .M. Zinoviev, Nucl. Phys. B **770**, 83 (2007) [hep-th/0609170]; Mod. Phys. Lett. A **24**, 17 (2009) [arXiv:0806.4030 [hep-th]]; Nucl. Phys. B **821**, 431 (2009) [arXiv:0901.3462 [hep-th]]. I. L. Buchbinder, T. V. Snegirev and Y. .M. Zinoviev, arXiv:1204.2341 [hep-th].
- [23] C. Lorce, Phys. Rev. D **79**, 113011 (2009) [arXiv:0901.4200 [hep-ph]].
- [24] M. Porrati, R. Rahman, Phys. Rev. **D80**, 025009 (2009). [arXiv:0906.1432 [hep-th]].
- [25] J. Scherk and J. H. Schwarz, Phys. Lett. B **82**, 60 (1979); Nucl. Phys. B **153**, 61 (1979). B. de Wit, P. G. Lauwers and A. Van Proeyen, Nucl. Phys. B **255**, 569 (1985). S. Deser and B. Zumino, Phys. Rev. Lett. **38**, 1433 (1977).
- [26] X. Bekaert, N. Boulanger and S. Leclercq, J. Phys. A **43**, 185401 (2010) [arXiv:1002.0289 [hep-th]].
- [27] M. Taronna, JHEP **1204**, 029 (2012) [arXiv:1107.5843 [hep-th]].
- [28] D. Francia and A. Sagnotti, Phys. Lett. B **543**, 303 (2002) [hep-th/0207002]; Class. Quant. Grav. **20**, S473 (2003) [hep-th/0212185].
- [29] X. Bekaert and N. Boulanger, Phys. Lett. B **561**, 183 (2003) [hep-th/0301243]; Commun. Math. Phys. **271**, 723 (2007) [hep-th/0606198].
- [30] R. Britto, F. Cachazo, B. Feng and E. Witten, Phys. Rev. Lett. **94**, 181602 (2005) [hep-th/0501052].
- [31] P. Benincasa and F. Cachazo, arXiv:0705.4305 [hep-th]. P. Benincasa and E. Conde, JHEP **1111**, 074 (2011) [arXiv:1106.0166 [hep-th]]; arXiv:1108.3078 [hep-th]. P. Benincasa, arXiv:1201.3191 [hep-th].

- [32] R. R. Metsaev, Phys. Lett. B **354**, 78 (1995); Phys. Lett. B **419**, 49 (1998) [hep-th/9802097]. A. Fotopoulos, K. L. Panigrahi and M. Tsulaia, Phys. Rev. D **74**, 085029 (2006) [hep-th/0607248].
- [33] K. Alkalaev, JHEP **1103**, 031 (2011) [arXiv:1011.6109 [hep-th]]. N. Boulanger and E. D. Skvortsov, JHEP **1109**, 063 (2011) [arXiv:1107.5028 [hep-th]]. N. Boulanger, E. D. Skvortsov and Y. M. Zinoviev, J. Phys. A **44**, 415403 (2011) [arXiv:1107.1872 [hep-th]]. M. A. Vasiliev, arXiv:1108.5921 [hep-th].
- [34] E. Joung and M. Taronna, Nucl. Phys. B **861**, 145 (2012) [arXiv:1110.5918 [hep-th]]. E. Joung, L. Lopez and M. Taronna, arXiv:1203.6578 [hep-th].
- [35] M. A. Vasiliev, Phys. Lett. B **243**, 378 (1990); Class. Quant. Grav. **8**, 1387 (1991); Class. Quant. Grav. **8**, 1387 (1991); Phys. Lett. B **567**, 139 (2003) [hep-th/0304049].
- [36] T. Damour and S. Deser, Annales Poincare Phys. Theor. **47**, 277 (1987).