

SKEW-PRODUCT SYSTEMS OVER INFINITE INTERVAL EXCHANGE TRANSFORMATIONS

HENK BRUIN AND OLGA LUKINA

ABSTRACT. We study the ergodic properties (recurrence, discrepancy, diffusion coefficients and ergodicity itself) of a class of \mathbb{Z} -extensions over infinite interval exchange transformations called rotated odometers. The choice of a skew-function is motivated by the use in the study of parallel flows on a particular staircase manifold of infinite genus.

1. INTRODUCTION

A rotated odometer $F_\pi : I \rightarrow I$ is a type of infinite interval exchange transformation (IIET) on the half-open unit interval $I = [0, 1)$, introduced in [7]. In this paper, we study \mathbb{Z} -extensions, that is, skew-products on $I \times \mathbb{Z}$ over these rotated odometers, and ask ourselves questions on their recurrence or transience, discrepancy, diffusion coefficients, and ergodicity.

1.1. Rotated odometers. The dyadic odometer is the group action on $\{0, 1\}^{\mathbb{N}}$ given by the addition of 1 to the left-most digit with infinite carry to the right. The dyadic odometer has a structure of a profinite group, and so admits a unique ergodic invariant measure (Haar measure). It is well known that, as a measure-preserving dynamical system, the dyadic odometer is isomorphic to the Lebesgue-measure preserving von Neumann-Kakutani map on the unit interval $I = [0, 1)$:

$$(1.1) \quad \mathbf{a}(x) = x - (1 - 3 \cdot 2^{-n}) \quad \text{if } x \in I_n = [1 - 2^{1-n}, 1 - 2^{-n}), \quad n \geq 1.$$

A *rotated odometer* is obtained as a composition of \mathbf{a} with a permutation of a partition of I into q intervals of equal length. Namely, take a positive integer q that is not a power of 2 (see [6] for the case when $q = 2^m$, $m \in \mathbb{N}$), and a permutation π on the symbols $\{0, \dots, q-1\}$. Divide the unit interval into q equal-sized half-open intervals $I_j = [j/q, (j+1)/q)$, $j = 0, \dots, q-1$, and define

$$R_\pi : I \rightarrow I, \quad I_j \mapsto I_{\pi(j)}.$$

For instance, a rotation by p/q , for $0 < p \leq q-1$, is an example of such a map, but not every R_π is induced by a rotation.

Definition 1.1. *A rotated odometer is an IIET given by the composition*

$$F_\pi = \mathbf{a} \circ R_\pi : I \rightarrow I,$$

where π is a permutation of q symbols.

Date: August 26, 2024.

2020 Mathematics Subject Classification: Primary: 37C83, 37E05, 28D05, Secondary: 37B10, 37E20, 37E35.

Keywords: infinite interval exchange transformation, substitution, skew-products, essential values, ergodicity, discrepancy, diffusion coefficient.

The dynamics of rotated odometers was studied in [7] using methods of symbolic dynamics, see Section 2 for details. In particular, associated to the aperiodic subsystem of a rotated odometer, there is an eventually periodic sequence of substitutions $(\chi_k)_{k \geq 1}$, and $q \times q$ -substitution matrices M_k , $k \geq 1$, which have non-negative integer entries. In this paper, we restrict to the case when the sequences of substitutions and the associated matrices are stationary, i.e., $\chi_k = \chi$ and $M_k = M$, $k \geq 1$, for some substitution χ and a $q \times q$ matrix M . We use the properties of χ and M in our analysis of the \mathbb{Z} -extensions of rotated odometers below.

Remark 1.2. As shown in [7, Theorem 1.1], every rotated odometer in Definition 1.1 can be realized as the first return map of a flow of rational slope on an infinite type translation surface L , where L has finite area, infinite genus, and at least one and at most a finite number of punctures. One of the punctures corresponds to a wild singularity in the metric completion of the surface, which is the accumulation point of the handles, and other (if any) punctures to cone angle singularities; see [24] for more on translation surfaces of infinite type.

1.2. Non-ergodicity of \mathbb{Z} -extensions of rotated odometers. Throughout this article, given a probability measure μ on (a subset of) $I = [0, 1]$ (most frequently, μ is Lebesgue) that is preserved by F_π , we denote by (I, F_π, μ) the corresponding measure-preserving dynamical system. We use the notation (I, F_π) when we only speak about the topological properties of the system.

To define the skew-product of a rotated odometer (I, F_π, Leb) , let $\psi : I \rightarrow \mathbb{Z}$ be a *skew-function* given by

$$(1.2) \quad \psi(x) = \begin{cases} 1, & 0 \leq x < \frac{1}{2}, \\ -1, & \frac{1}{2} \leq x < 1. \end{cases}$$

The skew-product of F_π and ψ is given by

$$(1.3) \quad T_\pi : I \times \mathbb{Z} \rightarrow I \times \mathbb{Z}, \quad (x, n) \mapsto (F_\pi(x), n + \psi(x)),$$

with invariant measure $\text{Leb} \otimes \nu$, where ν is the counting measure on \mathbb{Z} . The skew-product (1.3) is also called the \mathbb{Z} -*extension* of F_π with skew-function ψ .

Remark 1.3. The choice of the skew-function ψ in (1.2) is motivated by the interpretation of rotated odometers as the first return maps of flows of rational slope on a translation surface, see Remark 1.2. From this point of view, the skew-product (1.3) is interpreted as the first return map of the lifted flow on a \mathbb{Z} -fold covering space of the translation surface in Remark 1.2, to the section $I \times \mathbb{Z}$. The choice of the points of discontinuity for the step function ψ is motivated by the geometry of the staircase.

Problem 1.4. *Is the skew-product dynamical system (1.3) recurrent or transient?*

A necessary (but not sufficient) condition for the skew-product to be ergodic is that it is recurrent to a subsection $I \times \{0\}$, i.e., Leb-a.e. $x \in I \times \{0\}$ returns to $I \times \{0\}$ infinitely often. Since the invariant measure $\text{Leb} \otimes \nu$ is infinite, this recurrence doesn't follow directly from Poincaré recurrence, but by a result of Atkinson [2], T_π is recurrent if and only if $\int_I \psi d\text{Leb} = 0$.

If π is the identity permutation, then $F_\pi = \mathbf{a}$, and we denote the skew-product map by $T_{\mathbf{a}}$. In this case $(I \times \mathbb{Z}, T_{\mathbf{a}})$ is the first return map to $I \times \mathbb{Z}$ of the vertical flow on the staircase.

This flow itself, i.e., the \mathbb{R} -extension of the von Neumann-Kakutani map \mathbf{a} , has been studied for various classes of skew-functions, see for instance [16, 23] and references therein. In particular, [16] studies \mathbb{R} -extensions when the skew-product function is a step function with two discontinuities. For ψ as in (1.2), [16, Theorem 2] implies that the first return map of the vertical flow is not ergodic.

Problem 1.5. *Show that, for any $q \neq 2^n$, $n \in \mathbb{N}_0$, and any permutation π of q intervals, the skew-product $(I \times \mathbb{Z}, T_\pi, \text{Leb} \otimes \nu)$ is not ergodic.*

Since the family of all rotated odometers is rather large, and they exhibit diverse properties, we cannot give an affirmative answer to Problem 1.5 for a \mathbb{Z} -extension of any rotated odometer. Rather, we give a few criteria which allow us to rule out ergodicity, and present examples which satisfy these criteria. So far we have not found an example of an ergodic \mathbb{Z} -extension in this context.

Our approach to this problem is as follows. In Section 2.2 we recall from [7] that there is a sequence of substitutions $(\chi_k)_{k \geq 1}$, associated to a rotated odometer (I, F_π, Leb) , which assigns to each $i \in \mathcal{A} = \{0, \dots, q-1\}$ a finite word $\chi(i)$ in the alphabet \mathcal{A} . Since there are only finitely many permutations on q letters, $(\chi_k)_{k \geq 1}$ is (pre-)periodic. If this sequence is constant, then the rotated odometer (I, F_π) is called *stationary*, and we write $\chi_k = \chi_1 = \chi$.

Assumption 1.6. *Let (I, F_π) be a rotated odometer. In what follows, we assume that:*

- (1) *The rotated odometer (I, F_π) is stationary,*
- (2) *Lebesgue measure Leb is ergodic for (I, F_π) . Note that we do not assume that Leb is a unique invariant measure, and in fact most our examples are not uniquely ergodic.*

Using the coding procedure described in Proposition 2.3, for each $x \in I$ we can describe a finite piece of its orbit of arbitrary length as a subword of a substitution word $\chi^k(a)$, for some $a \in \mathcal{A}$, and $k \in \mathbb{N}$ large enough. Using the skew-function ψ from (1.2), we can associate to $\chi(a)$ its *weight* $\psi(\chi(a))$ which is an integer, see Section 2.4 for details. Similarly we can associate weights $\psi(\ell_j)$ to left eigenvectors ℓ_j , $0 \leq j \leq q-1$, of the matrix M describing the substitution. These weights play a central role in our proofs.

Remark 1.7. The standard approach to study the dynamics of systems given by sequences of substitutions is by associating to them an *S-adic subshift*, see Section 2.1 for details. In an *S-adic shift*, the phase space is obtained as the closure of a sequence $\rho = \lim_k \chi_1 \circ \dots \circ \chi_k(a_k)$ in $\mathcal{A}^{\mathbb{N}_0}$ with cylinder set topology, and the dynamics is given by a shift action. For rotated odometers, ρ corresponds to the orbit of 0 which is transitive in the minimal subset of I , but not transitive in I , unless (I, F_π, Leb) is uniquely ergodic, which is not always the case. That is the reason we do not use the formalism of *S-adic shifts* in our results.

We now state two criteria for non-ergodicity of rotated odometers. The first one is formulated in terms of the weights of the substitution words.

Theorem 1.8. *Let (I, F_π, Leb) be a stationary rotated odometer. Let $T_\pi : I \times \mathbb{Z} \rightarrow I \times \mathbb{Z}$ be its \mathbb{Z} -extension with skew-function ψ as in (1.2), and $\mathbf{d} := \gcd\{\psi(\chi(a)) : a \in \mathcal{A}\}$. If $\mathbf{d} \geq 2$, then the skew-product is not ergodic.*

Example 5.7 of a rotated odometer with $q = 11$ satisfies the hypothesis of Theorem 1.8 with $\mathbf{d} = 2$, and so it is not ergodic. Example 5.3 with $q = 9$ also satisfies Theorem 1.8 with

$\mathbf{d} = 2$. For the latter, we can give a more precise description of the ergodic properties using Theorem 1.9 below.

Theorem 1.9. *Let (I, F_π, Leb) be a stationary rotated odometer. Let $T_\pi : I \times \mathbb{Z} \rightarrow I \times \mathbb{Z}$ be its \mathbb{Z} -extension with skew-function ψ as in (1.2). Assume that for all eigenvalues λ_j of the associated matrix M of χ with norm $|\lambda_j| \geq 1$, the left eigenvectors have weights $\psi(\ell_j) = 0$, with the exception of one, say, for $\lambda_c > 1$, which is Pisot. Suppose the algebraic and the geometric multiplicities of λ_c are equal. Then Lebesgue measure Leb of the \mathbb{Z} -extension T_π has infinitely many ergodic components.*

Theorem 1.9 applies to uniquely ergodic rotated odometers in Examples 5.1 and 5.2 for $q = 3$, Example 5.5 for $q = 5$. In these examples, there are two eigenvalues of algebraic multiplicity 1 outside of the unit circle, and for the largest one $\psi(\ell_0) = 0$ (this is always true for rotated odometer with ergodic Lebesgue measure). Theorem 1.9 also applies to Example 5.3 for $q = 9$, where (I, F_π, Leb) is uniquely ergodic, and the second largest eigenvalue λ_1 is Pisot and has geometric and algebraic multiplicity 2. Example 5.5 for $q = 5$ is not uniquely ergodic, and satisfies the assumptions of the theorem. We refer the reader to Section 5 for further examples, to which Theorems 1.8 and 1.9 are not applicable.

1.3. Diffusion coefficient. The spread of particles over time, for example in a Brownian motion, is called diffusion, and if the (average or maximal) displacement of particles happens according to some power law, the exponent is called the *diffusion coefficient*. For Brownian motion, the average displacement is the square-root of time, so the diffusion coefficient is $\gamma = \frac{1}{2}$, and this is “the standard”. Non-standard diffusion is referred to as *anomalous diffusion*, and more precisely, it is called *super-diffusion* if the diffusion coefficient $\gamma > \frac{1}{2}$ and it is called *sub-diffusion* if $\gamma < \frac{1}{2}$. It is known that the Lorentz gas with circular obstacles and finite horizon (i.e, a particle will always hit an obstacle in bounded time) is diffusive [8], while the Lorentz gas with infinite horizon is super-diffusive [26], but only by a thin margin, because the diffusion coefficient is $\frac{1}{2}$ also here, but there is a logarithmic correction term that make the diffusion faster than for Brownian motion. Ehrenfest’s wind-tree model is similar to the Lorentz gas model, except that the obstacles are now aligned rectangles (or more general polygonal shapes, see [11]) situated at the lattice points of \mathbb{Z}^2 . The diffusion depends now on the direction of the flow. In the model with square obstacles, the diffusion coefficient for the typical direction is $\gamma = \frac{2}{3}$ [10], and this is found as the quotient of the largest two eigenvalues of a particular matrix. For more complicated polygonal shapes, the diffusion coefficients in the typical direction can be very different. For instance, [11] considers the a family of models with a certain geometric shape of an obstacle, where the diffusion coefficient depends only on the number of corners of the obstacle, and tends to zero as the number of corners increases. On the other hand, [11] also describes a model where the diffusion coefficient is very close to 1.

For \mathbb{Z} -extensions, the n -steps displacement is $\xi(T_\pi^n(x, k)) - \xi(x, k)$ for the projection $\xi(x, k) = k$. Clearly, the displacement is independent of $k \in \mathbb{Z}$, and thus the diffusion coefficient, as function of x , can be defined as

$$(1.4) \quad \gamma(x) = \limsup_{n \rightarrow \infty} \frac{\log(\xi(T_\pi^n(x, k)) - k)}{\log n}.$$

Unless F_π is uniquely ergodic, the dependence of $\gamma(x)$ on the point x cannot be avoided, so at best we can speak of the a.s. diffusion coefficient with respect to an ergodic measure (which will be Lebesgue, if χ is covering). Our main result about $\gamma(x)$ is the following.

Theorem 1.10. *Let (I, F_π, Leb) be a stationary rotated odometer. Suppose the associated matrix is diagonalizable, the largest eigenvalue λ_0 has multiplicity 1, and the second largest eigenvalue satisfies $|\lambda_1| > 1$, then for the corresponding \mathbb{Z} -extension, the diffusion coefficients satisfy*

$$\gamma(x) \leq \gamma_0 := \frac{\log |\lambda_1|}{\log \lambda_0} \text{ for Leb-a.e. } x.$$

1.4. Discrepancy. A sequence of points $(x_i)_{i \geq 1}$ in I is called uniformly distributed if, in the limit, the number of times the sequence hits a subinterval J of I is proportional to the length of J . It is well-known that, for a measure-preserving system with an ergodic invariant probability measure, the orbit of almost every point is uniformly distributed. Discrepancy is a quantitative characteristic of a sequence $(x_i)_{i \geq 1}$, which can be understood as characterizing the properties of the uniform distribution, i.e., how far it is from the ideal one. A classical reference for the results on discrepancy is the book [20]. In this paper, rigorous definitions of the concepts mentioned above are given in Section 3.1.

For substitutions, the discrepancy of sequences arising as fixed points of primitive substitutions and the corresponding shifts were studied by Adamczewski [1]. His results [1] were extended to S -adic transformations in [4]. Since our rotated odometers are often described by substitution matrices which are not primitive, we include the following estimate for the discrepancy of orbits of rotated odometers.

Theorem 1.11. *Suppose that F_π is covering and stationary, the associated matrix is diagonalizable, and the largest eigenvalue λ_0 has multiplicity 1. Then for Lebesgue-a.e. x , there is $C = C_x$ such that the F_π -orbit of x has discrepancy $\mathfrak{D}_R \leq C_x \cdot R^{\gamma_0 - 1}$ for $\gamma_0 := \max \left\{ \frac{\log |\lambda_1|}{\log \lambda_0}, 0 \right\}$.*

The rest of the paper is organized as follows. Section 2 recalls preliminaries, mostly from [7] but also about essential values from [25]. Section 3.1 computes the discrepancy of the rotated odometers, as well as the diffusion coefficients of the skew-products. Recurrence and (non-)ergodicity of the skew-products are covered in Section 4.2. In Section 5, we list the examples which illustrate our theorems.

2. PRELIMINARIES OF ROTATED ODOMETERS AND ESSENTIAL VALUES

We recall some results of [6, 7] on the properties of rotated odometers, and techniques to study them. Namely, [7, Theorem 1.2] proves that a rotated odometer (I, F_π) as in Definition 1.1 has the following properties:

- (1) There exists a decomposition $I = I_{per} \cup I_{np}$, with I_{per} possibly empty, such that every point in I_{per} is periodic under F_π , $0 \in I_{np}$, and the restriction $F_\pi : I_{np} \rightarrow I_{np}$ is invertible at every point except 0, and has no periodic points.
- (2) There is a unique minimal subsystem (I_{min}, F_π) of (I_{np}, F_π) .
- (3) If I_{per} is non-empty, then it is a finite or a countable union of half-open subintervals of I .

- (4) The minimal subsystem (I_{min}, F_π) is uniquely ergodic.
- (5) The aperiodic subsystem (I_{np}, F_π) is ergodic, with at most q invariant ergodic probability measures.

2.1. Substitutions. To study the dynamics of (I, F_π) further, we code the orbits of points in (I, F_π) by words in the alphabet $\mathcal{A} = \{0, \dots, q-1\}$. Let \mathcal{A}^* be the set of all non-empty words of finite length in this alphabet, and let $\Sigma = \mathcal{A}^{\mathbb{N}_0}$ be the set of one-sided infinite sequences in this alphabet.

Definition 2.1. A substitution $\chi : \mathcal{A} \rightarrow \mathcal{A}^*$ is a map that assigns to every $a \in \mathcal{A}$ a single word $\chi(a) \in \mathcal{A}^*$. It extends to \mathcal{A}^* and Σ by concatenation:

$$\chi(a_1 a_2 \dots a_r) = \chi(a_1) \chi(a_2) \dots \chi(a_r), \quad r \geq 1.$$

The $q \times q$ matrix M , where the (i, j) -th entry is the number of letters j in $\chi(i)$, is called the associated matrix of χ .

Definition 2.2. A substitution $\chi : \mathcal{A} \rightarrow \mathcal{A}^*$ is:

- primitive, if there is $r \geq 1$ such that for all $i \in \mathcal{A}$, $\chi^r(i)$ contains every letter in \mathcal{A} .
- proper if there exist two letters $a, b \in \{0, \dots, q-1\}$ such that for all $i \in \{0, \dots, q-1\}$, the first letter of $\chi(i)$ is a and the last letter of $\chi(i)$ is b .

The substitutions we consider are always *proper* and sometimes also *primitive*. Recall that a square matrix M is *primitive* if it has a power with strictly positive entries. Thus a substitution χ is primitive as in Definition 2.2 if and only if its associated matrix M is primitive.

If $\chi(a)$ starts with a , we get a fixed point of χ which (unless $\chi(a) = a$) is an infinite sequence

$$(2.1) \quad \rho = \rho_0 \rho_1 \rho_2 \rho_3 \dots = \lim_{j \rightarrow \infty} \chi^j(a) \in \Sigma.$$

For sequences $s = (s_k) \in \Sigma$, define the *left shift* by

$$(2.2) \quad \sigma : \Sigma \rightarrow \Sigma, \quad s_0 s_1 s_2 \dots \mapsto s_1 s_2 s_3 \dots$$

The dynamical system (X_ρ, σ) where $X_\rho = \overline{\{\sigma^n(\rho) : n \geq 0\}}$ is called a *substitution subshift*.

2.2. Renormalization and symbolic dynamics. We now briefly describe the renormalization procedure developed in [7], which allows us to associate to the rotated odometer (I, F_π) a substitution dynamical system. For that we restrict to the aperiodic subsystem (I_{np}, F_π) of the rotated odometer (I, F_π) .

Let $q \geq 2$ and let $F_\pi = \mathbf{a} \circ R_\pi$ be a rotated odometer as in Definition 1.1. Let

$$(2.3) \quad N = \min\{n \geq 1 : 2^N \geq q\} \quad \text{and} \quad L_k = [0, 2^{-kN}).$$

Denote by $\mathcal{P}_{0,q}$ the partition of $I = [0, 1)$ into q equal length intervals, which are the intervals of continuity for R_π in the definition of F_π . The half-open interval L_k is called the *k-th section*.

For each $q \geq 1$ and each $k \geq 1$, we define a partition $\mathcal{P}_{kN,q}$ of the unit interval into $q2^N$ intervals of equal length $q^{-1}2^{-N}$. This partition refines $\mathcal{P}_{mN,q}$, $0 \leq m \leq k-1$, and induces the partition $\mathcal{P}_{kN,q}^{cod}$ of the k -section L_k into q subintervals $L_{k,i}$, $0 \leq i \leq q-1$. We recall the following result.

Proposition 2.3. [7, Proposition 5.11, Lemma 5.15] *Let $F_{\pi,k} : L_k \rightarrow L_k$, $k \geq 1$ be the first return maps of the rotated odometer $F_\pi : I \rightarrow I$.*

Then for $k \geq 1$ there exist permutations π_k of q symbols, and finite IET $R_{\pi,k} : L_k \rightarrow L_k$ of the partitions $\mathcal{P}_{kN,q}^{\text{cod}}$, defined by π_k , such that

- (1) $F_{\pi,k} = \mathbf{a}_k \circ R_{\pi,k}$, where \mathbf{a}_k is a scaled copy of the von Neumann-Kakutani map \mathbf{a} , given by

$$\mathbf{a}_k(x) = \frac{1}{2^{kN}} \mathbf{a}(2^{kN}x).$$

- (2) *The sequence $(\pi_k)_{k \geq 1}$ is pre-periodic.*
 (3) *If $(\pi_k)_{k \geq 1}$ is strictly pre-periodic, and $S = \{\pi_1, \dots, \pi_{k_0-1}\}$ is the pre-periodic part of the sequence, then none of the permutations in S occurs in the periodic part.*
 (4) *For each set $L_{k,i}$, $0 \leq i \leq q-1$ of the coding partition $\mathcal{P}_{kN,q}$, the order in which $L_{k,i}$ visits the sets of $\mathcal{P}_{(k-1)N,q}$ under the first return map $F_{\pi,k} : L_{k-1} \rightarrow L_{k-1}$ defines a substitution word $\chi_k(i)$ with the alphabet $\mathcal{A} = \{0, \dots, q-1\}$. Thus associated to each section L_k , $k \geq 1$, there is a substitution $\chi_k : \mathcal{A} \rightarrow \mathcal{A}^*$.*
 (5) *Every substitution in the sequence $(\chi_k)_{k \geq 1}$ is proper, namely, every word $\chi_k(i)$ starts with 0, and ends with b_k , depending only on k and not on $i \in \{0, \dots, q-1\}$.*

Thus for each $i \in \mathcal{P}_{kN,q}$ the word $\chi_k(i)$ is the itinerary of the half-open interval $L_{k,i}$ in L_{k-1} , with respect to the partition $\mathcal{P}_{(k-1)N,q}$ of L_{k-1} , and, inductively, $\chi_1 \circ \chi_2 \circ \dots \circ \chi_k(i)$ is the itinerary of $L_{k,i}$ in $I = [0, 1)$ with respect to the partition $\mathcal{P}_{0,q}$.

We now can proceed similarly to the case of a single substitution in Section 2.1. By Proposition 2.3, item (5), the sequence

$$\rho = \lim_{k \rightarrow \infty} \chi_1 \circ \dots \circ \chi_k(0)$$

is well-defined and it is an S -adic fixed point of $(\chi_k)_{k \geq 1}$. The sequence ρ is also the itinerary of 0 in I with respect to the partition $\mathcal{P}_{0,q}$.

We define the S -adic subshift (X_ρ, σ) similarly to Section 2.1, formula (2.2), and the paragraph below (2.2). Recall that the rotated odometer (I, F_π) has a unique minimal subsystem (I_{\min}, F_π) , and the orbit of 0 under F_π lies dense in I_{\min} . Thus (X_ρ, σ) models the dynamics on the minimal set (I_{\min}, F_π) . More precisely, there is a homeomorphism onto its image $h : I_{\min} \rightarrow X_\rho$, such that $X_\rho \setminus h(X_\rho)$ is countable, and $h \circ F_\pi(x) = \sigma \circ h(x)$.

In many examples of rotated odometers I_{\min} is a proper subset of the aperiodic subsystem I_{np} , so we cannot restrict to considering only (X_ρ, σ) . Proposition 2.3 allows us to code the orbits outside the minimal set as well. Before we state the result which we will use, we have to introduce *covering* substitutions.

Definition 2.4. *Let $F_\pi : I \rightarrow I$ be a rotated odometer as in Definition 1.1, and let $F_{\pi,k} : L_k \rightarrow L_k$ be the sequence of first return maps as in Proposition 2.3. Then $F_{\pi,k}$ is covering if the orbits of the sets in $\mathcal{P}_{kN,q}^{\text{cod}}$ visit every element of $\mathcal{P}_{kN,q}$ in L_{k-1} , that is, if*

$$(2.4) \quad \bigcup_{i=0}^{q-1} \bigcup_{t=0}^{t_{k,i}-1} F_{\pi,k-1}^t(I_{k,i}) = L_{k-1}.$$

Alternatively, $F_{\pi,k}$ (or the associated substitution χ_k) is covering if $\sum_{i \in \mathcal{A}} |\chi_k(i)| = q2^N$.

The importance of *covering* substitutions is underlined by the following result from [7]:

Proposition 2.5. *Lebesgue measure is ergodic (and hence F_π is aperiodic on I) if and only if all the substitutions χ_k (or just χ if it is stationary) are covering.*

Thus, in our theorems, when we consider a rotated odometer with Lebesgue ergodic measure, we are automatically considering covering substitutions.

Let (X_χ, σ) denote the smallest subshift such that every subword W that it contains is a subword of $\chi_s \circ \cdots \circ \chi_1(a)$ for some $a \in \mathcal{A}$ and $s \in \mathbb{N}$; it symbolically describes the aperiodic part (I_{np}, F_π) of the rotated odometer.

Lemma 2.6 is a special case of [7, Theorem 5.10], when every substitution in $(\chi_k)_{k \geq 1}$ is covering. Since our main results are for ergodic Lebesgue measures, we include a simple proof of the existence of coding in this case for completeness. The proof of the general case in [7, Theorem 5.10] requires more involved techniques.

Lemma 2.6. *Let (I, F_π) be a rotated odometer, with associated sequence of substitutions $(\chi_k)_{k \geq 1}$. Suppose χ_k is covering for all $k \geq 1$. Then for any $x \in I$ and $n \geq 1$, there exists $s \in \mathbb{N}$, $a \in \mathcal{A}$ and a subword W of $\chi_s \circ \cdots \circ \chi_1(a)$ such that W is the itinerary of the orbit $\{F_\pi^k(x) : 0 \leq k \leq n\}$ with respect to the coding partition $\mathcal{P}_{0,q}$.*

Proof. Let $(L_k)_{k \geq 1}$ be the sequence of sections in Proposition 2.3, and let $(\chi_k)_{k \geq 1}$ be the sequence of associated substitutions. Let $x \in I$, and fix an $n \in \mathbb{N}_0$. For convenience, we introduce the notation

$$\chi^{(k)}(a) = \chi_k \circ \cdots \circ \chi_1(a), \quad a \in \mathcal{A}.$$

Since every χ_k is covering for every $k \geq 1$, there is $a_1 \in \mathcal{A}$ such that $x \in F_\pi^{j_0}(L_{k,a_1})$ for some $j_0 \geq 0$. Without loss of generality we may assume that k is so large that $x \notin L_k$, and thus $j_0 > 0$.

If $|\chi^{(k)}(a_1)| - j_0 > n$, then the part of the substitution word $\chi^{(k)}(a_1)$, starting from the j_0 -th symbol and of length n , codes the orbit of x of length n .

Otherwise we proceed as follows. The orbit of L_{k,a_1} returns to L_k after $|\chi^{(k)}(a_1)|$ iterations. Then the orbit of x returns to L_k after $j_1 = |\chi^{(k)}(a_1)| - j_0$ iterations. Since F_π is not invertible at 0, $F^{j_1}(x) \neq 0$, and thus there exists a half-open subinterval $V_1 \subset L_{k,a_1}$ with $x \in F_\pi^{j_0}(V_1)$ and such that $F^{j_0+j_1}(V_1) \subset L_{k,a_2}$ for some $a_2 \in \mathcal{A}$. If $|\chi^{(k)}(a_1)| - j_0 + |\chi^{(k)}(a_2)| > n$, then the part of the concatenated substitution word $\chi^{(k)}(a_1)\chi^{(k)}(a_2)$, starting from the j_0 -th symbol and of length n , codes the orbit of x of length n .

If not, then, continuing by induction, we can find a sequence of half-open subintervals $V_s \subset V_{s-1} \subset \cdots \subset V_1 \subset L_{k,a_1}$, and a sequence of letters a_1, a_2, \dots, a_s such that, for any $1 \leq i < s$, $x \in F^{j_0}(V_i)$, and $F^{j_0+\sum_{m=1}^i j_m}(V_i) \subset L_{k,a_{i+1}}$, and

$$|\chi^{(k)}(a_1)| - j_0 + \sum_{i=2}^s |\chi^{(k)}(a_i)| > n.$$

The previous inequality holds for some $s \geq 1$ since all substitution words have non-zero length. Thus the part of the concatenated substitution word $\chi^{(k)}(a_1) \cdots \chi^{(k)}(a_s)$ of length n starting from the j_0 -th symbol codes the orbit of x of length n . The choice of k in

this argument is not unique; taking larger values of k will result in a smaller value of s . Nevertheless, the coding is unique for a given $x \in I$, since it is always done with respect to the same partition $\mathcal{P}_{0,q}$. In our argument, we do not need to assume that x is a Lebesgue typical point; rather, any open neighborhood of x contains an interval and so has positive measure; therefore it contains a Lebesgue typical point, whose orbit of length n has the same coding as the orbit of x . \square

In Proposition 2.3, the sequence $(\chi_k)_{k \geq 1}$ is pre-periodic. Therefore, the properties of the corresponding shift are effectively the same as for the case of a single substitution. Namely, for k_0 the length of the pre-periodic part of $(\chi_k)_{k \geq 1}$ and p_0 the period, set

$$(2.5) \quad M = M_{k_0+p_0} \cdots M_{k_0+1}.$$

Thus we may reduce the problem to the study of a single substitution χ with associated matrix M . If $\chi_k = \chi$ for all $k \geq 1$, we say that $\chi_1 = \chi$ is stationary.

2.3. The associated matrix in Frobenius form. We now assume that the sequence of substitutions associated to the rotated odometer (I, F_π) is stationary. The matrix M of a covering substitution has column sums equal to 2^N , so $\lambda_0 = 2^N$ is also the leading eigenvalue, with a positive left eigenvector $\ell_0 = (1, 1, \dots, 1)$. This reflects the constant density of Lebesgue measure. However, even if Lebesgue measure is ergodic, it needs not be the only ergodic invariant measure. This can be seen from the *Frobenius form* of the matrix M : by conjugating with a permutation matrix Π , corresponding to relabeling the symbols in \mathcal{A} , we can obtain the form

$$(2.6) \quad M_F = \Pi^{-1} M \Pi = \begin{pmatrix} D_1 & O & \cdots & \cdots & O \\ A_{2,1} & D_2 & O & \cdots & O \\ A_{3,1} & A_{3,2} & D_3 & \cdots & \vdots \\ \vdots & \cdots & \cdots & \ddots & \vdots \\ A_{d,1} & \cdots & \cdots & A_{d,d-1} & D_d \end{pmatrix},$$

where the D_i are square diagonal blocks, which are either irreducible or zero matrices, O are (rectangular) blocks of zeroes and the $A_{i,j}$ are rectangular non-negative matrices. The block D_1 represents the (unique) minimal subsystem of the rotated odometer. Since $\chi(a)$ starts with 0 for each $a \in \mathcal{A}$, the first column of M is strictly positive, and so all $A_{i,1}$, $2 \leq i \leq d$, are non-zero.

Clearly

$$(2.7) \quad \det(M_F - \lambda I) = \prod_{k=1}^d \det(D_k - \lambda I_k),$$

where I and I_k stand for the identity matrices of the correct size. Hence every eigenvalue of M must be an eigenvalue of D_k for some k , and vice versa. In particular, as a consequence of the Perron-Frobenius Theorem, the number of eigenvalues λ_j of M that are strictly greater than 1 is at least the number of non-zero diagonal blocks in (2.6). It follows from [5] that the number of ergodic invariant measures for the rotated odometer (I, F_π) corresponds to the number of eigenvalues greater than 1 with non-negative eigenvectors. Such eigenvalues are called *distinguished* and every non-zero block D_k has a distinguished eigenvalue $\lambda_k > 1$ provided $A_{k,j}$ is zero for each $\lambda_j \geq \lambda_k$.

If a substitution is not covering, then M can still be put into the Frobenius form, and the same method of determining the number of ergodic invariant measures applies. The difference with the covering case is that the column sums no longer are equal to 2^N .

2.4. Doubling the middle symbol. We now describe a modification of the stationary substitution $\chi : \mathcal{A} \rightarrow \mathcal{A}^*$ associated to a rotated odometer $F_\pi : I \rightarrow I$, which we need to study its skew-product extensions.

Recall from (1.2) that the skew-function ψ has two intervals of continuity, $[0, \frac{1}{2})$ and $[\frac{1}{2}, 1)$. If the number q in the definition of F_π , and, consequently, the number of sets in the partition $\mathcal{P}_{0,q}$ of the unit interval used to define an IET R_π is odd, then the skew-function ψ is not constant on the middle interval $[\frac{m}{q}, \frac{m+1}{q})$.

For this reason, we split this middle interval into two halves, with symbols m^+ for $[\frac{m}{q}, \frac{1}{2})$ and m^- for $[\frac{1}{2}, \frac{m+1}{q})$, so that the new alphabet $\mathcal{A}' = \{0, 1, \dots, m^+, m^-, \dots, q-1\}$ has $q' = q + 1$ letters.

The corresponding $q' \times q'$ -matrix M' has identical m^+ -th and m^- -th rows, but the m^+ -th and m^- -th column need not be the same. Consequently, compared to the eigenvalues of M , M' will have one extra eigenvalue 0. If χ is covering, the leading eigenvalue is still 2^N , but the corresponding left eigenvector is $(1, \dots, 1, \frac{1}{2}, \frac{1}{2}, 1, \dots, 1)$ and the column sum is 2^N only if you leave out the contribution of the m^- -th row, and also add the m^+ -th and m^- -th columns together.

Definition 2.7. Let $\mathcal{A}' = \{0, 1, \dots, m^+, m^-, \dots, q-1\}$ be an extended alphabet, and $\chi : \mathcal{A}' \rightarrow (\mathcal{A}')^*$ be a substitution. Let $\psi : I \rightarrow \mathbb{Z}$ be the skew-function defined in (1.2). Then the weight of a letter $\chi(a)$, $a \in \mathcal{A}'$, is given by

$$\psi(a) = \begin{cases} +1, & a \in \{0, \dots, m-1, m^+\}, \\ -1, & a \in \{m^-, m+1, \dots, q-1\}. \end{cases}$$

The weight of a substitution word $\chi(a) = w_1 \cdots w_\ell$, for $\ell \in \mathbb{N}$, is given by

$$\psi(\chi(a)) = \sum_{i=1}^{\ell} \psi(w_i).$$

Alternatively, note that the number of occurrences of each letter in the substitution word $\chi(a)$, $a \in \mathcal{A}'$, is given by the a -th row m_a of the associated matrix M' . Then

$$\psi(\chi(a)) = m_a \left(\underbrace{1, \dots, 1}_{\text{from } 0 \text{ to } m^-}, \underbrace{-1, \dots, -1}_{\text{from } m^+ \text{ to } q-1} \right)^T.$$

In the rest of the paper, when we consider the skew-product system (1.3), we will often drop the prime accent and denote \mathcal{A}' and M' by \mathcal{A} and M , keeping in mind that they are then an alphabet with $q + 1$ letters, and a $(q + 1) \times (q + 1)$ -matrix respectively.

We will always order the eigenvalues by absolute value: $\lambda_0 \geq |\lambda_1| \geq \dots \geq |\lambda_q|$. The corresponding left eigenvectors are denoted by ℓ_j , and scaled in such a way that for some Jordan decomposition $M = UJU^{-1}$, the ℓ_j 's are the rows of U^{-1} , and also ℓ_0 is a non-negative vector.

2.5. Essential values. We start by recalling some basics of theory of essential values. Main references for this material are [21, 22, 25].

We denote by (X, \mathcal{B}, μ) the probability space, where \mathcal{B} is the collection of Borel sets in X , and μ is the probability measure.

Definition 2.8. Let (X, \mathcal{B}, μ) be the probability space, and $F : X \rightarrow X$ an ergodic transformation. A Borel map $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ is called a cocycle for F if

$$(2.8) \quad \Psi(x, n_1 + n_2) = \Psi(F^{n_2}(x), n_1) + \Psi(x, n_2).$$

for every $n_1, n_2 \in \mathbb{Z}$ and every $x \in X$, and $\mu(\bigcup_{n \in \mathbb{Z}} (\{x : F^n(x) = x\} \cap \{\Psi(x, n) \neq 0\})) = 0$.

A cocycle $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ is called a *coboundary* if there exists a Borel map $\phi : X \rightarrow \mathbb{Z}$ such that

$$(2.9) \quad \Psi(x, n) = \phi(F^n(x)) - \phi(x) \quad \text{for all } n \in \mathbb{Z}.$$

We now define essential values.

Definition 2.9. Let $F : X \rightarrow X$ be an ergodic transformation on the probability space (X, \mathcal{B}, μ) . Let $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ be a cocycle for F . Then an element $e \in \mathbb{Z}$ is called an essential value of Ψ if, for every $A \in \mathcal{B}$ with $\mu(A) > 0$,

$$\mu\left(\bigcup_{n \in \mathbb{Z}} A \cap F^{-n}(A) \cap \{x : \Psi(x, n) = e\}\right) > 0.$$

As described in [21], given any skew-function $\psi : X \rightarrow \mathbb{Z}$, one can define a cocycle $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ by

$$(2.10) \quad \Psi(x, n) = \begin{cases} \psi(F^{n-1}x) + \dots + \psi(F(x)) + \psi(x), & n \geq 1 \\ 0, & n = 0, \\ -\psi(F^n(x)) - \dots - \psi(F^{-1}(x)), & n \leq -1. \end{cases}$$

Conversely, given a cocycle Ψ , one can define a skew-function $\psi : X \rightarrow \mathbb{Z}$ by $\psi(x) = \Psi(x, 1)$.

Denote the partial ergodic sums by $S_n \psi(x) = \sum_{i=0}^{n-1} \psi(F^i(x))$. The following is well-known.

Lemma 2.10. Let $F : X \rightarrow X$ be an ergodic transformation on the probability space (X, \mathcal{B}, μ) . Let $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ be a cocycle for F , and let $\psi : X \rightarrow \mathbb{Z}$ be the skew-function defined by $\psi(x) = \Psi(x, 1)$. Then the element $e \in \mathbb{Z}$ is an essential value of the cocycle Ψ if and only if for every positive measure $A \in \mathcal{B}$ there exists an $n \in \mathbb{Z}$ such that

$$(2.11) \quad \mu(A \cap F^{-n}(A) \cap \{x \in X : S_n \psi(x) = e\}) > 0.$$

In particular, $e = 0$ is always an essential value, by taking $n = 0$ in (2.11).

Definition 2.11. Let $F : X \rightarrow X$ be an ergodic transformation on the probability space (X, \mathcal{B}, μ) , and let $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ be a cocycle for F . We call the essential value 0 non-trivial, if (2.11) holds for $e = 0$ and some $n_A \neq 0$, for any $A \in \mathcal{B}$ with $\mu(A) > 0$.

If 0 is a non-trivial essential value, then the cocycle $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ is recurrent.

If Ψ is not recurrent, then Ψ is transient.

Let $\overline{\mathbb{Z}} = \mathbb{Z} \cup \{\infty\}$ be the one-point compactification of \mathbb{Z} .

Definition 2.12. We say that ∞ is an essential value for the cocycle $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ (with associated skew-function ψ) if for every $N \in \mathbb{N}$, and every positive measure set $A \in \mathcal{B}$ there exists an $n \in \mathbb{Z}$ such that

$$\mu(A \cap F^{-n}(A) \cap \{x \in X : |S_n \psi(x)| \geq N\}) > 0.$$

Denote by $\overline{E}(\Psi)$ the set of essential values of the cocycle Ψ in $\overline{\mathbb{Z}}$. One of the applications of essential values is to the study of *skew-products* of dynamical systems.

Definition 2.13. Let (X, \mathcal{B}, μ) be a probability measure space, and $F : X \rightarrow X$ be a measure-preserving transformation. Let ν be the counting measure on \mathbb{Z} . Consider a measure space $(X \times \mathbb{Z}, \mathcal{B} \times 2^{\mathbb{Z}}, \mu \otimes \nu)$, let $\psi : X \rightarrow \mathbb{Z}$ be a skew-function, and $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ be the associated cocycle, as in (2.10). Then the skew-product of F is the transformation

$$(2.12) \quad T : X \times \mathbb{Z} \rightarrow X \times \mathbb{Z}, \quad (x, n) \mapsto (F(x), n + \psi(x)),$$

or, alternatively, denoting $T^m = T \circ \dots \circ T$,

$$(2.13) \quad T^m : X \times \mathbb{Z} \rightarrow X \times \mathbb{Z}, \quad (x, n) \mapsto (F^m(x), n + \Psi(x, m)).$$

Remark 2.14. Formula (2.13) explicitly defines the skew-product of the action of any element m of the acting group \mathbb{Z} , while formula (2.12) only specifies the skew-product of the action of a generator of \mathbb{Z} . Both definitions can be found in the literature. Formula (2.13) is suited to define skew-products of actions of finitely generated groups, which may have more than one generator.

We recall the following properties of essential values from [25, Theorem 3.9, Propositions 3.14 and 3.15, Corollary 5.4].

Theorem 2.15. Let $F : X \rightarrow X$ be an ergodic transformation on the probability space (X, \mathcal{B}, μ) , and let $\Psi : X \times \mathbb{Z} \rightarrow \mathbb{Z}$ be a cocycle for T . Then the following is true:

- (1) $\overline{E}(\Psi)$ is a closed non-empty subset of $\overline{\mathbb{Z}}$, and $E(\Psi) = \overline{E}(\Psi) \cap \mathbb{Z}$ is a subgroup of \mathbb{Z} .
- (2) $E(\Psi) = \mathbb{Z}$ if and only if the skew-product of F defined in Definition 2.13 is ergodic.
- (3) Ψ is a coboundary if and only if $\overline{E}(\Psi) = \{0\}$, with 0 being a non-trivial essential value.
- (4) Ψ is transient if and only if $\overline{E}(\Psi) = \{0, \infty\}$, with 0 being a trivial essential value.

2.6. Coboundaries for rotated odometers. In the setting of rotated odometers, one easy sufficient condition for the cocycle Ψ to be a coboundary is given in Lemma 2.16 below.

The weights of substitution words are defined in Definition 2.7. In the lemma, we denote by \mathcal{A} the alphabet with the doubled middle symbol, see Section 2.4.

Lemma 2.16. Let (I, F_π) be a stationary covering rotated odometer with the associated substitution χ . Let ψ be the skew-function defined in (1.2), and let $\Psi : I \times \mathbb{Z} \rightarrow \mathbb{Z}$ be the associated cocycle. If $\psi(\chi(a)) = 0$ for each $a \in \mathcal{A}$, then 0 is the only non-trivial essential value and Ψ is a coboundary.

This result is well-known, cf. [25]. In the proof below, we just specify the coboundary.

Proof. As discussed in Section 2.2, the orbits of (I, F_π) are coded by sequences of letters in \mathcal{A} , and there is the smallest subshift (X_χ, σ) such that every subword W that it contains is a subword of an iterated substitution word.

By (2.9), we need to find ϕ such that the coboundary equation (2.9) holds. Let L_1 be a section as in Proposition 2.3, and $L_{1,a}$, $a \in \mathcal{A}$, be the sets in the partition of L_k . For $x \in L_{1,a}$, $a \in \mathcal{A}$, the itinerary of the orbit of x for the first $m = |\chi(a)|$ steps is given by the substitution word. Let $\chi(a) = w_0 \dots w_{m-1}$, and set $\phi(x) = 0$ for all $x \in L_{1,a}$. Then for $1 \leq n \leq m$, all $x \in L_{1,a}$ and all $a \in \mathcal{A}$ define

$$(2.14) \quad \phi(F_\pi^n(x)) = \sum_{j=0}^{n-1} \psi(F_\pi^j(x)) = \sum_{j=0}^{n-1} \psi(w_j).$$

Since χ is covering, ϕ is now defined on I . Since $\psi(\chi(a)) = 0$ for all $a \in \mathcal{A}$, the coboundary equation extends for all $x \in I$ and all $n \in \mathbb{Z}$ by the same formula (2.14). \square

3. DISCREPANCY AND DIFFUSION COEFFICIENTS

In this section we compute the upper bounds on the discrepancy and the diffusion coefficient of a stationary covering rotated odometer (I, F_π, Leb) .

3.1. Discrepancy. Recall (see for instance [20, Chapter 1, Sections 1 and 5]), that a sequence $(x_n)_{n \geq 1} \subset [0, 1]$ is *uniformly distributed* if

$$\lim_{n \rightarrow \infty} \frac{1}{n} \#\{1 \leq j \leq n : x_j \in J\} = |J|$$

for each interval $J \subset [0, 1]$ and *well-distributed* if, for each interval $J \subset [0, 1]$,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \#\{k+1 \leq j \leq k+n : x_j \in J\} = |J|, \quad \text{uniformly in } k.$$

By the Birkhoff ergodic theorem, if (X, F, μ) is ergodic, then for μ -almost all $x \in X$ the orbit $\text{orb}(x)$ is uniformly distributed.

The *discrepancy* of $(x_n)_{n \geq 1}$ is a measure of how far the sequence deviates from the exact averages. More precisely:

Definition 3.1. *Let $(x_n)_{n \geq 1} \subset [0, 1]$ be a sequence of real numbers. Then the number*

$$\mathfrak{D}_R = \sup_{0 \leq a \leq b \leq 1} \left| \frac{1}{R} \#\{1 \leq j \leq R : x_j \in [a, b)\} - (b - a) \right|.$$

is called the discrepancy of $(x_n)_{n \geq 1}$.

One can also define

$$(3.1) \quad \mathfrak{D}_R^* = \sup_k \sup_{0 \leq a \leq b \leq 1} \left| \frac{1}{R} \#\{k+1 \leq j \leq k+R : x_j \in [a, b)\} - (b - a) \right|.$$

Clearly $\mathfrak{D}_R \leq \mathfrak{D}_R^*$. Note that Definition 3.1 does not require the sequence $(x_n)_{n \geq 1}$ to be uniformly (or well-) distributed.

A definition of discrepancy for substitution shifts, using cylinder sets instead of intervals, was used by Adamczewski [1], in his study of discrepancies of primitive substitutions. Adamczewski's results [1] were extended to S -adic transformations in [4]. For many rotated

odometers the substitution matrices of the associated shifts are not primitive, therefore, we cannot use the results of [1, 4]. In Theorem 1.11 below we obtain an estimate on the discrepancy of the sequence given by the orbit of $x \in I$ in the case when the associated substitution is covering, which means that Lebesgue measure is ergodic. Theorem 3.2 below is a restatement of Theorem 1.11 of the introduction, added here for the convenience of the reader.

Theorem 3.2. *Suppose that F_π is covering and stationary, the associated matrix is diagonalizable, and the largest eigenvalue λ_0 has multiplicity 1. Then for Lebesgue-a.e. x , there is $C = C_x$ such that the F_π -orbit of x has discrepancy $\mathfrak{D}_R \leq \mathfrak{D}_R^* \leq C_x \cdot R^{\gamma_0-1}$ for $\gamma_0 := \max \left\{ \frac{\log |\lambda_1|}{\log \lambda_0}, 0 \right\}$.*

If λ_0 has algebraic multiplicity ≥ 2 , then the diffusion coefficient could be 1. The assumption that J is diagonalizable is for simplicity, mostly. However, if the Jordan block associated to λ_1 is nontrivial, then we would need logarithmic correction factors as Adamczewski obtains for primitive substitution matrices, [1]. We have no examples where λ_1 has a non-trivial Jordan block, or λ_0 algebraic multiplicity ≥ 2 .

If F_π is uniquely ergodic, then C_x in the above theorem can be taken uniformly over all $x \in [0, 1)$, but no uniformity can be expected if there is a proper minimal subsystem, as for instance in Example 5.5, because there the \mathbb{Z} -extension of the minimal part is transient, with diffusion coefficient $1 > \gamma_0 = 0.694\dots$. The points in the minimal part are not Lebesgue typical, so there is no contradiction to Theorem 3.2.

Proof. In this proof, we work with the rotated odometer (I, F_π, Leb) , and so we do not have to split the middle interval of the partition $\mathcal{P}_{0,q}$ for q odd.

Let $M = UJU^{-1}$ be the Jordan decomposition of M , where J is such that the eigenvalues on the diagonal are in decreasing order, i.e., $\lambda_0 > |\lambda_1| \geq \dots \geq |\lambda_{q-1}|$. Then the first row ℓ_0 of U^{-1} , which is the left eigenvector associated to $\lambda_0 = 2^N$, is constant, because χ is covering. Assume it is scaled so that $(1, \dots, 1)$ is this row. Then for every symbol $a \in \mathcal{A} = \{0, \dots, q-1\}$, the abelianization of $\chi^n(a)$ is

$$(3.2) \quad V_n := (0, \dots, \underbrace{1}_{\text{position } a}, \dots, 0)UJ^nU^{-1} = \underbrace{U_a}_{a\text{-th row}} J^nU^{-1} = \sum_{j=0}^{q-1} U_{a,j} \lambda_j^n \ell_j,$$

where the last equality follows because J is a diagonal matrix. The length of the substitution word obtained from $a \in \mathcal{A}$ after n -th application of the substitution rule is

$$(3.3) \quad |\chi^n(a)| = \left\| \sum_{j=0}^{q-1} U_{a,j} \lambda_j^n \ell_j \right\|_1 \\ = \left\| U_{a,0} q 2^{nN} \left(1 + \sum_{j=1}^{q-1} \frac{U_{a,j}}{U_{a,0}} \left(\frac{\lambda_j}{\lambda_0} \right)^n \frac{\ell_j}{q} \right) \right\|_1 \sim |U_{a,0} q 2^{nN}|,$$

where at the last step we use that $\lambda_0 = 2^N > |\lambda_j|$, $1 \leq j \leq q-1$, and so all summands in the sum in the brackets are negligible for n large.

We now compute \mathfrak{D}_R^* , see (3.1), for the orbit of a Lebesgue typical point in I .

Recall that, for $k \geq 0$, we have the partition $\mathcal{P}_{kN,q}$ of the unit interval into half-open subintervals of length $(q2^{kN})^{-1}$, where N is defined in (2.3). We call an interval G a $q2^{kN}$ -adic interval if $G \in \mathcal{P}_{kN,q}$. Recall that for each $k \geq 1$, L_k is the union of the first q subintervals in the partition $\mathcal{P}_{kN,q}$. Denote by $L_{k,a}$, $0 \leq a \leq q-1$, these subintervals.

Recall from Proposition 2.3 that, associated to the rotated odometer F_π , there is a sequence of substitutions $(\chi_j)_{j \geq 1}$ with alphabet \mathcal{A} . Namely, the sets $L_{k,a}$, $0 \leq a \leq q-1$, form a coding partition $\mathcal{P}_{kN,q}^{\text{cod}}$, which codes the orbits of the $q2^{(k+1)N}$ -adic intervals in the partition $\mathcal{P}_{(k+1)N,q}^{\text{cod}}$. That is, one defines a substitution $\chi_{k+1} : \mathcal{A} \rightarrow \mathcal{A}^*$, where the substitution word $\chi_{k+1}(a)$ records the order in which the sets in $\mathcal{P}_{kN,q}^{\text{cod}}$ are visited by the orbit of $L_{k+1,a}$ under the first return map $F_{\pi,k} : L_k \rightarrow L_k$. For $i \geq 1$, the composition $\chi_{k+i} \circ \cdots \circ \chi_{k+1}(a)$ codes the orbit of $L_{k+i,a}$ with respect to $\mathcal{P}_{kN,q}$, and the orbit of $L_{k+i,a}$ with respect to $\mathcal{P}_{0,q}$ is coded by the composition $\chi_{k+i} \circ \cdots \circ \chi_1$ of substitutions.

The assumption of the theorem is that $(\chi_i)_{i \geq 1}$ is stationary, i.e., there is a substitution $\chi : \mathcal{A} \rightarrow \mathcal{A}^*$ such that $\chi_i = \chi$ for all $i \geq 1$. Then, given a $q2^{(k+n)N}$ -adic interval $L_{k+n,a}$, $0 \leq a \leq q-1$, the coding of its orbit relative to the partition $\mathcal{P}_{kN,q}$ is given by the substitution word $\chi^n(a)$.

We will compute (3.1) in two steps: first for any $q2^{kN}$ -adic interval G , and then for an interval of arbitrary length.

So let G be a $q2^{kN}$ -adic interval, let $n \geq 1$ and let $y \in L_{k+n,a}$, $a \in \mathcal{A}$, be a Lebesgue typical point, i.e., the orbit $\text{orb}(y)$ returns to L_k infinitely often. The easiest case is when $G \in \mathcal{P}_{kN,q}$, then the number of visits of the orbit of y of length $R = |\chi^{k+n}(a)|$ to G is bounded above by $|\chi^n(a)|$. Note that this bound is not sharp: indeed, $|\chi^n(a)|$ counts visits to any $q2^{kN}$ -adic interval in L_k , not just G . Under assumptions of the theorem, this bound is true for *any* $q2^{kN}$ -adic interval G in $\mathcal{P}_{kN,q}$: indeed, since χ is covering, there exists a unique $a \in \mathcal{A}$ such that $G = F_\pi^t(L_{k,a})$ for some $t \in \mathbb{N}$. By construction, the number t is less than the first return time of $L_{k,a}$ to L_k , and G appears in the orbit of $L_{k,a}$, before it returns to L_k , only once. Then $\text{orb}(y)$ visits $L_{k,a}$ if and only if it visits G , which implies that the bound on the number of visits to $L_{k,a}$ also holds for G . Thus we obtain, for any $q2^{kN}$ -adic interval G , the following estimate.

Denote by $\mathfrak{D}_R^*(y, G)$ an element of the set over which the supremum is taken in (3.1). Then we have

$$\begin{aligned}
(3.4) \quad \mathfrak{D}_R^*(y, G) &:= \frac{1}{R} \#\{0 \leq j < R : F_\pi^j(y) \in G\} - |G| \leq \frac{|\chi^n(a)|}{|\chi^{k+n}(a)|} - |G| \\
&= \frac{1}{|\chi^{k+n}(a)|} \left(U_{a,0} \lambda_0^n + \sum_{j=1}^{q-1} U_{a,j} |\lambda_j|^n \|\ell_j\|_1 \right) - \frac{1}{q2^{kN}} \\
&= \left(\frac{U_{a,0} \lambda_0^n + \sum_{j=1}^{q-1} U_{a,j} |\lambda_j|^n \|\ell_j\|_1}{U_{a,0} \lambda_0^{k+n} + \sum_{j=1}^{q-1} U_{a,j} |\lambda_j|^{k+n} \|\ell_j\|_1} \right) - \frac{1}{q2^{kN}} \\
&= \frac{U_{a,0} \lambda_0^n}{q U_{a,0} \lambda_0^{k+n}} \left(\frac{1 + \sum_{j=1}^{q-1} \frac{U_{a,j}}{U_{a,0}} \left(\frac{|\lambda_j|}{\lambda_0} \right)^n \|\ell_j\|_1}{1 + \sum_{j=1}^{q-1} \frac{U_{a,j}}{U_{a,0}} \left(\frac{|\lambda_j|}{\lambda_0} \right)^{k+n} \frac{\|\ell_j\|_1}{q}} \right) - \frac{1}{q2^{kN}} \\
(3.5) \quad &\sim \frac{1}{q2^{kN}} \left(1 + \sum_{j=1}^{q-1} \frac{U_{a,j}}{U_{a,0}} \left(\frac{|\lambda_j|}{\lambda_0} \right)^n \|\ell_j\|_1 - \sum_{j=1}^{q-1} \frac{U_{a,j}}{U_{a,0}} \left(\frac{|\lambda_j|}{\lambda_0} \right)^{k+n} \frac{\|\ell_j\|_1}{q} \right) - \frac{1}{q2^{kN}} \\
(3.6) \quad &\leq C |G| \left(\frac{|\lambda_1|}{\lambda_0} \right)^n \leq C' |G| R^{\gamma_0-1} < C' R^{\gamma_0-1},
\end{aligned}$$

for some $C, C' > 0$ independent of a . Here we use (3.2) to obtain (3.4), the approximation $\frac{1}{1+x} \sim 1 - x$ for small x and the fact that $\left(\frac{|\lambda_j|}{\lambda_0} \right)^n \rightarrow_n 0$ to obtain (3.5), and the fact that λ_1 dominates the eigenvalues λ_j , $j = 2, \dots, q-1$, in the absolute value, to obtain (3.6).

Now consider the orbit of arbitrary length R . As G is $q2^{kN}$ -adic, it suffices to take $R \geq \max_{a \in \mathcal{A}} |\chi^k(a)|$.

Let $u = u_1 \dots u_R$ be a subword of the infinite sequence in X_χ which codes the orbit of a Lebesgue typical point $y \in I$ as in Lemma 2.6, i.e., $u_\ell \in \mathcal{A}$ for each $1 \leq \ell \leq R$. This corresponds to an orbit of length R for a point $y' \in \text{orb}(y)$. By abuse of notation, and to distinguish with the case of orbits of specific length above, denote $\mathfrak{D}_R^*(u) := \frac{1}{R} \#\{1 \leq j \leq R : F_\pi(y) \in G\} - |G|$.

Take $n \geq 0$ maximal such that there is a letter $a \in \mathcal{A}$ such that $\chi^{k+n}(a)$ is a subword of u . Once this n has been found, we can take a block $B_0 \in \mathcal{A}^*$ maximal such that $\chi^{k+n}(B_0)$ is a subword of u .

Next find the $s \in \mathbb{N}$ minimal and blocks $B_{\pm 1}, B_{\pm 2}, \dots, B_{\pm s} \in \mathcal{A}^*$ maximal such that u is a subword of

$$(3.7) \quad v := \chi^{k+n-s}(B_{-s}) \dots \chi^{k+n-1}(B_{-1}) \chi^{k+n}(B_0) \chi^{k+n-1}(B_1) \dots \chi^{k+n-s}(B_s).$$

Here we allow the blocks $B_{\pm i}$, $1 \leq i \leq s$, to be empty, for the sake of convenience of notation, so that we could have a formula symmetric in s . Also, we choose $B_{\pm i}$ recursively, i.e., $B_{\pm i}$ can only be chosen after $B_{\pm(i-1)}$ maximal are found, for $1 \leq i \leq s$.

This procedure ensures that the lengths $b_i := |\chi^{k+n-|i|}(B_i)|$, $1 \leq |i| \leq s$, are bounded. Indeed, note that the lengths $|B_{\pm i}|$ are bounded above by $H = \max\{|\chi(a)| : a \in \mathcal{A}\}$, the length of the longest word of the substitution. Indeed, let B_{-i} be the first non-empty block such that $\chi^{k+n-i}(B_{-i})$ is adjacent to $\chi^{k+n}(B_0)$ on the left. Then $\chi^{k+n-i}(B_{-i})$ is contained

in a longer substitution word $\chi^{k+n}(a)$, for some $a \in \mathcal{A}$, and so the block B_{-i} must be a proper subword of the substitution word $\chi^i(a)$. Moreover, if $c \in \mathcal{A}$ is the rightmost letter in the subword $\chi^{i-1}(a)$, then B_{-i} is a proper subword of $\chi(c)$, since otherwise we would have that $\chi^{k+n-i+1}(c)$ is a subword of u adjacent to $\chi^{k+n}(B_0)$ on the left. This means that either $B_{-i+1} = c$ (if $i \geq 2$), or the block B_0 can be increased by adding c on the left. Both contradict the maximality of $B_{\pm i}$, for $0 \leq i \leq s$. The argument proceeds by induction on i . Next, since there is only a finite number of options for B_i , the lengths b_i are bounded. Namely, $b_i < \max\{|\chi^{k+n-|i|+1}(a)| : a \in \mathcal{A}\}$.

Then there is $C > 0$ and $\beta \in (0, 1)$, such that, independently of n and s , we have $b_i \leq C\beta^{|i|}b_0$ for all $-s \leq i \leq s$. In particular,

$$|u| \leq |v| \leq \frac{2C}{1-\beta}b_0.$$

The estimate (3.6) applies to b_i , $-s \leq i \leq s$, so there exists $C' \geq 1$ such that

$$b_i \mathfrak{D}_{b_i}^*(\chi^{k+n-|i|}(B_i)) \leq C' b_i^{\gamma_0} \leq C' C^{\gamma_0} \beta^{\gamma_0 |i|} b_0^{\gamma_0}.$$

Hence

$$\mathfrak{D}_R^*(u) \leq \frac{1}{R} \sum_{i=-s}^s b_i \mathfrak{D}_{b_i}^*(\chi^{k+n-|i|}(B_i)) \leq \frac{1}{R} \frac{2C' C_0^{\gamma_0}}{1-\beta^{\gamma_0}} b_0^{\gamma_0} \leq \frac{2C' C_0^{\gamma_0}}{1-\beta^{\gamma_0}} R^{\gamma_0-1},$$

which proves the theorem for any $q2^{kN}$ -adic interval, for $k \geq 1$.

Now, let G be any interval in I , and let $R = |\chi^{k+n}(a)|$, for $n \geq 1$ and $a \in \mathcal{A}$. Note that G can be represented as a (possibly infinite) union of $q2^{kN}$ -adic intervals, i.e., there are non-intersecting intervals K_1, K_2, \dots , such that $G = \bigcup_{t \geq 1} K_t$. Each K_t has length $c_t = (q2^{ktN})^{-1}$ for some $k_t \in \mathbb{N}$, and $|G| = \sum_{t \geq 1} c_t$. Then using (3.6) we obtain

$$\begin{aligned} \mathfrak{D}_R^*(y, G) &= \frac{1}{R} (\{0 \leq j \leq R : F_\pi^j(y) \in G\} - R|G|) \\ &= \frac{1}{R} \sum_{t \geq 1} (\{0 \leq j \leq R : F_\pi^j(y) \in K_t\} - R c_t) = \frac{1}{R} \sum_{t \geq 1} R \mathfrak{D}_R^*(y, K_t) \\ (3.8) \quad &\leq \sum_{t \geq 1} C' c_t R^{\gamma_0-1} = C' |G| R^{\gamma_0-1} \leq C' R^{\gamma_0-1}, \end{aligned}$$

since $|G| \leq 1$. The proof for arbitrary $R > 0$ proceeds similarly to the case of the $q2^{kN}$ -adic interval G , with (3.8) instead of (3.6). This concludes the proof of the theorem. \square

3.2. Diffusion coefficient. If the cocycle Ψ defined by the skew-function ψ is a coboundary, then all orbits are bounded, and hence the diffusion coefficient is 0. For the other case, we compute the upper bound on the diffusion coefficient in Theorem 1.10. Recall that we assume that the largest eigenvalue λ_0 has multiplicity 1, i.e., we assume that $\lambda_0 > |\lambda_1| \geq \dots \geq |\lambda_q|$. We also assume for the second largest eigenvalue that $|\lambda_1| > 1$. This holds, in particular, if the matrix M of the substitution χ has at least two non-zero diagonal blocks in the Frobenius form, see Section 2.3. Theorem 3.3 below is a restatement of Theorem 1.10 of the introduction, added here for the convenience of the reader.

Theorem 3.3. *If the substitution χ is covering and stationary, the associated matrix M is diagonalizable, the largest eigenvalue λ_0 has multiplicity 1, and the second largest eigenvalue satisfies $|\lambda_1| > 1$, then for the corresponding \mathbb{Z} -extension, the diffusion coefficients satisfy*

$$\gamma(x) \leq \gamma_0 := \frac{\log |\lambda_1|}{\log \lambda_0} \quad \text{for Leb-a.e. } x.$$

The assumptions that M is diagonalizable and that $|\lambda| \neq 1$ are not essential. Without them, there may appear extra logarithmic terms in the diffusion.

Proof. In this proof, we work with the skew-product dynamical system (1.3) over a rotated odometer (I, F_π, Leb) . Therefore, if q is odd, we split the middle state $m := \lfloor q/2 \rfloor$ in two states m^+ and m^- , according to the value of the skew-function, see Section 2.4 for details. The substitution and the associated matrix are still called χ and M . Recall that, as a consequence of the Perron-Frobenius Theorem, the left eigenvector $\ell_0 = (1, \dots, 1, \frac{1}{2}, \frac{1}{2}, 1, \dots, 1)$, where $\frac{1}{2}$ correspond to the split symbol.

Let ℓ_j , $j = 1, \dots, q-1$, be the left eigenvectors associated to eigenvalues λ_j .

We note that, in the definition of the diffusion coefficient (1.4) by (2.13) we have for $x \in I$

$$\xi(T_\pi^n(x, k)) - k = \xi(F_\pi^n(x), k + \Psi(x, n)) - k = \Psi(x, n) = \sum_{i=0}^{n-1} \psi(F_\pi^i(x)),$$

where Ψ is the cocycle associated to the skew-function ψ . Using the coding of orbits, the piece $\{x, F_\pi(x), \dots, F_\pi^{n-1}(x)\}$ is determined by a substitution word $W = w_1 \dots w_n$, and the corresponding displacement in the skew-product system is given by

$$\psi(W) := \sum_{i=1}^n \psi(w_i),$$

where we compute the weights $\psi(w_i)$ of symbols as in Definition 2.7.

Now we use a similar construction for W as was applied in the proof of Theorem 1.11 to the word u . Namely, take $s \geq 0$ maximal such that there is a letter $a \in \mathcal{A}$ such that $\chi^s(a)$ is a subword of W . Once such an s has been found, we can take a block $B_0 \in \mathcal{A}^*$ maximal such that $\chi^s(B_0)$ is a subword of W . Next find the blocks $B_{\pm 1}, B_{\pm 2}, \dots, B_{\pm s} \in \mathcal{A}^*$, possibly empty, such that

$$(3.9) \quad W := B_{-s} \chi(B_{-s+1}) \cdots \chi^{s-1}(B_{-1}) \chi^s(B_0) \chi^{s-1}(B_1) \cdots \chi(B_{s-1}) B_s.$$

Let $B_i = b_{i,1} \cdots b_{i,t_i}$ for $-s \leq i \leq s$, and note that $t_i < H = \max\{|\chi(a)| : a \in \mathcal{A}\}$ by the argument in the paragraph below the paragraph containing formula (3.7).

Let $\vec{u}_i = (u_{i,0}, \dots, u_{i,q})$ be the row vector with $u_{i,a} = \#\{1 \leq k \leq t_i : b_{i,k} = a\}$, $a \in \mathcal{A}$, i.e., the a -th components of \vec{u}_i is the number of occurrences of the letter $a \in \mathcal{A}$ in the block B_i . Take coefficients $c_{i,j} \in \mathbb{R}$ such that $\vec{u}_i = \sum_{j=0}^q c_{i,j} \ell_j$, and note that, since there is a finite number of subwords of length at most H in the alphabet \mathcal{A} , $E_j = \sup_i |c_{i,j}| < \infty$, $0 \leq j \leq q$, is independent of W .

Given a vector \vec{u}_i , we can compute its weight as the inner product \cdot of \vec{u}_i with the vector $(1, \dots, 1, -1, \dots, -1)$, which assigns weights to the states of the partition $\mathcal{P}_{0,q}$ (with the

middle interval split into 2 for q odd), with the middle interval split in the case of odd q ; in particular, we have

$$\psi(\ell_0) = \ell_0 \cdot (1, \dots, 1, -1 \cdots -1) = 0$$

for the left eigenvector associated to the largest eigenvalue λ_0 . Next, we estimate the absolute value of the weight of the piece of the orbit corresponding to each block B_i , that is,

$$\begin{aligned} |\psi(\chi^{s-|i|}(B_i))| &= \left| \left(\vec{u}_i M^{s-|i|} \right) \cdot (1, \dots, 1, -1 \dots, -1) \right| \\ &= \left| \left(\sum_{j=0}^q c_{i,j} \lambda_j^{s-|i|} \ell_j \right) \cdot (1, \dots, 1, -1 \dots -1) \right| \leq E |\lambda_1|^{s-|i|}, \end{aligned}$$

because $|\lambda_1| \geq |\lambda_j|$ for $2 \leq j \leq q$, and $\psi(\ell_0) = 0$. Here the constant $E \geq \sup\{E_j |\psi(\ell_j)| : 1 \leq j \leq q\}$.

Then, using (3.9) and by assumption $|\lambda_1| > 1$ we obtain

$$|\psi(W)| \leq \sum_{i=-s}^s |\psi(\chi^{s-|i|}(B_i))| \leq E \sum_{i=-s}^s |\lambda_1|^{s-|i|} \leq \frac{2E}{1 - 1/|\lambda_1|} |\lambda_1|^s.$$

Note that for $|B_0| = \|\vec{u}_0\|_1 = \|\sum_{j=0}^q c_{0,j} \ell_j\|_1$, a computation similar to (3.3) shows that

$$|\chi^s(B_0)| \sim |c_{0,0} \lambda_0^s|,$$

so, in particular, since $|\chi^s(B_0)|$ increases with s , $|c_{0,0}| \neq 0$. Since there is at most a finite number of words of length at most H , there exists a constant $E' > 0$ such that

$$|W| \geq |\chi^s(B_0)| \geq E' |c_{0,0}| \lambda_0^s.$$

Taking into account that $\lambda_0^{\gamma_0} = |\lambda_1|$, we obtain

$$|\psi(W)| \leq \frac{2E}{E'^{\gamma_0}} \frac{|\lambda_1|}{|\lambda_1| - 1} \frac{1}{|c_{0,0}|^{\gamma_0}} |W|^{\gamma_0},$$

as claimed. \square

Remark 3.4. If a substitution is not covering, but $\psi(\ell_\mu) = 0$, where μ is an ergodic measure, and ℓ_μ is the left eigenvector associated with the largest (in absolute value) eigenvalue ℓ_μ , then the same proof applies.

If a rotated odometer is not uniquely ergodic, then by the discussion in Section 2.3, the substitution M can be put into the Frobenius form M_F , where the first diagonal block D_1 corresponds to the minimal subsystem. Let μ be the measure supported on the minimal subset of (I, F_π) , and let λ_μ and ℓ_μ denote the leading eigenvalue of D_1 and its corresponding left eigenvector. If $\psi(\ell_\mu) = 0$, then a proof similar to that of Theorem 3.3 applies, and one obtains that the diffusion coefficient is bounded above by $\max\left\{\frac{\log |\lambda'_\mu|}{\log |\lambda_\mu|}\right\}$, where λ'_μ is the second largest eigenvalue of D_1 .

However, so far we haven't found any example where both $D_1 \neq M$ and $\psi(\ell_\mu) = 0$. Instead, in all our examples that are not uniquely ergodic, the minimal subsystem becomes transient on the skew-product, with diffusion coefficient $\gamma = 1$, because

$$0 < \frac{1}{C} |\psi(\ell_\mu)| \leq \frac{|\xi(T_\pi^n(x, k)) - k|}{n} \leq C |\psi(\ell_\mu)| < \infty \quad \mu\text{-a.e.},$$

for $C = \frac{\max\{U_{a,\mu} : a \in \mathcal{A}\}}{\min\{U_{a,\mu} : U_{a,\mu} > 0\}}$, and where $\xi : I \times \mathbb{Z} \rightarrow \mathbb{Z}$ is the projection.

4. RECURRENCE AND NON-ERGODICITY OF SKEW-PRODUCTS

In this section, we study the ergodic properties of the skew-product (1.3). Recall from Section 2 that a rotated odometer (I, F_π) , where π is a permutation of q symbols, admits at most q invariant ergodic measures, and Lebesgue measure Leb on I is ergodic if and only if (I, F_π) is covering, that is, $F_\pi : I \rightarrow I$ has no periodic points. In this section, we concentrate mostly on the study of skew-products on $I \times \mathbb{Z}$ with respect to the invariant measure $\text{Leb} \otimes \nu$ (where ν is counting measure on \mathbb{Z}), using theory of essential values, see Section 2.5 or [22, 25].

4.1. Recurrence. We assume that a rotated odometer (I, F_π) is covering, that is, Leb is an ergodic invariant measure. The following result goes back at least to Atkinson [2], but is used also in [17, 9].

Theorem 4.1. *Let (X, \mathcal{B}, μ, F) be an ergodic transformation on a probability space. Then the skew-function $\psi : X \rightarrow \mathbb{Z}$ has integral $\int_X \psi d\mu = 0$ if and only if the associated cocycle Ψ is recurrent, i.e., $\Psi(x, n) = \sum_{j=0}^{n-1} \psi \circ F^j(x) = 0$ infinitely often, for μ -a.e., $x \in X$.*

In comparison, for \mathbb{Z}^2 -extensions, as, for instance, in the wind-tree model, Atkinson's result fails. Nonetheless, Avila & Hubert [3] show that for typical (although not all) direction of the flow in a wind-tree model, the motion is recurrent.

As we are working with a \mathbb{Z} -extension, Atkinson's result is applicable and we obtain the following corollary. Recall that ν denotes counting measure on \mathbb{Z} .

Lemma 4.2. *Let (I, F_π, Leb) be a rotated odometer, and let $\psi : I \rightarrow \mathbb{Z}$ be the skew-function defined in (1.2). If χ is covering, then the skew-product $(I \times \mathbb{Z}, T_\pi, \text{Leb} \otimes \nu)$ is recurrent.*

If Lebesgue is not the only invariant measure, i.e., there is a proper minimal subsystem (I_{\min}, F_π, μ) , we obtain the result:

Lemma 4.3. *Let (I, F_π, μ) be a rotated odometer restricted to its (uniquely ergodic) minimal set, and let $\psi : I \rightarrow \mathbb{Z}$ be the skew-function defined in (1.2). Let ℓ denote the leading left eigenvector of the diagonal block D_1 in the Frobenius form (2.6) of M . If D_1 is diagonalizable, then the skew-product $(I \times \mathbb{Z}, T_\pi, \mu \otimes \nu)$ is recurrent if and only if $\psi(\ell) = 0$.*

Proof. The minimal part of the odometer is represented by upper left diagonal block D_1 in the Frobenius form. Using the permutation Π of (2.6) for the alphabet \mathcal{A} , we can assume that D_1 is the associated matrix of the substitution χ_1 acting on the first q_1 letters, describing the minimal part of F_π . Then D_1 is a $q_1 \times q_1$ -matrix, with eigenvalues $\lambda_0 > |\lambda_1| \geq \dots \geq \lambda_{q_1-1}$, and corresponding left (generalized) eigenvectors $\ell_0, \dots, \ell_{q_1-1}$, where we note that since D_1 is primitive, then by the Perron-Frobenius Theorem the multiplicity of λ_0 is 1. This substitution then acts on the alphabet $\mathcal{A}_1 = \{0, \dots, q_1 - 1\}$. Applying ψ to (3.3), we can write

$$\psi(\chi_1^n(0)) = \sum_{j=0}^{q_1-1} \psi(\ell_j) U_{0,j} \lambda_j^n \quad \text{and} \quad |\chi_1^n(0)| = \left\| \sum_{j=0}^{q_1-1} \ell_j U_{0,j} \lambda_j^n \right\|_1.$$

By minimality, $U_{0,0} \neq 0$. Since $\lambda_0 > |\lambda_j|$ for $1 \leq j < q_1$,

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \frac{\psi(\chi^n(0))}{|\chi^n(0)|} &= \lim_{n \rightarrow \infty} \frac{\sum_{j=0}^{q_1-1} \psi(\ell_j) U_{0,j} \lambda_j^n}{\left\| \sum_{j=0}^{q_1-1} \ell_j U_{0,j} \lambda_j^n \right\|_1} \\
 (4.1) \qquad \qquad \qquad &= \lim_{n \rightarrow \infty} \frac{\psi(\ell_0) \left(1 + \sum_{j=1}^{q_1-1} \frac{U_{0,j} \lambda_j^n}{U_{0,0} \lambda_0^n} \right)}{\|\ell_0\|_1 \left(1 + \sum_{j=1}^{q_1-1} \frac{\|\ell_j\|_1 U_{0,j} \lambda_j^n}{\|\ell_0\|_1 U_{0,0} \lambda_0^n} \right)} = \frac{\psi(\ell_0)}{\|\ell_0\|_1}.
 \end{aligned}$$

By Atkinson's result, $(I \times \mathbb{Z}, T_\pi, \mu \otimes \nu)$ is recurrent if and only if $\int \psi d\mu = 0$. But μ is the unique invariant probability measure of the minimal subsystem, so Oxtoby's Theorem and finally (4.1) give for the fixed point $\rho = \rho_0 \rho_1 \rho_2 \cdots = \lim_{n \rightarrow \infty} \chi^n(0)$:

$$\int \psi d\mu = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{j=0}^{N-1} \psi(\rho_j) = \lim_{n \rightarrow \infty} \frac{1}{|\chi^n(0)|} \sum_{j=0}^{|\chi^n(0)|-1} \psi(\rho_j) = \lim_{n \rightarrow \infty} \frac{\psi(\chi^n(0))}{|\chi^n(0)|} = \frac{\psi(\ell_0)}{\|\ell_0\|_1}.$$

Therefore $(I \times \mathbb{Z}, T_\pi, \mu \otimes \nu)$ is recurrent if and only if $\psi(\ell_0) = 0$. \square

This lemma applies also if χ is not covering (so Lebesgue is not ergodic), such as in Examples 5.9 and 5.10.

4.2. Non-ergodicity. In this section we establish two criteria for non-ergodicity of skew-products over rotated odometers, based on the properties of the associated substitutions. We recall from Theorem 2.15 or [25, Corollary 5.4] that the skew-product (1.3) is ergodic if and only if $E(\Psi) = \mathbb{Z}$.

Below we prove Theorem 1.8, restating it in a more technical way in Theorem 4.4 for the convenience of the reader. In this proof, we work with the skew-product dynamical system (1.3) over a rotated odometer (I, F_π, Leb) . Therefore, if q is odd, we split the middle state $m := \lfloor q/2 \rfloor$ in two states m^+ and m^- , according to the value of the skew-function, see Section 2.4 for details. The substitution and the associated matrix are still called χ and M .

Theorem 4.4. *Let $F_\pi : [0, 1) \rightarrow [0, 1)$ be a rotated odometer without intervals of periodic points, and suppose the associated substitution χ is stationary, so Lebesgue measure is ergodic. Let $\Psi : [0, 1) \times \mathbb{Z} \rightarrow \mathbb{Z}$ be the cocycle, and let $\mathbf{d} := \gcd\{\psi(\chi(a)) : a \in \mathcal{A}\}$. Then the subgroup $E(\Psi)$ of essential values is contained in $\mathbf{d}\mathbb{Z}$. In particular, if $\mathbf{d} > 1$, then 1 is not an essential value.*

Note that Theorem 4.4 does not imply that $E(\Psi)$ is equal to $\mathbf{d}\mathbb{Z}$; rather, it gives a necessary condition for an integer $e \in \mathbb{Z}$ to be an essential value of the cocycle Ψ . In particular, $E(\Psi)$ can still be a trivial subgroup of \mathbb{Z} , see Theorem 4.6 and examples after that theorem.

Proof. Recall from [25] that $e \in \mathbb{Z}$ is an essential value of Ψ if and only if for every Borel set $B \subset [0, 1)$ with $\text{Leb}(B) > 0$ we have

$$(4.2) \qquad \text{Leb} \left(\bigcup_{n \in \mathbb{Z}} (B \cap F_\pi^{-n}(B) \cap \{x \in B : \Psi(x, n) = e\}) \right) > 0.$$

We will show that there exists a set B such that if (4.2) holds, then $e \in \mathbf{d}\mathbb{Z}$, and then the statement of the theorem follows.

Take $B = L_k$ for some $k \geq 1$, where L_k is the k -th section. Recall that F_π is invertible except on the orbit of 0, and so $\text{Leb}(B \cap F_\pi^{-n}(B)) > 0$ for some $n > 0$ if and only if $\text{Leb}(B \cap F_\pi^n(B)) > 0$.

The section $B = L_k$ of length 2^{-kN} is subdivided into the intervals $L_{k+1,a}$, $0 \leq a \leq q-1$, of length $q^{-1}2^{-kN}$. The length of the substitution word $|\chi(a)|$ determines the first return time of $L_{k+1,a}$ to L_k under iterations of the rotated odometer map F_π , that is, we have

$$F_\pi^{|\chi(a)|}(L_{k+1,a}) \subset L_k, \quad \text{and} \quad F_\pi^s(L_{k+1,a}) \cap L_k = \emptyset \quad \text{for } 0 < s < |\chi(a)|.$$

We claim that the intersection

$$S_{k+1,a,b} = F_\pi^{|\chi(a)|}(L_{k+1,a}) \cap L_{k+1,b}, \quad b \in \mathcal{A},$$

is either empty, or is at most a finite union of half-open intervals. To see that recall that $F_\pi = \mathbf{a} \circ R_\pi$, where R_π is a finite IET, and \mathbf{a} is the von Neumann-Kakutani map. The restriction of each F_π^i , $0 \leq i < |\chi(a)|$, to $L_{k+1,a}$ is a translation; the interval $L_{k+1,a}$ is re-arranged under F_π^i (by the von Neumann-Kakutani part of the composition) only after its orbit reaches the set $R_\pi^{-1}([1 - 2^{-kN}, 1))$, which happens for $i = |\chi(a)| - 1$. We then have $F_\pi^{|\chi(a)|}(L_{k+1,a}) \subset L_k$.

Recall from (1.1) that I is subdivided into intervals I_ℓ , $\ell \geq 1$, of length $|I_\ell| = 2^{-\ell}$, on which \mathbf{a} is continuous, and which are re-arranged under \mathbf{a} in the opposite order. Since $q \neq 2^r$, $r \in \mathbb{N}$, the image $R_\pi \circ F_\pi^{|\chi(a)|-1}(L_{k+1,a})$ may be contained in the union of more than one interval I_ℓ , and there is a single letter $\iota \in \mathcal{A}$ such that $R_\pi \circ F_\pi^{|\chi(a)|-1}(L_{k+1,a}) \supset \bigcup_{\ell > \ell_\iota} I_\ell$, where $\ell_\iota = 2^{kN+1}$. Then $\mathbf{a}(I_\ell) \subset L_{k+1,0}$ for $\ell > \ell_\iota$, and, for $\ell \leq \ell_\iota$, I_ℓ is mapped into a finite union of the sets $L_{k+1,b}$, $b \in \mathcal{A}$. It follows that either $S_{k+1,a,b} = \emptyset$, or $S_{k+1,a,b}$ is a finite union of half-open intervals.

Next, note that $L_k \cap F_\pi^n(L_k)$ is non-empty if and only if one of $L_{k,a}$, $a \in \mathcal{A}$, is mapped into L_k by a power of F_π . That is, $L_k \cap F_\pi^n(L_k)$ is non-empty if and only if $n = \sum_{i=1}^t |\chi(a_i)|$ for some $t \geq 1$, and $a_i \in \mathcal{A}$, $1 \leq i \leq t$. Let $D \subset L_k$ be a subinterval for which $F_\pi^n(D) \subset L_k$; then the itinerary of every point $x \in D$ is described by the concatenated substitution word $W = \chi(a_1) \cdots \chi(a_t)$, and for this word we have

$$\psi(W) = \psi(\chi(a_1)) + \cdots + \psi(\chi(a_t)).$$

Since every weight $\psi(\chi(a_i))$ is a multiple of \mathbf{d} , for all $x \in D_r$ we have $\Psi(x, n) \in \mathbf{d}\mathbb{Z}$. Since this argument is true for any $n \in \mathbb{Z}$ such that $\text{Leb}(L_k \cap F_\pi^{-n}(L_k)) > 0$, we have that $E(\Psi) \subset \mathbf{d}\mathbb{Z}$. \square

Example 4.5. By Theorem 4.4 the skew-products over the following rotated odometers are not ergodic:

- (1) For $q = 3$ and $\pi = (012), (021)$ in Examples 5.1 and 5.2, and for $q = 9$ and $\pi = (1, 7, 4)(2, 5)(3, 6)$ in Example 5.3 the rotated odometers are uniquely ergodic with Lebesgue invariant measure. In these examples $\mathbf{d} := \gcd\{\psi(\chi(a)) : a \in \mathcal{A}\} = 2$, so $E(\Psi) \subset 2\mathbb{Z}$ and the skew-product $(I \times \mathbb{Z}, T_\pi, \text{Leb} \otimes \nu)$ is not ergodic.
- (2) For $q = 11$ in Example 5.7, the rotated odometer determined by the permutation $\pi = (0, 2, 7, 6, 5, 4, 3, 8, 10, 1, 9)$ is not uniquely ergodic. Here again $\mathbf{d} = \gcd\{\psi(\chi(a)) : a \in \mathcal{A}\} = 2$, so $E(\Psi) \subset 2\mathbb{Z}$ and the skew-product $(I \times \mathbb{Z}, T_\pi, \text{Leb} \otimes \nu)$ is not ergodic.

We now prove Theorem 1.9, restating it below in Theorem 4.6. Recall that a Pisot number is an algebraic integer > 1 such that its Galois conjugates are all strictly inside the unit circle. If λ_j is Pisot, this does not preclude that there are other eigenvalues outside the unit disc, even if the weight of the corresponding left eigenvector is non-zero. After all, the characteristic polynomial of M need not be irreducible.

In the proof below M is a $(q+1) \times (q+1)$ -matrix, i.e., M corresponds to the substitution χ with doubled middle symbol, see (2.4). The alphabet in Theorem 4.6 is given by $\mathcal{A} = \{0, 1, \dots, m^-, m^+, \dots, q-1\}$, i.e., $\#\mathcal{A} = q+1$, and the numbering in the sums below is from 0 to q .

Theorem 4.6. *Let F_π be stationary and covering, and assume the following:*

- (1) *for all eigenvalues λ_j of the associated matrix M with norm $|\lambda_j| \geq 1$, the left eigenvectors have weights $\psi(\ell_j) = 0$, with the exception of one, say, for $\lambda_c > 1$.*
- (2) *λ_c is Pisot, and the algebraic and geometric multiplicities of λ_c are equal.*

Then the essential values $\overline{E}(\Psi) = \{0, \infty\}$, with 0 a non-trivial essential value. In particular, Lebesgue measure on the staircase has infinitely many ergodic components.

Example 4.7. This theorem applies to permutations (012) and (021) for $q = 3$ in Examples 5.1 and 5.2, thanks to the fact that in these examples, all left eigenvectors to eigenvalues 1 have weight $\psi(\ell_j) = 0$. Why this latter property holds for all our examples is not clear to us, but see Lemma 4.8 for some discussion. A more interesting example is (1, 7, 4)(2, 5)(3, 6) in Example 5.3 because it has the second largest eigenvalue of multiplicity 2. The rotated odometer corresponding to the substitution (02431) for $q = 5$ in Example 5.5 has two ergodic invariant measures, and a non-integer Pisot eigenvalue. In all these examples Theorem 4.6 applies, and so $\overline{E}(\Psi) = \{0, \infty\}$, with non-trivial essential value 0.

Proof. Let $t \geq 1$ be the algebraic and geometric multiplicity of λ_c . By assumption, λ_c has t linearly independent left eigenvectors $\ell_c, \dots, \ell_{c+t-1}$. For the smallest $c' > c$ with $\psi(\ell_{c'}) \neq 0$, we have by assumption that $|\lambda_{c'}| < 1 < \lambda_c$. Apply the weight function ψ to (3.3) to obtain

$$\psi(\chi^n(b)) = \sum_{a=0}^q \psi(\ell_c) U_{b,a} (\lambda_a^n + \epsilon_a n \lambda_a^{n-1}) = \left(\sum_{j=0}^{t-1} \psi(\ell_{c+j}) U_{b,c+j} \right) \lambda_c^n + R(n),$$

where $U_{b,a}$ is the (b, a) -entry in the matrix U of some Jordan block decomposition $M = UJU^{-1}$, and $\epsilon_a = 1$ if the Jordan block of λ_a is non-diagonal, and $\epsilon_a = 0$ otherwise. Since λ_c is the only eigenvalue of absolute value ≥ 1 , whose left eigenvector has non-zero weight, only that term remains in the rightmost expression, and the remainder term $R(n)$ is exponentially small. In particular, since $\lambda_a < \lambda_{c'} < 1 < \lambda_c$ for all $a \geq c'$, there exists n_0 such that for all $n > n_0$ and all such a we have $n\lambda_a^{n-1} < \lambda_c^n$.

Now $\sum_{j=0}^{t-1} \psi(\ell_{c+j}) U_{b,c+j} \neq 0$ for at least one b , because otherwise the columns $c, \dots, c+t-1$ of U are linearly dependent, contradicting that U is invertible. This implies that $|\psi(\chi^n(b))| \rightarrow_{n \rightarrow \infty} \infty$, and some $(x, k) \in I \times \mathbb{Z}$ has an unbounded T_π -orbit. This shows that ψ is not a coboundary and $\infty \in \overline{E}(\Psi)$. Since χ is covering, Lemma 4.2 shows that $0 \in \overline{E}(\Psi)$. It remains to show that $\overline{E}(\Psi)$ contains no integer $e \neq 0$.

Let $H = \max_{a \in \mathcal{A}} |\chi(a)|$, and $E = \sup\{|U_{b,a} \psi(\ell_a)| : b, a \in \mathcal{A}\}$. The eigenvalue λ_c is algebraic of degree $d \leq q$, so the corresponding eigenvectors can be chosen in the field extension $\mathbb{Q}(\lambda_c)$,

and we can write, for each $0 \leq j \leq t-1$,

$$(4.3) \quad \psi(\ell_{c+j})U_{a,c+j} = \frac{1}{\Delta_{c+j}} \sum_{k=0}^{d-1} p_{k,a}^j \lambda_c^k$$

for integers $\Delta_{c+j} \neq 0$ and $p_{k,a}^j$. Let

$$\Delta_c = \prod_{j=0}^{t-1} \Delta_{c+j}, \quad \Delta_{c \div j} = \frac{\Delta_c}{\Delta_{c+j}}, \quad 0 \leq j \leq t-1.$$

Set $G_j = 2 \sum_{a \in \mathcal{A}} \sum_{k=0}^{d-1} |p_{k,a}^j| \in \mathbb{N}$, and

$$G = t \Delta_c \max\{G_j : 0 \leq j \leq t-1\}.$$

Since λ_c is Pisot, by the Garsia Separation Lemma [14, Lemma 1.51], there is $C = C(G) > 0$ such that for all $\alpha \in \mathbb{N}$ and all $\varepsilon_j \in \{-G, \dots, G\}$, $0 \leq j \leq \alpha$, we have

$$(4.4) \quad \left| \sum_{j=0}^{\alpha} \varepsilon_j \lambda_c^{-j} \right| \geq C \lambda_c^{-\alpha} \quad \text{or} \quad \left| \sum_{j=0}^{\alpha} \varepsilon_j \lambda_c^{-j} \right| = 0.$$

Take $k_0 \in \mathbb{N}$ so large that $k_0 > n_0$, and

$$(4.5) \quad |\lambda_c|^{k_0} > \frac{2\Delta_c|e|}{C} \quad \text{and} \quad |\lambda_{c'}|^{k_0} < \frac{1-|\lambda_{c'}|}{4HEq}.$$

Thus, for a fixed F_π , the constant k_0 depends only on the choice of $e \in \mathbb{Z}$.

Now let $A \subset [0, 1]$ be an arbitrary set of positive measure. To show that $e \neq 0$ is an essential value, for every such A we need to find an n such that

$$(4.6) \quad \text{Leb} \left(A \cap F_\pi^{-n}(A) \cap \left\{ x \in A : \sum_{m=0}^{n-1} \psi \circ F_\pi^m(x) = e \right\} \right) > 0.$$

In fact, it is not a restriction to assume that A is a $q2^{kN}$ -adic interval for some $k \in \mathbb{N}$, since every A would contain such an interval, for k large enough. We will prove that it is possible to choose A so that (4.6) does not hold for any $n \in \mathbb{N}$. This will show that e cannot be an essential value.

Since F_π is covering, there is a $q2^{kN}$ -adic subinterval A' of L_k such that $F_\pi^{k'}$ maps A' continuously onto A for some $k' \geq 0$. Pulling A back by $F_\pi^{-k'}$, (4.6) is equivalent to finding an n such that

$$\text{Leb} \left(A' \cap F_\pi^{-n}(A') \cap \left\{ x \in A' : \sum_{m=0}^{n-1} \psi \circ F_\pi^m(x) = e \right\} \right) > 0.$$

The return times to the set L_k satisfy $n = \sum_{i=1}^r |\chi^k(a_i)|$ for some $r \geq 1$ and $a_i \in \mathcal{A}$. We now assume that k is large enough (and, consequently, A is small enough) so that $\min\{|\chi^k(a)| : a \in \mathcal{A}\} > k_0$, where k_0 is the constant we chose in (4.5).

Since the rotated odometer (I, F_π, Leb) is ergodic, A contains a full measure set of points returning to A infinitely often. Let y be a Lebesgue typical point in A , and suppose $F^n(y) \in A$ for some n . By the choice of A we have $n > k_0$, and by the coding procedure in Proposition 2.3, the itinerary of y is coded by a word W in the alphabet \mathcal{A} of length n . Since F_π

acts by interval exchanges, there is a positive measure set $A_W \subset A$ (in fact, a subinterval) of points with the same itinerary as y .

Similarly to what we did in the proofs of Theorems 1.10 and 1.11, we can find (possibly empty) maximal blocks $B_i = b_{i,1} \dots b_{i,h_i}$, $-s \leq i \leq s$, of length $h_i \leq H$ such that

$$W = \chi^{k_0}(B_{-s})\chi^{k_0+1}(B_{-s+1}) \dots \chi^{k_0+s-1}(B_{-1})\chi^{k_0+s}(B_0)\chi^{k_0+s-1}(B_1) \dots \chi^{k_0+1}(B_{s-1})\chi^{k_0}(B_s).$$

Thus, for all $x \in A_W$,

$$\sum_{m=0}^{n-1} \psi \circ F_\pi^m(x) = \sum_{i=-s}^s \sum_{j=1}^{h_i} \sum_{a=c}^q \psi(\ell_a) U_{b_{i,j},a} (\lambda_a^{k_0+s-|i|} + \epsilon_a n \lambda_a^{k_0+s-|i|-1}) = S_1 + S_2$$

where

$$\begin{aligned} |S_1| &:= \left| \sum_{i=-s}^s \sum_{j=1}^{h_i} \sum_{a=c'}^q \psi(\ell_a) U_{b_{i,j},a} \left(\lambda_a^{k_0+s-|i|} + \epsilon_a n \lambda_a^{k_0+s-|i|-1} \right) \right| \\ &\leq 2HEq \sum_{i=-s}^s |\lambda_{c'}|^{k_0+s-|i|} \leq \frac{4HEq |\lambda_{c'}|^{k_0}}{1 - |\lambda_{c'}|} < 1, \end{aligned}$$

because $|\lambda_a| \leq |\lambda_{c'}| < 1$ for all $a < c'$, and

$$\begin{aligned} (4.7) \quad S_2 &:= \lambda_c^{k_0+s} \sum_{i=-s}^s \sum_{j=1}^{h_i} \sum_{z=0}^{t-1} \psi(\ell_{c+z}) U_{b_{i,j},c+z} \lambda_c^{-|i|} \\ &= \lambda_c^{k_0+s} \sum_{i=-s}^s \sum_{j=1}^{h_i} \left(\sum_{z=0}^{t-1} \frac{1}{\Delta_{c+z}} \sum_{k=0}^{d-1} p_{k,b_{i,j}}^z \lambda_c^{k-|i|} \right) \\ &= \frac{\lambda_c^{k_0+s+d}}{\Delta_c} \underbrace{\sum_{u=1}^{s+d} \sum_{z=0}^{t-1} \Delta_{c+z} \sum_{k=0}^{d-1} \left(\sum_{j=1}^{h_{u+k-d}} p_{k,b_{u+k-d},j}^z + \sum_{j=1}^{h_{-(u+k-d)}} p_{k,b_{-(u+k-d),j}^z} \right)}_{\text{integers in } \{-G, \dots, G\}} \lambda_c^{-u} \\ &= 0 \quad \text{or} \quad |S_2| \geq \frac{|\lambda_c|^{k_0+s+d}}{\Delta_c} C |\lambda_c|^{-s-d} \geq |\lambda_c|^{k_0} \frac{C}{\Delta_c}, \end{aligned}$$

depending on which possibility of the Garsia Separation Lemma holds. In the first case, $|S_1 + S_2| < 1$, and in the second case, $|S_1 + S_2| > -1 + 2|e| \geq |e|$. Thus we found a set for which (4.6) fails, which shows that $e \neq 0$ is not an essential value. \square

4.3. Left eigenvectors of zero weight. The above theorem requires, among other assumptions, that the eigenvalues $|\lambda_j| = 1$ have weight $\psi(\ell_j) = 0$. This looks an artificial assumption, but all the examples that we know (including the ones in the next section) have this property. As a partial explanation for this, we consider in the next lemma some situations where $\lambda_j \in \{0, -1, 1\}$ and $\psi(\ell_j) = 0$. The proof is straightforward.

Let $|\chi(a)|_b$ be the number of occurrences of letter b in $\chi(a)$. We denote by 1_a the vector with 1 in the a -th entry, and 0 in the other entries.

Lemma 4.8. *Let χ be a stationary substitution, and M the associated matrix.*

- (1) *If $\chi(a) = \chi(b)$ for some $a < b$, then $1_a - 1_b$ is a left eigenvector to eigenvalue 0.*
- (2) *If $|\chi(a)|_a = |\chi(b)|_a + 1$ and $|\chi(a)|_b = |\chi(b)|_b - 1$ for some $a < b$, and $|\chi(a)|_j = |\chi(b)|_j$ for $j \neq a, b$, then $1_a - 1_b$ is a left eigenvector to eigenvalue 1.*
- (3) *If $|\chi(a)|_a = |\chi(b)|_a - 1$ and $|\chi(a)|_b = |\chi(b)|_b + 1$ for some $a < b$, and $|\chi(a)|_j = |\chi(b)|_j$ for $j \neq a, b$, then $1_a - 1_b$ is a left eigenvector to eigenvalue -1 .*
- (4) *If $|\chi(a)|_a = |\chi(b)|_a + 1$ and $|\chi(a)|_b = |\chi(b)|_b - 1$ for some $a < b$ (like in (2)), $\chi(c) = \chi(d)$, $|\chi(a)|_c = |\chi(b)|_c + 1$ and $|\chi(a)|_d = |\chi(b)|_d - 1$ and $|\chi(a)|_j = |\chi(b)|_j$ for all other indices j , then $1_a - 1_b + 1_c - 1_d$ is a left eigenvector to eigenvalue 1.*

If additionally $0 \leq a < b \leq m^+$ or $m^- \leq a < b \leq q$, then the left eigenvectors mentioned in (1)-(3) have weight 0. A similar extended statement holds for case (4).

The conditions in this lemma may look cumbersome and arbitrary, but they do occur because of certain structure of the (stationary) rotated odometer F_π and its renormalization. Indeed, if $L_{1,a}$ and $L_{1,b}$ are $q2^N$ -adic subintervals of L_1 that have the same return time $n = |\chi(a)| = |\chi(b)|$ to L_1 , and F_π^j is an isometry on $L_{1,a} \cup L_{1,b}$, then if $\frac{a+1}{q}$ lies between $F_\pi^j(L_{1,a})$ and $F_\pi^j(L_{1,b})$ for some j , and no other $\frac{a'}{q}$, $a' \in \mathcal{A}$, has this property, then the condition of part (2) holds. Naturally, this is easiest if $b = a + 1$, and it applies to Examples 5.10, 5.11, 5.12, 5.13 (with $a = 0, b = 4$)

A similar phenomenon occurs in Example 5.5 with $a = 0, b = 3$, leading to case (3) of the lemma.

5. EXAMPLES

In the tables below we list the properties of several stationary (and one non-stationary) substitutions, whether they are covering or not, and if covering, then whether they are uniquely ergodic or not.

We give the characteristic polynomial and eigenvalues of the associated matrix M that emerges after doubling the middle symbol m to m^+ and m^- , see Section 2.4. The number of ergodic measures of the rotated odometers can easily be read off from the Frobenius form of M , see Section 2.3. The quantities γ_0 and $\overline{E}(\Psi)$ are the Lebesgue typical diffusion coefficient and the set of essential values.

In the last line we give the weights of those left eigenvectors of M that belong to non-zero eigenvalues, as weights of eigenvectors of the eigenvalue 0 don't affect anything. The scaling of the eigenvectors is arbitrary, as given by a program in Sage which we used to compute the examples. For proofs of our theorems it only matters if the weights are zero or non-zero. Non-rational values are marked by $*$, and we only give them up to at most two first digits. In the case of irrational eigenvalues it is usually clear which (quadratic or cubic) polynomial factors they are a root of. Values which are not reported are marked by $**$.

5.1. Uniquely ergodic rotated odometers.

Example 5.1.

$\pi = (012), \quad q = 3$ covering $\#\{\text{ergodic measures}\} = 1$ $\gamma_0 = \frac{1}{2}, \quad \overline{E}(\Psi) = \{0, \infty\}$	$M = \begin{pmatrix} 1 & 0 & 1 & 2 \\ 1 & 0 & 1 & 2 \\ 1 & 0 & 1 & 2 \\ 2 & 2 & 0 & 0 \end{pmatrix}$
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	$\begin{array}{c c c c} 0 & 1^+ & 1^- & 2 \\ \hline -2 & -2 & -2 & 4 \end{array}$
Characteristic polynomial	$(x - 4)(x + 2)x^2$
Eigenvalues Weights of left eigenvectors	$\begin{array}{c c c} 4 & -2 & 0(\times 2) \\ \hline 0 & -1 & ** \end{array}$

Example 5.2.

$\pi = (021), \quad q = 3$ covering $\#\{\text{ergodic measures}\} = 1$ $\gamma_0 = \frac{1}{2}, \quad \overline{E}(\Psi) = \{0, \infty\}$	$M = \begin{pmatrix} 2 & 2 & 2 & 4 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	$\begin{array}{c c c c} 0 & 1^+ & 1^- & 2 \\ \hline -2 & -2 & -2 & 4 \end{array}$
Characteristic polynomial	$(x - 4)(x + 2)x^2$
Eigenvalues Weights of left eigenvectors	$\begin{array}{c c c} 4 & -2 & 0(\times 2) \\ \hline 0 & -1 & ** \end{array}$

In both Examples 5.1 and 5.2, the rotated odometers are stationary and covering, and so all our theorems apply. The weights of all substitution words are even, thus Theorem 1.8 gives $E(\Psi) \subset 2\mathbb{Z}$ in both cases. By Theorem 1.9 we conclude that $\overline{E}(\Psi) = \{0, \infty\}$.

Example 5.3.

$\pi = (1, 7, 4)(2, 5)(3, 6)$ $q = 9$ covering $\#\{\text{ergodic measures}\} = 1$ $\gamma_0 = \frac{1}{2}$ $\overline{E}(\Psi) = \{0, \infty\}$	$M = \begin{pmatrix} 4 & 3 & 1 & 2 & 0 & 1 & 3 & 4 & 1 & 3 \\ 3 & 6 & 3 & 3 & 1 & 1 & 3 & 3 & 2 & 3 \\ 1 & 2 & 3 & 3 & 1 & 1 & 1 & 1 & 2 & 1 \\ 1 & 2 & 3 & 3 & 1 & 1 & 1 & 1 & 2 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 1 \\ 1 & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 2 & 1 \\ 3 & 3 & 2 & 1 & 1 & 0 & 4 & 3 & 1 & 4 \end{pmatrix}$																				
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	<table border="1"> <tbody> <tr> <td>0</td><td>1</td><td>2</td><td>3</td><td>4⁺</td><td>4⁻</td><td>5</td><td>6</td><td>7</td><td>8</td> </tr> <tr> <td>-2</td><td>4</td><td>4</td><td>4</td><td>-2</td><td>-2</td><td>-2</td><td>-2</td><td>-2</td><td>-2</td> </tr> </tbody> </table>	0	1	2	3	4 ⁺	4 ⁻	5	6	7	8	-2	4	4	4	-2	-2	-2	-2	-2	-2
0	1	2	3	4 ⁺	4 ⁻	5	6	7	8												
-2	4	4	4	-2	-2	-2	-2	-2	-2												
Characteristic polynomial	$(x - 16)(x - 4)^2(x - 1)^2x^5$																				
Eigenvalues Weights of left eigenvectors	<table border="1"> <tbody> <tr> <td>16</td><td>4($\times 2$)</td><td>1($\times 2$)</td><td>0($\times 5$)</td> </tr> <tr> <td>0</td><td>$-\frac{1}{2}, 1$</td><td>0, 0</td><td>**</td> </tr> </tbody> </table>	16	4($\times 2$)	1($\times 2$)	0($\times 5$)	0	$-\frac{1}{2}, 1$	0, 0	**												
16	4($\times 2$)	1($\times 2$)	0($\times 5$)																		
0	$-\frac{1}{2}, 1$	0, 0	**																		

In this example, the rotated odometer is stationary and covering, and so all our theorems apply. Since the weights of all substitution words are even, by Theorem 1.8 we have $E(\Psi) \subset 2\mathbb{Z}$. The second largest eigenvalue $\lambda_1 = 4$ has both the algebraic and the geometric multiplicity equal to 2. Thus Theorem 1.9 applies, and we conclude that $\overline{E}(\Psi) = \{0, \infty\}$.

5.2. A coboundary.

Example 5.4.

$\pi = (01234), \quad q = 5$ covering $\#\{\text{ergodic measures}\} = 2$ ψ is coboundary: $\gamma_0 = 0$ $\overline{E}(\Psi) = \{0\}$	$M = \begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 4 & 8 & 4 & 4 & 4 & 8 \end{pmatrix}$												
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	<table border="1"> <tbody> <tr> <td>0</td><td>1</td><td>2⁺</td><td>2⁻</td><td>3</td><td>4</td> </tr> <tr> <td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td> </tr> </tbody> </table>	0	1	2 ⁺	2 ⁻	3	4	0	0	0	0	0	0
0	1	2 ⁺	2 ⁻	3	4								
0	0	0	0	0	0								
Characteristic polynomial	$(x - 8)(x - 2)x^4$												
Eigenvalues Weights of left eigenvectors	<table border="1"> <tbody> <tr> <td>8</td><td>2</td><td>0($\times 4$)</td> </tr> <tr> <td>0</td><td>0</td><td>**</td> </tr> </tbody> </table>	8	2	0($\times 4$)	0	0	**						
8	2	0($\times 4$)											
0	0	**											

For $\pi = (01234)$, the cocycle Ψ is a coboundary because the weight of all substitution words is 0. Hence all orbits in $(I \times \mathbb{Z}, T_\pi, \text{Leb} \otimes \nu)$ are bounded. The minimal subset (I_{\min}, F_π)

corresponds to the symbols $\{0, 3\}$, and lifts to the suspension over a dyadic odometer in $(I \times \mathbb{Z}, T_\pi, \text{Leb} \otimes \nu)$, which has zero measure with respect to $\text{Leb} \otimes \nu$. Lebesgue itself is the only other ergodic F_π -measure.

Note that the weight of the left eigenvector associated to the second largest eigenvalue is 0, which shows again that the orbits of T_π stay within a bounded subset (compare with Theorem 1.9).

5.3. Non-uniquely ergodic covering substitutions.

Example 5.5.

$\pi = (02431), \quad q = 5$ covering $\#\{\text{ergodic measures}\} = 2$ $\gamma_0 = 0.694\dots$ $\overline{E}(\Psi) = \{0, \infty\}$	$M = \begin{pmatrix} 1 & 1 & 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 2 & 1 & 0 & 1 & 0 & 1 \\ 2 & 1 & 0 & 1 & 0 & 1 \\ 3 & 5 & 2 & 1 & 8 & 5 \\ 1 & 1 & 0 & 1 & 0 & 0 \end{pmatrix}$												
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	<table style="border-collapse: collapse; margin: auto;"> <tr> <td style="border-right: 1px solid black; padding: 2px 5px;">0</td> <td style="border-right: 1px solid black; padding: 2px 5px;">1</td> <td style="border-right: 1px solid black; padding: 2px 5px;">2⁺</td> <td style="border-right: 1px solid black; padding: 2px 5px;">2⁻</td> <td style="border-right: 1px solid black; padding: 2px 5px;">3</td> <td style="padding: 2px 5px;">4</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px 5px;">1</td> <td style="border-right: 1px solid black; padding: 2px 5px;">1</td> <td style="border-right: 1px solid black; padding: 2px 5px;">1</td> <td style="border-right: 1px solid black; padding: 2px 5px;">1</td> <td style="border-right: 1px solid black; padding: 2px 5px;">-4</td> <td style="padding: 2px 5px;">1</td> </tr> </table>	0	1	2 ⁺	2 ⁻	3	4	1	1	1	1	-4	1
0	1	2 ⁺	2 ⁻	3	4								
1	1	1	1	-4	1								
Characteristic polynomial	$(x - 8)(x + 1)^2(x - 2 - \sqrt{5})(x - 2 + \sqrt{5})x$												
Eigenvalues Weights of left eigenvectors	<table style="border-collapse: collapse; margin: auto;"> <tr> <td style="border-right: 1px solid black; padding: 2px 5px;">8</td> <td style="border-right: 1px solid black; padding: 2px 5px;">2 + $\sqrt{5}$</td> <td style="border-right: 1px solid black; padding: 2px 5px;">-1($\times 2$)</td> <td style="border-right: 1px solid black; padding: 2px 5px;">2 - $\sqrt{5}$</td> <td style="padding: 2px 5px;">0</td> </tr> <tr> <td style="border-right: 1px solid black; padding: 2px 5px;">0</td> <td style="border-right: 1px solid black; padding: 2px 5px;">$\frac{1}{2}(\frac{3}{\sqrt{5}} - 1)$</td> <td style="border-right: 1px solid black; padding: 2px 5px;">0, 0</td> <td style="border-right: 1px solid black; padding: 2px 5px;">$-\frac{1}{2}(\frac{3}{\sqrt{5}} + 1)$</td> <td style="padding: 2px 5px;">**</td> </tr> </table>	8	2 + $\sqrt{5}$	-1($\times 2$)	2 - $\sqrt{5}$	0	0	$\frac{1}{2}(\frac{3}{\sqrt{5}} - 1)$	0, 0	$-\frac{1}{2}(\frac{3}{\sqrt{5}} + 1)$	**		
8	2 + $\sqrt{5}$	-1($\times 2$)	2 - $\sqrt{5}$	0									
0	$\frac{1}{2}(\frac{3}{\sqrt{5}} - 1)$	0, 0	$-\frac{1}{2}(\frac{3}{\sqrt{5}} + 1)$	**									

The rotated odometer in this example has two ergodic invariant measures, one supported on the minimal set, another one Lebesgue. The minimal set corresponds to the symbols $\{0, 1, 2, 4\}$, all of which have positive weights. Thus the lift of (I_{\min}, F_π, μ) , where μ is the measure supported on the minimal set, is the skew-product $(I \times \mathbb{Z}, T_\pi, \mu \otimes \nu)$ which is transient, namely, $\overline{E}(\Psi|_{\mu \otimes \nu}) = \{0, \infty\}$, with 0 a trivial essential value. This means that the orbits of typical points with respect to $\mu \otimes \nu$ escape to ∞ without returning to 0. Here, as before, ν is the counting measure on \mathbb{Z} .

The situation is different for (I, F_π, Leb) . Since the rotated odometer is stationary, and Lebesgue measure is ergodic, Atkinson's result in Lemma 4.2 applies, and 0 is a non-trivial essential value of (I, F_π, Leb) . There are two eigenvalues outside of the unit circle, with one of left eigenvectors having non-zero weight. Thus Theorem 1.9 gives $\overline{E}(\Psi) = \{0, \infty\}$. This means that the orbits of typical points with respect to $\text{Leb} \otimes \nu$ 'oscillate' in $I \times \mathbb{Z}$, going further and further away from 0, but also returning to 0 infinitely many times.

Example 5.6.

$\pi = (0, 6, 5, 8, 4, 7, 3)$ $q = 9$ covering $\#\{\text{ergodic measures}\} = 2$ $\gamma_0 = 0.5515\dots$ $\overline{E}(\Psi) = \{0, \infty\} \cup ?$	$M = \begin{pmatrix} 2 & 1 & 0 & 1 & 0 & 0 & 2 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 & 2 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 & 2 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 5 & 7 & 10 & 10 & 4 & 8 & 3 & 12 & 8 & 8 \\ 2 & 2 & 3 & 1 & 1 & 0 & 2 & 2 & 4 & 4 \\ 2 & 2 & 3 & 1 & 1 & 0 & 2 & 2 & 4 & 4 \end{pmatrix}$																				
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	<table border="1"> <tbody> <tr> <td>0</td><td>1</td><td>2</td><td>3</td><td>4⁺</td><td>4⁻</td><td>5</td><td>6</td><td>7</td><td>8</td> </tr> <tr> <td>2</td><td>1</td><td>1</td><td>1</td><td>2</td><td>2</td><td>2</td><td>-3</td><td>-3</td><td>-3</td> </tr> </tbody> </table>	0	1	2	3	4 ⁺	4 ⁻	5	6	7	8	2	1	1	1	2	2	2	-3	-3	-3
0	1	2	3	4 ⁺	4 ⁻	5	6	7	8												
2	1	1	1	2	2	2	-3	-3	-3												
Characteristic polynomial	$(x - 16)(x - 4)(x - 1)(x^3 - 5x^2 + 2x - 1)x^4$																				
Eigenvalues	16 4.614* 4 1 0.19 - 0.42i* 0.19 + 0.42i* 0($\times 4$)																				
Weights of left eigenvectors	0 0.48* 1 0 0.26 - 0.36i* 0.26 + 0.36i* **																				

The rotated odometer in this example has two ergodic invariant measures, as we can see from the Frobenius form of M .

For the rotated odometer (I, F_π, Leb) , 0 is a non-trivial essential value by Lemma 4.2, so Lebesgue typical points with respect to $\text{Leb} \otimes \nu$ return infinitely often to 0. The substitution matrix M has three eigenvalues outside of the unit circle, two integer and one irrational (Pisot) eigenvalue. Left eigenvectors corresponding to two of these eigenvalues have non-zero weights, so Theorem 1.9 does not apply, but we still conjecture the absence of non-zero integer essential values.

Example 5.7.

$\pi = (0, 2, 7, 6, 5, 4, 3, 8, 10, 1, 9)$ $q = 11$ covering $\#\{\text{ergodic measures}\} = 2$ $\gamma_0 = 0.625\dots$ $\overline{E}(\Psi) \subseteq 2\mathbb{Z} \cup \{\infty\}$	$M = \begin{pmatrix} 2 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 2 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 0 & 0 & 0 & 0 & 2 & 0 & 0 \\ 3 & 6 & 6 & 6 & 4 & 4 & 4 & 8 & 4 & 3 & 8 & 8 \\ 3 & 6 & 6 & 6 & 4 & 4 & 4 & 8 & 4 & 3 & 8 & 8 \end{pmatrix}$																								
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	<table border="1" style="margin: auto;"> <tr> <td>0</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5⁺</td><td>5⁻</td><td>6</td><td>7</td><td>8</td><td>9</td><td>10</td> </tr> <tr> <td>2</td><td>4</td><td>2</td><td>2</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>2</td><td>-6</td><td>-6</td> </tr> </table>	0	1	2	3	4	5 ⁺	5 ⁻	6	7	8	9	10	2	4	2	2	0	0	0	0	0	2	-6	-6
0	1	2	3	4	5 ⁺	5 ⁻	6	7	8	9	10														
2	4	2	2	0	0	0	0	0	2	-6	-6														
Characteristic polynomial	$(x - 16)(x^3 - 8x^2 + 16x - 16)(x - 1)^2x^6$																								
Eigenvalue λ_j Weight $\psi(\ell_j)$	<table border="1" style="margin: auto;"> <tr> <td>16</td><td>5.679*</td><td>1.161 - 1.213i*</td><td>1.161 + 1.213i*</td><td>1($\times 2$)</td><td>0($\times 6$)</td> </tr> <tr> <td>0</td><td>0.34*</td><td>-0.17 + 0.20i*</td><td>-0.17 - 0.20i*</td><td>0, 0</td><td>**</td> </tr> </table>	16	5.679*	1.161 - 1.213i*	1.161 + 1.213i*	1($\times 2$)	0($\times 6$)	0	0.34*	-0.17 + 0.20i*	-0.17 - 0.20i*	0, 0	**												
16	5.679*	1.161 - 1.213i*	1.161 + 1.213i*	1($\times 2$)	0($\times 6$)																				
0	0.34*	-0.17 + 0.20i*	-0.17 - 0.20i*	0, 0	**																				

The rotated odometer in this example has two ergodic invariant measures, with one of the measures, μ , supported on the minimal set $I_{min} \subset I$, corresponding to the part of the matrix with symbols $\{0, 1, 2, 3, 4, 7, 8\}$. The weights of the substitution words corresponding to these symbols are all positive or zero, thus the skew-product $(I \times \mathbb{Z}, T_\pi, \mu \otimes \nu)$ is transient, namely, $\overline{E}(\Psi|_{\mu \otimes \nu}) = \{0, \infty\}$, with 0 a trivial essential value. The orbits of typical points with respect to $\mu \otimes \nu$ tend to ∞ without returning to 0.

For the rotated odometer (I, F_π, Leb) , 0 is a non-trivial essential value by Lemma 4.2, so $\text{Leb} \otimes \nu$ -typical points return infinitely often to 0. Non-zero weights of substitution words are even, which implies by Theorem 1.8 that $\overline{E}(\Psi) \subset 2\mathbb{Z} \cup \{\infty\}$, so 1 is not an essential value, and $\text{Leb} \otimes \nu$ is not ergodic. The second largest eigenvalue is not Pisot, so Theorem 1.9 does not apply, and we cannot rule out the existence of non-zero integer essential values.

Example 5.8.

$\pi = (0, 8, 7, 3, 2, 1)(4, 9)(5, 10, 6)$ $q = 11$ covering $\#\{\text{ergodic measures}\} = 3$ $\gamma_0 = 0.694\dots$ $\overline{E}(\Psi) = \{0\} \cup ?$	$M = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 2 & 2 & 3 & 1 & 2 & 1 & 1 & 2 & 2 & 3 & 0 & 1 \\ 1 & 2 & 1 & 1 & 2 & 0 & 1 & 1 & 1 & 2 & 0 & 1 \\ 2 & 3 & 2 & 1 & 2 & 1 & 1 & 3 & 1 & 2 & 0 & 2 \\ 4 & 4 & 8 & 13 & 9 & 1 & 1 & 8 & 11 & 8 & 16 & 11 \\ 1 & 2 & 2 & 0 & 1 & 1 & 1 & 2 & 1 & 1 & 0 & 1 \end{pmatrix}$																								
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	<table border="1" style="margin: auto;"> <tr> <td>0</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5⁺</td><td>5⁻</td><td>6</td><td>7</td><td>8</td><td>9</td><td>10</td> </tr> <tr> <td>2</td><td>2</td><td>2</td><td>2</td><td>1</td><td>1</td><td>1</td><td>2</td><td>1</td><td>2</td><td>-16</td><td>1</td> </tr> </table>	0	1	2	3	4	5 ⁺	5 ⁻	6	7	8	9	10	2	2	2	2	1	1	1	2	1	2	-16	1
0	1	2	3	4	5 ⁺	5 ⁻	6	7	8	9	10														
2	2	2	2	1	1	1	2	1	2	-16	1														
Characteristic polynomial	$(x - 16)(x^2 - 7x + 1)(x^2 + x + 1)(x^3 - 2x^2 - 3x - 1)x^4$																								
Eigenvalue λ_j Weight $\psi(\ell_j)$	<table border="1" style="margin: auto;"> <tr> <td>16</td><td>6.85*</td><td>3.08*</td><td>0.14*</td><td>$-0.5 \pm 0.86i^*$</td><td>$-0.54 \pm 0.18i$</td><td>0($\times 4$)</td> </tr> <tr> <td>0</td><td>1.4*</td><td>1.6*</td><td>3.6*</td><td>0, 0</td><td>$1.2 \mp 1.3i^*$</td><td>**</td> </tr> </table>	16	6.85*	3.08*	0.14*	$-0.5 \pm 0.86i^*$	$-0.54 \pm 0.18i$	0($\times 4$)	0	1.4*	1.6*	3.6*	0, 0	$1.2 \mp 1.3i^*$	**										
16	6.85*	3.08*	0.14*	$-0.5 \pm 0.86i^*$	$-0.54 \pm 0.18i$	0($\times 4$)																			
0	1.4*	1.6*	3.6*	0, 0	$1.2 \mp 1.3i^*$	**																			

The rotated odometer in this example has three ergodic invariant measures, which can be seen from the Frobenius form of M :

$$\left(\begin{array}{cccc|cccc|cccc|c} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 2 & 2 & 1 & 1 & 2 & 1 & 3 & 2 & 2 & 3 & 1 & 0 & 0 \\ 1 & 2 & 1 & 1 & 2 & 1 & 1 & 1 & 1 & 2 & 1 & 0 & 0 \\ 2 & 3 & 1 & 1 & 2 & 1 & 2 & 3 & 1 & 2 & 2 & 0 & 0 \\ 1 & 2 & 1 & 0 & 1 & 0 & 2 & 2 & 1 & 1 & 1 & 0 & 0 \\ \hline 4 & 4 & 1 & 1 & 9 & 13 & 8 & 8 & 11 & 8 & 11 & 16 & 16 \end{array} \right)$$

Denote by μ_1 and μ_2 the measures corresponding to the first and the second non-zero block, with leading eigenvalues $3.08\dots < 6.85\dots$, respectively. The weights of the symbols in these blocks are all positive, which means that $(I \times \mathbb{Z}, T_\pi, \mu_1 \otimes \nu)$ and $(I \times \mathbb{Z}, T_\pi, \mu_2 \otimes \nu)$ are both transient. Since the rotated odometer is covering, $(I \times \mathbb{Z}, T_\pi, \text{Leb} \otimes \nu)$ is recurrent by Lemma 4.2. However, $\gcd\{\psi(\chi(a)) : a \in \mathcal{A}\} = 1$ so Theorem 1.8 does not allow us to rule out any integer essential values. Since M has multiple eigenvalues with non-zero weights of left eigenvectors outside of the unit circle, Theorem 1.9 does not apply, and we cannot decide if $\text{Leb} \otimes \nu$ is ergodic or not.

5.4. Substitutions which are not covering.

Example 5.9.

$\pi = (0, 6, 5, 4, 3, 2, 1)$ $q = 7$ not covering $\#\{\text{ergodic measures}\} = \infty$ transient: $\gamma_0 = 1$ $\overline{E}(\Psi) = \{\infty\}$	$M = \begin{pmatrix} 2 & 2 & 0 & 0 & 1 & 1 & 0 & 2 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	$\begin{array}{c c c c c c c c} 0 & 1 & 2 & 3^+ & 3^- & 4 & 5 & 6 \\ \hline 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{array}$
Characteristic polynomial	$(x^2 - 2x - 6)x^6$
Eigenvalues Weights of left eigenvectors	$\begin{array}{c c c} 1 + \sqrt{7} & 1 - \sqrt{7} & 0(\times 4) \\ \hline \frac{1}{3}(4 - \sqrt{7}) & \frac{1}{3}(4 + \sqrt{7}) & ** \end{array}$

The rotated odometer in this example is not covering, which verified by noticing that the column sums in the substitution matrix are not $2^N = 8$, and which means that $I = I_{per} \cup I_{np}$, where I_{per} is a non-empty union of intervals of periodic points. Since the substitution is stationary, I_{per} is an infinite union of intervals, with infinite number of periods. Since $\psi(\chi(a)) = 1 > \psi(\chi(0)) = 0$ for all $a \neq 0 \in \mathcal{A}$, the z -extension $(I_{np} \times \mathbb{Z}, T_\pi)$ is transient.

Example 5.10.

$\pi = (04123), \quad q = 5$ not covering $\#\{\text{ergodic measures}\} = 2$	$M = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 \\ 2 & 3 & 2 & 1 & 4 & 4 \\ 2 & 3 & 2 & 1 & 4 & 4 \\ 2 & 3 & 2 & 1 & 4 & 4 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	$\begin{array}{c c c c c c} 0 & 1 & 2^+ & 2^- & 3 & 4 \\ \hline 2 & -2 & -2 & -2 & 1 & 1 \end{array}$
Characteristic polynomial	$(x - 6)(x - 2)x^4$
Eigenvalues Weights of left eigenvectors	$\begin{array}{c c c} 6 & 2 & 0(\times 4) \\ \hline \frac{1}{6} & \frac{1}{2} & ** \end{array}$

The weight of the leading left eigenvector is positive, which is possible since this substitution is not covering. The minimal part is a dyadic odometer; it lifts to a subsystem on the staircase that is also transient to $+\infty$, with diffusion coefficient 1.

5.5. A non-stationary example.

Example 5.11.

$\pi = (01243), \quad q = 5$ covering $\#\{\text{ergodic measures}\} = 2$ $\gamma_0 = \frac{1}{3}, \quad \overline{E}(\Psi) = \{0, \infty\}$	$M = \begin{pmatrix} 2 & 0 & 0 & 0 & 0 & 0 \\ 2 & 4 & 2 & 2 & 4 & 4 \\ 2 & 4 & 2 & 2 & 4 & 4 \\ 2 & 4 & 2 & 2 & 4 & 4 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$
$a \in \mathcal{A}$ Weight $\psi(\chi(a))$	$\begin{array}{c c c c c c} 0 & 1 & 2^+ & 2^- & 3 & 4 \\ \hline 2 & -2 & -2 & -2 & 1 & 1 \end{array}$
Characteristic polynomial	$(x - 8)(x - 2)x^4$
Eigenvalues Weights of left eigenvectors	$\begin{array}{c c c} 8 & 2 & 0(\times 4) \\ \hline 0 & -2 & ** \end{array}$

For $\pi = (01243)$, the renormalization of R_π gives $R_{\pi'}$ for $\pi' = (01234)$, see Example 5.4. In that example, Ψ is a coboundary, so every orbit of the \mathbb{Z} -extension is bounded. The composition with F_π changes this. For example, the minimal set of $F_{\pi'}$ in Example 5.4 is the dyadic odometer, the minimal subsystem corresponds to the subsystem $\chi'(0) = \chi'(3) = 03$. Because $\psi(\chi(0)) = 2$ and $\psi(\chi(3)) = 1$ for F_π , the \mathbb{Z} -extension of the minimal subsystem of F_π , which is also the dyadic odometer, becomes transient.

REFERENCES

- [1] B. Adamczewski, *Symbolic discrepancy and self-similar dynamics*. Ann. de l'Institut Fourier, **54** (2004), 2201–2234.
- [2] G. Atkinson, *Recurrence of co-cycles and random walks*. J. London Math. Soc. (2), **13** (1976), 486–488.
- [3] A. Avila, P. Hubert, *Recurrence for the wind-tree model*. Ann. Inst. H. Poincaré C Anal. Non Linéaire **37** (2020), 1–11.
- [4] V. Berthé, V. Delecroix, *Beyond substitutive dynamical systems: S-adic expansions*. RIMS Lecture note “Kokuroku Bessatu” B46 (2014), 81–123.
- [5] S. Bezuglyi, J. Kwiatkowski, K. Medynets, B. Solomyak, *Invariant measures on stationary Bratteli diagrams*, Ergod. Th. & Dynam. Sys., **30** (2010), 973–1007.
- [6] H. Bruin, O. Lukina, *Rotated odometers and actions on rooted trees*, Fund. Math. **260** (2023), 233–249.
- [7] H. Bruin, O. Lukina, *Rotated odometers*, J. Lond. Math. Soc., **107**(6) (2023), 1983–2024.
- [8] L. Bunimovich, Y. Sinai, *Statistical properties of Lorentz gas with periodic configuration of scatterers*. Comm. Math. Phys. **78:4** (1980/81), 479–497.
- [9] J. Chaika, D. Robertson, *Ergodicity of skew products over linearly recurrent IETs*. J. London Math. Soc. (2) **100** (2019), 223–248.
- [10] V. Delecroix, P. Hubert, S. Lelièvre, *Diffusion for the periodic wind-tree model*. Ann. Sci. Ec. Norm. Sup. **47:4** (2014), 1085–1110.
- [11] V. Delecroix, A. Zorich, *Cries and whispers in wind-tree forests*. Ann. of Math. Stud. **205** (2020), 83–115.
- [12] F. Durand, *Combinatorics on Bratteli diagrams and dynamical systems*, Combinatorics, Automata and Number Theory, Series Encyclopedia of Mathematics and its Applications **135**, Cambridge University Press (2010), 338–386.

- [13] K. Frączek, C. Ulcigrai, *Non-ergodic \mathbb{Z} -periodic billiards and infinite translation surfaces*. *Invent. Math.* **197** (2014), 241–298.
- [14] A. Garsia, *Arithmetic properties of Bernoulli convolutions*, *Trans. Amer. Math. Soc.* **102** (1962), 409–432.
- [15] W. Gottschalk, G. Hedlund, *Topological dynamics*, Amer. Mat. Soc. Colloquium **36** AMS, Providence RI (1955).
- [16] P. Hellekalek, *Ergodicity of a class of cylinder flows related to irregularities of distribution*, *Compositio Mathematica*, **61** (1987), 129–136.
- [17] P. W. Hooper, B. Weiss, *Generalized staircases: recurrence and symmetry*. *Ann. Inst. Fourier (Grenoble)* **62** (2012), no. 4, 1581–1600.
- [18] K. Frączek, P. Hubert, *Recurrence and non-ergodicity in generalized wind-tree models*. *Math. Nachr.* **291** (2018), 1686–1711.
- [19] A. Katok, B. Hasselblatt, *Introduction to the modern theory of dynamical systems*, Cambridge Univ. Press, Cambridge (1995).
- [20] L. Kuipers, H. Niederreiter, *Uniform distribution of sequences*, Pure Appl. Math. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, (1974), xiv+390 pp.
- [21] M. Lemanczyk, M. Mentzen, *Topological Ergodicity of Real Cocycles over Minimal Rotations*, *Monatsh. Math.* **134** (2002), 227–246.
- [22] M. Mentzen, *Some applications of groups of essential values of cocycles in topological dynamics*. *Topol. Methods Nonlinear Anal.* **23** (2004), no. 2, 357–375.
- [23] D. A. Pask, *Ergodicity of certain cylinder flows*, *Israel J. Math.* **76** (1991), 129–152.
- [24] A. Randecker, *Geometry and topology of wild translation surfaces*, PhD. thesis, Karlsruhe Institute of Technology, 2016.
- [25] K. Schmidt, *Cocycles on ergodic transformation groups*. Macmillan Lectures in Mathematics, **Vol. 1**. Macmillan Co. of India, Ltd., Delhi, 1977.
- [26] D. Szász, T. Varjú, *Limit laws and recurrence for the planar Lorentz process with infinite horizon*. *J. Stat. Phys.* **129** (2007), no. 1, 59–80.

HENK BRUIN, FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA

Email address: `henk.bruin@univie.ac.at`

OLGA LUKINA, MATHEMATICAL INSTITUTE, LEIDEN UNIVERSITY, P.O. BOX 9512, 2300 RA LEIDEN, THE NETHERLANDS

Email address: `o.lukina@math.leidenuniv.nl`